



## ANTIBACTERIAL ACTIVITIES OF PHYTOCHEMICAL EXTRACTS FROM THE LEAVES OF *JUSTICIA GENDARUSSA* BURM.F

**A.SIVASAKTHI\* AND VIJAYALAKSHMI M**

*Department of Biotechnology, Dr.M.G.R. Educational and Research Institute university,  
Maduravoyal, Chennai,Tamilnadu, India.*

### ABSTRACT

Phytochemicals are chemical compounds that occur naturally in plants and protect plants against bacteria, viruses, and fungi. They are non-nutritive plant chemicals that have protective and disease preventive properties. The aim of the present study was to investigate the antibacterial activities of phytochemical extracts from the leaves of *Justicia gendarussa*. The antibacterial activities of *Justicia gendarussa* phytochemical extracts were determined by agar disc diffusion and minimum inhibitory concentration (MIC) against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Vibrio cholerae*. The highest zone of inhibitory activity was observed with alkaloid extract of the plant against the test bacteria while less antibacterial activity was observed in terpenoid, flavanoid and glycoside extracts. Therefore, it was confirmed that alkaloid extract of *Justicia gendarussa* contains a high level of antibactericidal compound which can be used as a novel chemotherapeutic drug for bacterial diseases.

**KEYWORDS:** *Justicia gendarussa*, Antibacterial activities, Alkaloid extract and novel chemotherapeutic drug.



**A.SIVASAKTHI**

Department of Biotechnology, Dr.M.G.R. Educational and Research Institute university,  
Maduravoyal, Chennai,Tamilnadu, India.

## INTRODUCTION

India is an emporium of medicinal plants. It is one of the richest countries in the world with regard to its genetic resources of medicinal plants. In recent years, the secondary plant metabolites (phytochemicals), previously with unknown pharmacological activities, have been extensively investigated as a source of medicinal agents<sup>1</sup>. Infectious diseases are the leading cause of death world-wide. Antibiotic resistance has become a global concern<sup>2</sup>. The clinical efficacy of many existing antibiotics is being threatened by the emergence of multidrug-resistant pathogens<sup>3</sup>. There is a continuous need to discover new antimicrobial compounds with diverse chemical structures and novel mechanisms of action<sup>4</sup>. Therefore, researchers are increasingly turning their attention to folk medicine, looking for new leads to develop better drugs against microbial infections<sup>5</sup>. *Justicia gendarussa* belongs to the family *Acanthaceae*. The family *Acanthaceae* is a taxon of dicotyledonous flowering plants containing almost 240 genera and 2200 species<sup>6</sup>. Most are tropical herbs, shrubs or twining vines, some are epiphytes. Only a few species are distributed in temperate region. It is commonly called as 'Neernotchi' in Tamil. Leaves are linear lanceolate, glabrous; flower small, white with pink or purple spots inside. The plant is used in traditional medicinal practice for chronic rheumatism, inflammations, bronchitis, vaginal discharges, dyspepsia, eye diseases, fever, hemiplegia, headache, earache, muscle pain, respiratory disorder and digestive trouble. The roots of this plant are also used to treat cough, jaundice, arthritis, cephalgia, facial paralysis, otalgia, hemicranias and liver disorders<sup>7</sup>. The dried leaves are used to repel insects from clothing.

## MATERIALS AND METHODS

### *Plant material*

*Justicia gendarussa* leaves were collected from Medavakkam near Tambaram, Tamilnadu, India. The plant was identified and authenticated by Dr. S.Sankaranarayanan,

Head of the department, Department of Medicinal Botany, Sri Sairam Siddha Medical College, Tambaram, Chennai. The plant material was air-dried under shade at room temperature, ground with an electric grinder into fine powder and stored in airtight containers for analysis.

### *Bacterial strains*

The bacterial strains used were Gram positive: *Staphylococcus aureus* MTCC 29213; and Gram negative: *Escherichia coli* MTCC 25922, *Proteus mirabilis* MTCC 13315 and *Vibrio cholerae* MTCC 12657. All bacterial strains were obtained from Microbial Type Culture Collection and Gene Bank, Institute of Microbial Technology Sector 39-A, Chandigarh – 160036, India.

### *Phytochemical analysis of the plant extract*

The screening of plant secondary metabolites such as alkaloids, terpenoids, flavonoids, glycosides, anthroquinones, tannins and saponnins was performed using the standard procedure given by Trease and Evans<sup>8</sup> and Harborne<sup>9</sup>.

### *Extraction of phytochemicals from the leaves of J.gendarussa*

#### *Extraction of Alkaloid*

Finely powdered leaf material was extracted with cold distilled methanol with occasional swirling. After filtration, the solvent was removed under reduced pressure at 40 °C, to minimise any thermal degradation of the alkaloids. The crude alkaloid mixture was then separated from neutral and acidic materials, and from water solubles, by initial extraction with aqueous acetic acid followed by dichloromethane. The basification was done on the aqueous solution and the organic layer of dichloromethane contained crude alkaloid extract<sup>10</sup>.

**Extraction of Terpenoids**

Ground leaves were extracted with hot (60 °C) 95% ethanol. After filtration, the dark green solvent was evaporated to dryness under reduced pressure at 40 °C. The residue was partitioned between water and chloroform. The organic layer was separated and condensed to yield dark green syrup. The chloroform extract was then partitioned between hexane and 10% aqueous methanol. The aqueous methanol extract was then used for antibacterial activity as terpenoid extract<sup>11</sup>.

**Extraction of Flavonoids**

The dried leaf powder of *J.gendarussa* was defatted with petroleum ether (40-60 °C). The extract was then percolated with methanol until exhaustion at 40 °C by rotary evaporator. The condensed material was partitioned using ethyl acetate. The ethyl acetate extract contains crude flavonoids<sup>12</sup>.

**Extraction of Glycosides**

The powdered leaf material of the plant was extracted three times with methanol at 25 °C for 24 hours and then concentrated in vacuum. The extract was washed with n-hexane and then the methanol layer was further concentrated to a gummy mass. The later was suspended with water and extracted with equal volume of ethyl acetate to obtain glycoside extract of the plant<sup>13</sup>.

**Antibacterial activities of four different phytochemical extracts of *J.gendarussa* tested against pathogenic bacteria****Agar disc diffusion assay**

The antibacterial activity was studied using the disc-diffusion method<sup>14</sup>. Bacteria were grown overnight on Muller Hinton (MH) agar plates. Five young colonies were suspended with 5ml of sterile saline (0.9%) and the density of the suspension adjusted to approximately  $3 \times 10^8$  colony forming units (CFU). The swab was used to inoculate the dried surface of MH agar plate by streaking four times over the surface of the agar, rotating the plate

approximately by 90 ° to ensure an even distribution of the inoculums. The medium was allowed to dry for about 3 min before adding a sterile paper disc of 5 mm diameter. Each disc was tapped gently down onto the agar to provide uniform contact. Phytochemical extracts (50µg) were weighed and dissolved in 1ml of 7% Methanol. 5, 10, 15 and 20 microlitres of the compounds were introduced on each disc (five replicates) and 7% methanol alone served as a negative control. The plates were incubated at 37 °C for 24 h; inhibition zones were measured and calculated.

**Minimum inhibitory concentration (MIC)**

The MIC of the isolated compounds was determined by dilution method<sup>15</sup>. The strains were grown in MH broth to exponential phase with an A560 of 0.8, representing  $3 \times 10^8$  CFU/ml. Different dilutions of the *J.gendarussa* phytochemical extracts were prepared to give concentrations at 5, 10, 15 and 20 µg/ml respectively. A 0.5 ml of each concentration was added into separate test tubes containing 4ml of MH broth inoculated with 0.5 ml bacterial suspension at a final concentration of  $10^8$  CFU/ml. Each MIC was determined from five independent experiments performed in duplicate. The tubes containing 4.5 ml of bacterial inoculates and 0.5 ml of 7% methanol were used as bacterial controls, 4.5 ml of uninoculated MH broth and 0.5 ml plain broth solution served as a blank. The tubes were incubated at 37 °C for 18 h; inhibition of bacterial growth was determined by measuring the absorbance at A560 nm.

**RESULTS AND DISCUSSION**

The phytochemical screening of the *J.gendarussa* showed the presence of alkaloids, terpenoids, flavanoids, glycosides, tannins and saponins (Table-1). The presence of starch, glycosides, saponins, tannins, phenolic compounds, terpenoids, steroids and flavonoids in *J.gendarussa* was already recorded.<sup>16</sup>

**Table 1**  
**Phytochemical screening of leaf extract of *J.gendarussa***

SL NO.	PHYTO CONSTITUENTS	TEST PERFORMED	OBSERVATION	PRESENT/ABSENT
1	Alkaloids	Dragendorff's test	Orange / red precipitate	++
		Mayers test	Yellowish precipitation	
2	Flavonoids	Alkali reagent test	Intense yellow colour	++
3	Glycosides	Keller-killani test	Formation of brown ring interface	++
4	Tannin	FeCl <sub>3</sub> test	Brownish green coloration	++
5	Saponins	Frothing test	Foam	++
6	Terpenoids	Salkowski test	Reddish brown colouration of the interface	++
7	Antraquinones	Benzene Ammonia Test	Pink colour	--

-- = Negative (absent); ++= Positive (present)

**Table 2**  
**Antibacterial activities of phytochemical extracts of *J.gendarussa* against bacterial species [Diameter of the inhibition zone (mm) ]**

Type of extract	<i>Staphylococcus aureus</i>				<i>Vibrio cholerae</i>			
	5µg/ml	10µg/ml	15µg/ml	20µg/ml	5µg/ml	10µg/ml	15µg/ml	20µg/ml
Alkaloid extract	12.33±0.58	13.9±0.5	14±0	15.33±0.58	9±0	9.33±0.58	10.58±0.41	11.9±0.58
Terpenoid extract	9±0	9.33±0.58	10.03±0.37	11.6±0.55	8.3±0.81	9.3±0.40	10.03±0.37	11.6±0.55
Flavonoid extract	8.3±0.81	9±0	10.9±0.28	11.5±0.5	7.3±0.58	8.38±0.81	9.33±0.58	9.97±0.37
Glycoside extract	7±0.5	8.1±0.5	9.33±0.58	10.4±0.61	6.9±0.65	8.5±0.40	8.9±0.9	9.32±0.27

Type of extract	<i>Escherichia coli</i>				<i>P.aeruginosa</i>			
	5µg/ml	10µg/ml	15µg/ml	20µg/ml	5µg/ml	10µg/ml	15µg/ml	20µg/ml
Alkaloid extract	10.33±0.58	11.9±0.5	12±0	13.33±0.58	8.03±0.55	9.6±0.32	10.2±0.60	11.06±0.60
Terpenoid extract	7.5±0.81	8.3±1.17	9.1±0.28	10.9±0.36	8±0	9.31±0.58	10.03±0.37	10.6±0.55
Flavonoid extract	6.6±0.57	7.5±0.86	8.8±0.76	9.5±0.5	5.7±0.64	6.6±0.79	7.4±0.40	8.05±0.45
Glycoside extract	5.7±0.64	6.6±0.79	7.4±0.40	8.05±0.45	5.6±0.57	6.5±0.86	6.83±0.76	7.58±0.5

\*The antimicrobial activity was determined by measuring the diameter of zone of inhibition that is the mean of triplicates ± SD of three replicates.

Phytochemical extracts thus obtained were further subjected for determination of minimal inhibitory concentration by two-fold micro broth dilution method against the bacteria studied. Table-3 indicates that the alkaloid extract was found to be most significant inhibitor than the other extracts. MIC of this extract showed gradient value against the concentration used to inhibit the bacteria. Furthermore, gram positive bacterial species was found most sensitive as compared to gram negative species.

**Table 3**  
**Minimal Inhibitory Concentration (MIC) of different phytochemical extracts against bacteria [Optical density values at (560nm) ]**

Type of extract	<i>Staphylococcus aureus</i>				<i>Vibrio cholerae</i>			
	5µg/ml	10µg/ml	15µg/ml	20µg/ml	5µg/ml	10µg/ml	15µg/ml	20µg/ml
Alkaloid extract	0.326±0.05	0.225±0.01	0.142±0.01	0.106±0.03	0.514±0.02	0.425±0.01	0.315±0.01	0.213±0.01
Terpenoid extract	0.544±0.02	0.425±0.01	0.315±0.01	0.213±0.01	0.579±0.05	0.402±0.07	0.396±0.01	0.233±0.04
Flavonoid extract	0.579±0.05	0.462±0.07	0.296±0.01	0.233±0.04	0.660±0.04	0.569±0.09	0.469±0.04	0.409±0.06
Glycoside extract	0.678±0.05	0.462±0.07	0.396±0.01	0.233±0.04	0.647±0.03	0.557±0.02	0.424±0.01	0.332±0.01

Type of extract	<i>Escherichia coli</i>				<i>P.aeruginosa</i>			
	5µg/ml	10µg/ml	15µg/ml	20µg/ml	5µg/ml	10µg/ml	15µg/ml	20µg/ml
Alkaloid extract	0.559±0.05	0.462±0.07	0.396±0.01	0.233±0.04	0.524±0.02	0.425±0.01	0.335±0.01	0.213±0.01
Terpenoid extract	0.614±0.02	0.425±0.01	0.315±0.01	0.313±0.01	0.679±0.05	0.402±0.07	0.396±0.01	0.233±0.04
Flavonoid extract	0.660±0.04	0.569±0.09	0.469±0.04	0.409±0.06	0.660±0.04	0.569±0.09	0.469±0.04	0.409±0.06
Glycoside extract	0.647±0.03	0.557±0.02	0.424±0.01	0.332±0.01	0.747±0.03	0.557±0.02	0.424±0.01	0.332±0.01

\*The Minimal Inhibitory Concentration was determined by measuring the turbidity of the bacterial culture that is the mean of triplicates ± SD of three replicates.

## CONCLUSION

The present study clearly proved that the alkaloid compound of *J.gendarussa* posses greater antibacterial activity on both gram positive and gram negative bacteria. Development or synthesis of novel antibacterial drugs from this medicinal plant for treating bacterial diseases may be further researched for the benefit of mankind.

## ACKNOWLEDGMENT

The authors gratitude to Gloris Biomed Research Centre (P) Ltd for the support and encouragement for this work.

## REFERENCES

1. Krishnaraju AV, Rao TVN, Sundararaju D, Assessment of bioactivity of Indian medicinal plants using Brine shrimp (*Artemia salina*) lethality assay. Int J Appl Sci Eng 2: 125-134, 2005.
2. Westh H, Zinn CS, Rosdahl VT, An international multicenter study of antimicrobial consumption and resistance in *Staphylococcus aureus* isolates from 15 hospitals in 14 countries, Microb Drug Resist 10: 169-176, 2004.
3. Bandow JE, Brotz H, Leichert LIO, Proteomic approach to understanding antibiotic action. Antimicrob Agents Chemother 47: 948-955, 2003.
4. Rojas R, Bustamante B, Bauer J, Antimicrobial activity of selected Peruvian medicinal plants. J Ethnopharmacol 88: 199-204, 2003.
5. Benkeblia N. Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium*

- sativum). *Lebensm-Wiss u-Technol* 37: 263-268, 2004.
6. Pandey B.P, College Botany, Volume II, S.CHAND publishing, New Delhi, page no 290, 2010
  7. Sankaranarayanan S. Medical Taxonomy of Angiosperms: Recent trends in Medicinal uses and Chemical Constituents. Harishi Publications, Chennai-73, 2009.
  8. Trease GE, Evans WC: Pharmacognosy 11th edn, Brailliar Tiridel Can. Macmillian publishers, 1989.
  10. Harborne JB. Phytochemical methods, London. Chapman and Hall, Ltd. pp. 49-188. 1973.
  11. Surya Hadi and John BB. Initial Studies on Alkaloids from Lombok Medicinal Plants. *Molecules* 6: 117-129, (2001)
  12. Paul PB, Po-ming hon HC, Dominic Chan TW, Bo-mu W, Thomas CWM. and Chun-Tao C. Sesquiterpene lactones from *Elephantopus scaber*. *Phytochemistry*; 44: 113-116(1997).
  13. Amal MYM, Ahmed IK, Mahmoud AS. Isolation, structural elucidation of flavonoid constituents from *Leptadenia pyrotechnica* and evaluation of their toxicity and antitumor activity. *Pharmaceutical Biology* 2009; 47: 539–552.
  14. Aya H, Katsuyoshi M, Hideaki O. and Yoshio T. Flavonol glycosides from the leaves of *Indigofera zollingeriana*. *J. Nat Med.* (2011); 65: 360.
  15. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *American Journal of Clinical Pathology* (1996); 45: 493–496.
  15. Brantner A, Grein E. Antibacterial activity of plant extracts used externally in traditional medicine. *Journal of Ethnopharmacology* (1994); 44: 35–40.
  16. Subramanian N et.al Antimicrobial activity and preliminary phytochemical screening of *justicia gendarussa* (burm. F.) Against human pathogens