



**ANTIOXIDANT ACTIVITY OF LEAF, UNRIPE AND
RIPE FRUITS OF *LANTANA CAMARA* L**

S LAYAVARJITHA^{*1} AND D KADAMBAN²

1. M.Phil. Scholar Department of Botany, Kanchi Mamunivar Centre For Post Graduate Studies, Puducherry, India.
2. Associate Professor of Botany and Research Supervisor, Department of Botany, Kanchi Mamunivar Centre For Post Graduate Studies, Puducherry, India.

ABSTRACT

The antioxidant activity of leaf, unripe and ripe fruits of *Lantana camara* L. were determined using H₂O₂ radical scavenging activity and Ferric Reducing antioxidant power (FRAP) assay. Ethanol, methanol, ethyl acetate and water extracts prepared from dry leaf powder and fresh fruits of both unripe and ripe fruits were used to find out the antioxidant activity. All the four extracts of leaf, unripe and ripe fruits showed considerable antioxidant effect. Water extract of powdered leaf showed a maximum activity of 94.81% where as aqueous extract of unripe fruit showed the least of 0.05% in FRAP activity. In H₂O₂ radical scavenging activity water extract of unripe showed the highest activity of 14.53% and methanol, ethyl acetate and water extracts of leaf powder showed the least activity of 3.05%. As the leaf, unripe and ripe fruits of *Lantana camara* have antioxidant potential they can be studied from pharmaceutical and nutraceutical aspects.

KEY WORDS: Antioxidant activity, FRAP, H₂O₂, *Lantana camara*



S LAYAVARJITHA

M.Phil. Scholar Department of Botany, Kanchi Mamunivar
Centre For Post Graduate Studies, Puducherry, India.

INTRODUCTION

The Verbenaceae species *Lantana camara* L. is a native to tropical and subtropical America and has been dispersed throughout the world as a popular ornamental plant, becoming one of the World's worst weeds¹. However, it is listed as one of the important medicinal plants of the world and it has been used in many parts of the world to treat a wide variety of disorders². In India the leaves of the plant are boiled for tea and the decoction is a remedy against cough and it is used as a lotion for wounds and Pounded leaves are applied to cuts, ulcers and swellings³. Antioxidants are compounds that protect cells against the damaging effects of reactive oxygen species. They are phytochemicals, vitamins and other nutrients that protect our cells from damage caused by free radicals. The principal agents responsible for the protective effects could be the presence of antioxidant substances that exhibit their effects as free radical scavengers, hydrogen-donating compounds, singlet oxygen quenchers and metal ion chelators⁴. The majority of the antioxidant activity is due to the flavones, isoflavones, flavonoids, anthocyanin, coumarin, lignans, catechins and isocatechins⁵. Antioxidant-based drug formulations are used for the prevention and treatment of complex diseases like atherosclerosis, stroke, diabetes, Alzheimer's disease and cancer⁶.

Some of the reactive oxygen species, including hydrogen peroxide, singlet oxygen, hydroxyl and superoxide radicals, have positive roles in energy production *in vivo* systems, phagocytosis, intercellular signal transfer, regulation of cell growth and the synthesis of important biological compounds⁷. Additionally, reactive oxygen species modify DNA and membranes by attacking the lipids, proteins, and carbohydrates in cell membranes and tissues⁸. In the organism, the rates of production and removal of free radicals are in balance, known as oxidative balance. An increase in the rate of production or a decrease in the rate of removal disrupts this balance and increases the levels of reactive oxygen species. This condition, which is called oxidative stress,

indicates a serious imbalance between the production of free radicals and the antioxidant defense systems, resulting in tissue damage⁹. Antioxidants are compounds that protect cells against the damaging effects of reactive oxygen species.

L. camara contains lantadenes, the pentacyclic triterpenes which is reported to possess number of useful biological activities. Several previous reports have described antifungal¹⁰, antiproliferative¹¹, and antimicrobial activities of *L. camara*² including termicidal activities¹². Ethanolic extract of dried flowers of *Lantana camara* were screened for its protective effect against the Cisplatin-induced kidney damage in rats¹³. The antioxidant activity of methanolic extract of *Lantana camara* has been already studied using DPPH radical scavenging activity and nitric oxide free radical scavenging method¹⁴. The aim of present study is to determine the percentage of free radical scavenging activities of antioxidants using H₂O₂ radical scavenging and FRAP assay of leaf, unripe and ripe fruits of the plant, *Lantana camara*, in four different extracts viz. ethanol, methanol, ethyl-acetate and water .

MATERIALS AND METHODS

Plant Material

The aerial parts of *Lantana camara* L. were collected in March 2013 in Lawspet, Puducherry and identified at the French Institute, Puducherry for its correct identity. Voucher Specimen was deposited in the Herbarium of the Department of Plant science and Biotechnology, Kanchi Mamunivar centre for Post Graduate Studies, Puducherry.

Preparation of plant extracts

The leaves of the plants were air dried at room temperature for 10 days then powdered using a mixer grinder. Plant extracts were prepared for all the parts of the plant (ethanol, methanol, ethyl-acetate and water) following a standard protocol.

HYDROGEN PEROXIDE SCAVENGING EFFECT

The scavenging activity of hydrogen peroxide by the plant extracts was determined by the method of Ruch et al. (1989)¹⁵.

Principle

The UV light absorption of hydrogen peroxide can be easily measured at 230nm. On scavenging of hydrogen peroxide by the plant extracts, the absorption decreases at this wavelength, which property can be utilized to quantify their H₂O₂ scavenging ability.

Reagents

1. Phosphate buffer (40mM, pH 7.4),
2. H₂O₂ in phosphate buffer (40mM)

Procedure

A solution of hydrogen peroxide (40 mM) was prepared in phosphate buffer (pH 7.4). Plant extracts at the concentration of 10mg/10µl were added to 0.6ml of H₂O₂ solution. The total volume was made up to 3ml with phosphate buffer. The absorbance of the reaction mixture was recorded at 230nm. The blank solution contained phosphate buffer without H₂O₂. The percentage of H₂O₂ scavenging by the plant extracts was calculated as

$$\% \text{ scavenged hydrogen Peroxide} = \frac{A_0 - A_1}{A_0} \times 100$$

Where,

A₀ - Absorbance of control

A₁ - Absorbance in the presence of plant extract


REDUCING POWER ASSAY –FRAP

The reducing power of different fractions was determined by the method of Oyaizu (1986)¹⁶.

Principle

The reducing power assay was used to assess the reduction potential of fractions as this assay involves the reduction of ferricyanide ion i.e. [Fe (CN)₆]³⁻ to ferrocyanide ion i.e. [Fe (CN)₆]⁴⁻ by electron donation from polyphenols . The

polyphenols are oxidized in this reaction. The ferrocyanide ions combines with Fe (III) in acidic medium to give a Prussian blue complex i.e. ferri ferrocyanide complex, Fe₄ [Fe (CN)₆]³⁻, the intensity of which is measured spectrophotometrically at 700 nm (Graham, 1992). The intensity of coloured complex increases with the electron or H. donating ability of extract and its different fractions. The redox reaction may be summarized as follows:



Procedure

1 ml of extract of different concentrations was mixed with 2.5 ml of phosphate buffer (200mM, pH 6.6) and 2.5 ml of 1% potassium ferricyanide. The mixture was incubated at 50°C for 20 minutes. A volume of 2.5 ml of 10% TCA was then added to the mixture and centrifuged at 3000 rpm for 10 minutes. 2.5 ml of supernatant was mixed with 2.5 ml of distilled water and 0.5

ml of FeCl₃ (0.1%) and the absorbance was measured spectrophotometrically at 700nm. Increase in absorbance of the reaction mixture was interpreted as increase in reducing activity of the extract and the results were compared with Gallic acid that was used as a positive control. The percentage reduction of the sample as compared to standard was calculated using the formula:

$$\text{Reduction (\%)} = 1 - \frac{\text{Absorbance of Gallic Acid} - \text{Absorbance of Sample}}{\text{Absorbance of Gallic Acid}} \times 100$$

Here, absorbance of Gallic acid refers to the absorbance value exhibited by Gallic acid at the highest tested concentration i.e. 200µg / ml.

RESULTS AND DISCUSSION

All the four extracts viz. ethanol, methanol, ethyl-acetate and water of leaf, unripe and ripe fruits of the plant, *Lantana camara* shows antioxidant activity. Percentage of antioxidant activity of different extracts of *Lantana camara* leaf, raw and ripe fruit are given in Table 1 and 2. Percentage of antioxidant activity of different extracts of *Lantana camara* leaf, raw and ripe fruits are given in Fig 1 and 2.

H₂O₂ scavenging activity

The antioxidant activities of H₂O₂ scavenging ability of ethanol, methanol, ethyl-acetate and water extracts are reported in Table-1 and Fig-1. The antioxidant activity of unripe fruit decreased in the order of water extract (14.53%)> methanol extract (10.83%)> ethyl acetate extract (4.28%)> ethanol extract (2.79%). In ripe fruit the antioxidant activity decreased in the order of ethanol extract (6.81%)> methanol extract (4.67%)> ethyl acetate extract (3.24%)> water extract (2.79%). In the case of leaf powder the

order of decrease in the antioxidant activity is ethanol extract (4.54%)> methanol extract, ethyl acetate extract, water extract (3.05%). Water extract of unripe showed the highest activity of 14.53% and water extracts of ripe fruit and ethanol extract of unripe fruit showed the least activity of 2.79%.

Antioxidant activity by using FRAP method

The results of the FRAP assay are reported in Table 2 and Fig.2. All the four extracts of leaf, unripe and ripe fruits of *Lantana camara* showed considerable antioxidant activity. The antioxidant activity of unripe fruit decreased in the order of ethanol extract (56.78%)> ethyl acetate extract (20.16%)> methanol extract (14.94%)> water extract (0.05%). In ripe fruit the antioxidant activity decreased in the order of methanol extract (60.70%)> water extract (53.32%)> ethanol extract (25.92%)> ethyl acetate extract (23.29%). In the case of leaf powder the order of decrease in the antioxidant activity is water extract (94.81%)> ethanol extract (71.22%)> methanol extract (65.59%)> ethyl acetate extract (60.38%). Water extract of powdered leaf showed a maximum activity of 94.81% where as aqueous extract of unripe fruit showed the least of 0.05% in FRAP activity.

Percentage of antioxidant activity of different extracts of Lantana camara leaf, unripe and ripe fruits.



TABLE 1
H₂O₂ scavenging activity

SAMPLE	SOLVENTS			
	ETHANOL	METHANOL	ETHYL ACETATE	WATER
Unripe fruit	2.79	10.83	4.28	14.53
Ripe fruit	6.81	4.67	3.24	2.79
Leaf powder	4.54	3.05	3.05	3.05

TABLE 2
FRAP Method

SAMPLE	SOLVENTS			
	ETHANOL	METHANOL	ETHYL ACETATE	WATER
Unripe fruit	56.78	14.94	20.16	0.05
Ripe fruit	25.92	60.70	23.29	53.32
Leaf powder	71.22	65.59	60.38	94.81

Figure 1

H₂O₂ scavenging activity of various extracts of parts of Lantana camara





Figure 2

Antioxidant activity by using FRAP method of various parts of Lantana camara in various solvents



Antioxidant-based drug formulations are used for the prevention and treatment of complex diseases like atherosclerosis, stroke, diabetes, Alzheimer's disease and cancer⁶. There is an increasing interest in antioxidants, particularly in those intended to prevent the presumed deleterious effects of free radicals in the human body, as well as the deterioration of fats and other constituents of food stuffs¹⁷. Recently

there has been an upsurge of interest in the therapeutic potential of medicinal plants as antioxidants in reducing such free radical induced tissue injury. Antioxidants thus play an important role of protecting the human body against damage by reactive oxygen species¹⁸. Based on the research finding leaf, unripe and ripe fruits of *Lantana camara* can be studied from pharmaceutical and nutraceutical aspects.

REFERENCES

- Day MD, Willey CJ, Playford J and Zalucki MP. *Lantana: Current Management Status and Future Prospects*; Australian Centre for International Agricultural Research: Canberra, Australian, (2003).
- Ross IA. Chemical constituents, traditional and modern medical uses. *Medicinal plants of the world*. Humana Press, New Jersey. (1999).
- Verma RK and Verma SK. Phytochemical and termiticidal studies of *Lantana camara var aculeata* leaves. *Fitoterapia* 77: 466-468. (2006).
- Okawa M, Kinjo J, Nohara T and Ono M. DPPH (1,1-Diphenyl-2-Picrylhydrazyl) radical
- Scavenging activity of flavonoids obtained from some medicinal plants. *Boil Pharm Bull* 24: 1202-1205. (2001).
- Aqil F, Ahmed I and Mehmood Z. Antioxidant and free radical scavenging properties of twelve traditionally used Indian medicinal plants. *Turk J Biol* 30: 177-183, (2006).
- Devasagayam TPA, Tilak JC, Boloor KK, et al. Review: Free radical and antioxidants in human health. *Curr Stat Fut Pros JAPI* 53: 794-804, (2004).
- Packer L, Cadenas E, and Davies KJA. Free radicals and exercise: An introduction. *Free Radical Biology and Medicine*, 44, pp. 123-125. (2008).
- Jung T, Höhn A, Catalgol B, and Grune T. Age-related differences in oxidative protein-damage in young and senescent fibroblasts. *Archives of Biochemistry and Biophysics*, 483, pp. 127-135. (2009).
- Wells PG, McCallum GP, Chen CS, Henderson JT, Lee CJ, Perstin J, Preston TJ, Wiley MJ, and Wong AW. Oxidative stress in developmental origins of disease: Teratogenesis, neurodevelopmental deficits, and cancer. *Toxicological Sciences*, 108, pp. 4-18. (2009).
- WHO expert committee on Diabetes mellitus, Technical reports series World Health Organisation. Geneva: (1980).
- Sharma OP, Dawra RK and Makkar HPS. Isolation and partial purification of *Lantana (Lantana camara L.)* toxins. *Toxicology Letters*. 37:165-172. (1987).
- Tripathi AK and Shukla BN. Antifungal activity of some plant extracts against *Fusarium oxysporum* sp. causing wilt of linseed. *Journal of Mycology and Plant Pathology*. 32:266-7. (2002).
- Khalid S, Khan H, Anwar AA and Abrar S. Research Article. Nephroprotective effect of the ethanolic extract of *Lantana camara Linn* flower on acute dose of Cisplatin induced renal injured rats. *RGUHS J Pharm Sci* . Vol 2, Issue 2, pg 68-77. (2012).
- Mayee R and Thosar A. Evaluation of *Lantana Camara* for antiurolithiatic and antioxidant activities in rats. *Pharm. Clinical. Res.*, 3: 10-14. (2011).
- Ruch RJ, Cheng S, and Klaunig JE. Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea, *Carcinogenesis*, 10, 1003-1008. (1989).
- Oyaizu M. Studies on product of browning reaction prepared from glucose amine. *Japanese Journal of Nutrition* 44: 307-15. (1986).
- Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin J Sci Technol* 26: 211-219. (2004).
- Gopalakrishnan G, Dhanapal CK, and Manavalan R. *In-vitro* antioxidant activities of methanolic extract of root of *coleous vettiveroides* (Jacob), *International Journal of Pharma and Bio Sciences*: Vol.2, Issue-4.P-353 -7, (2011).