



---

## MULTIFARIOUS ACTIONS OF DIETARY FLAVONOIDS – IMPLICATIONS IN CANCER AND CATARACT

**DURGA.M, NATHIYA.S AND DEVASENA.T\***

*Centre for Nanoscience and Technology, A.C.Tech Campus,  
Anna University, Chennai-600025, India.*

### ABSTRACT

Flavonoids are widely present in fruits and vegetables. Different categories of flavonoids such as flavonals, flavones, flavonones, isoflavone, catechin, anthocyanin and isoflavone are present in plants. It has been discovered that they have many beneficial actions on body cells by enhancing the activity of many enzyme systems. The aim of this present review is to summarize the significant multifunctional and biological activities of flavonoids such as anti-allergic, anti-toxic, anti-microbial, anti-cataract and anti-cancer activity and to present the recent advancements in the pharmacological and chemical properties of flavonoids.

**KEYWORDS:** Flavonoids, anti-cancer, quercetin, cataract and anti-microbial.



**DEVASENA.T**

Centre for Nanoscience and Technology, A.C.Tech Campus,  
Anna University, Chennai-600025, India.

tdevasenabio@sify.com

tdevasenabio@annauniv.edu

## INTRODUCTION

Diverse beneficial effects of plants have been related to the presence of phytochemicals such as polysaccharides and flavonoids<sup>1</sup>. Flavonoids are bioactive polyphenols of low molecular weight<sup>2,3</sup> that play an important role in photosynthetic plant cells<sup>4</sup>. Flavonoids are identified by their flavan nucleus<sup>5</sup> and carbon skeleton<sup>6,7</sup>. Until nearly 50 years ago, the mechanism of flavonoid action was not clear. Research in flavonoids began in early 1930s; Albert Szent-Gyorgi a Hungarian scientist, discovered a new substance from oranges. It was first termed as "Vitamin P". Later, it was clarified that this substance was "Rutin" (a Flavonoid), since then a great interest in flavonoids was evidenced. They possess a

phenyl benzopyrene nucleus (C6-C3-C6) and are classified into different groups such as flavanols, flavones, isoflavones, flavones, and flavanonols based on their chemical structures<sup>8,9</sup> (Table 1). It was discovered that the normal human intake of flavonoids is a few 100s of mg/day<sup>10</sup>. These compounds are widely present in medicinal plants and have been an important part of folk medicine globally in India and China<sup>11-15</sup>. These bioactive compounds showcase a significant spectrum of biological actions that include anti-cancer, anti-allergic, anti-mutagenic, anti-cataract, anti-toxic and the modulation of enzyme actions<sup>16,17</sup>. This review article focuses on the multifarious actions of dietary flavonoids.

**Table 1**  
**Different subclasses of flavonoids and their sources**<sup>3, 18</sup>

Sl.no	Sub-classes of flavonoids	Important flavonoids present	Source
1.	Flavonones	Naringen, Herpesitin	Citrus peels
2.	Flavonol	Quercetin, Rutin	Tea, Onion
3.	Catechins	Catechins, Epicatechins	Red wine, Tea, Fruits
4.	Isoflavones	Daidzen, Glycindin	Beans
5.	Flavones	Apigenin, Luteolin, Quercetin	Citrus fruit, Parsley
6.	Anthocyanidins	Cyanidin, Delphinidin, Malvidin	Grapes, apple skin, celery, Berries, Olives, grapes, Tea.

## MULTIFARIOUS ACTIONS OF DIETARY FLAVONOIDS

### 1. Flavonoids and Carcinogenesis

Any disturbance in the growth metabolism leads to cancer<sup>19</sup>. The main contributing factors are inadequate anti-oxidant systems and the presence of high levels of ROS and free radicals<sup>20,21</sup>. These ROS directly interfere with cell growth and signaling. The damage by ROS at cellular level induces mitosis and hence increases the chances of the damaged DNA to form mutations. Many studies have revealed potent anti-oxidants that suppress carcinogenesis. Studies involving fenugreek seed extracts proved to be potent anti-colon cancer agents by altering the expression of phospholipase A and by decreasing the levels of cholesterol<sup>22</sup>. Anti-tumourigenic activity of fenugreek seed extracts were also revealed in 1,2-dimethylhydrazine (DMH) induced wistar

rats<sup>23</sup>. Flavonoids are effective bioactive compounds that can interfere with the progression of cancer by various modulations (Fig 1). Flavonoids such saponins were found to be significant inhibitors of cell proliferation<sup>23</sup>. Studies showed an inverse correlation between occurrence of lung cancer and intake of flavonoid quercetin. Studies in mice revealed the anti-metastatic capacity of apigenin and quercetin in mice<sup>24</sup>. Angiogenesis is a controlled process in the body, that is activated mainly during wound healing. Unregulated angiogenesis leads to carcinogenesis<sup>25</sup>. Flavonoids were found to interfere with the endothelial cell migration, protein kinase production and lumen formation and hence proved to be potential angiogenesis inhibitors<sup>23, 26, 27</sup>. Various research groups showed the *invitro* anti-cancer effect of various flavonoids on different cell lines (Table 2).


**Table 2**  
**Invitro anti-cancer effects of flavonoids**

Sl.no	Type of cancer	Cell line used	Flavonoid	Reference
1.	Human lung cancer	A549, H441, H661, SW900	Quercetin, flavones	28, 29
2.	Human oral cancer	HSC-2, HSG, SCC-25	Quercetin, chalcones	30-34
3.	Human thyroid cancer	ARO,WRO,NPA	Apigenin, kaempferol, catechin, luteolin,genistein	35, 36
4.	Human prostate cancer	DU145, PC3	Silymarin, myricetin, apigenin, luteolin, epicatechin, catechin	37-40
5.	Human breast cancer	MCF-7	Quercetin, luteolin, genistein	41, 42
6.	Human leukaemia	HL-60, K562	Quercetin, myricetin, apigenin, chalcones	43-46
7.	Human colon cancer	HCT-15, IEC-6,HT-29	Genistein, quercetin, anthocyanin	47-51
8.	Mouse melanoma cancer	4A5	Chalcones	52

2,4-Dimethoxybenzaldehyde (DMBA) induced cancer was inhibited by consuming dietary quercetin in hamster<sup>53</sup> and rats<sup>54</sup>. Studies by Khanduja *et al.*, also revealed that ellagic acid and quercetin inhibited lung cancer in mice<sup>55</sup>. Quercetin inhibited tumour formation in the small intestine<sup>56</sup>. In animal models, rats fed with 0.05% of chalcone, 2-hydroxy chalcone or quercetin in diet prevented 4-Nitroquinoline-1-oxide (4NQO) induced cancer. Studies involving A/J mice (57) showed that lung tumours were inhibited by catechin rich tea.

Investigations of 4-Methyl Nitrosamine-1-3, Pyridyl-1 Butanone (NNK) induced mice given black or green tea prior to the administration of the toxin significantly reduced the total number of tumours formed in mice. With intra-peritoneal administration of apigenin or quercetin, it was found that melanoma cancer was inhibited<sup>29</sup>. The preventive effect of apigenin was found to be due to the prevention of phosphorylation process of Mitogen Activated Protein Kinases (MAPK).

### Pathway of carcinogenesis and the potential benefits of flavonoids



**Figure 1**  
**Pathway of carcinogenesis and the potential benefits of flavonoids**

## II. Anti-Cataract Activity of Flavonoids

One of the major complications of diabetes mellitus is cataract. 60% of patients with diabetes are prone to this eye condition. The correlation between Diabetes and formation of cataract was discovered in many research studies<sup>59</sup>. The glucose from the extracellular region diffuses into the lens because of the uncontrolled action of the hormone insulin and hence leads to the formation of cataract. Cataractogenesis is a multi-functional process. It was recently found that many pharmacological actions of flavonoids can act in the inhibition of cataract. *In vitro* Studies using rat lens organ culture with hydrogen peroxide showed that small levels of quercetin prevented loss of transparency of lens induced by calcium and sodium influx<sup>60</sup>. Studies by Cornish *et al.*, 2002, showed that quercetin and 3'-O-methyl quercetin (its metabolite) were responsible for preventing damage in the eye lens<sup>61</sup>. Cataract induced by the compound selenite was prevented by an isolated bioflavonoid (methylated quercetin) from *Cochlospermum religiosum* leaves<sup>62</sup>, similarly the flavonoid from *Vitex negundo* leaves also protected rat eye lens from the cataract induced by selenite<sup>63</sup>. Venoruton, decreased the extent of leakage of lactate dehydrogenase and opacification of rat lens hence reducing cataract effects<sup>64</sup>. *In vivo* Studies The selenite induced cataract model in rats was mainly used to study the anti-cataract effects of flavonoids<sup>65, 66</sup>. In streptozotocin induced rodents the effect of quercetin rhamnoside was studied early as 1977<sup>67</sup>. The animals in the study did not develop cataract when treated with quercetin in contrast to the control animal. Similarly, isoflavone significantly reduced the prevalence of cataract and death in diabetic animals<sup>68</sup>. Studies involving a group of bioflavonoids and flavangenol showed the prevention of cataract in diabetic animals<sup>69</sup>. Studies on neonatal galactosemic rats involving topically applied quercetin on the orbital pouch showed reduced cataractogenesis in rat lens. Quercetin acted by inhibiting extracellular fluid and decreasing intracellular edema<sup>70</sup>. Similar studies by

Mohan *et al.*, showed the anti-cataract activity by topically applied myricetin and quercetin<sup>71</sup>. The flavonoid genistein decreased the development of cataract in rats induced by galactose in diet<sup>72</sup>. Studies by Isai *et al.*, showed that rutin prevented cataract induced by selenite in rat pups<sup>73</sup>. Studies involving the instillation of crude onion juice into the eyes of rats were found to inhibit the formation of cataract by 75%. Onion was found to contain quercetin in the form of quercetin 3,4' – diglucoside and quercetin 4' – glucoside<sup>74</sup>. The onion juice was also found to increase the anti-oxidant enzyme levels in rat lens.

Isolated flavonoids from black and green tea<sup>75</sup>, grape seeds, *Ginkgo biloba*<sup>76, 77, 78</sup>, *Vitex negundo*<sup>79</sup> were also found to have anti-cataract action in experimental selenite induced cataract animal models. Radiation induced cataract was found to be inhibited in rats by *Ginkgo biloba* extracts<sup>80</sup>. The role of flavonoid, catechin as an anti-cancer agent was found in N-methyl- N nitrosurea induced cataract rat models<sup>81</sup>. Flavanoid rich grape seed extract decreased apoptosis of human lens cells induced by hydrogen peroxide<sup>82</sup>. UV-induced stress was prevented by flavonoid, fisetin by modulating the MAPK pathways<sup>83</sup>. Other Medicinal Properties of Flavonoids

## III. Flavonoids and Thrombosis

Thromboxanes and endoperoxidases are formed from the metabolism of Arachidonic Acid (AA) that is formed during inflammation. This condition contributes to platelet aggregation and activation, which further leads to the formation of platelet thrombus and atherosclerosis. Studies in 1960's showed that coagulation of blood and platelet aggregation was reduced by a tea pigment<sup>84</sup>. Studies on monkeys and dogs<sup>85</sup> showed that flavonoids like myricetin, quercetin and kaempferol inhibited the aggregation of platelets. Investigations showed that free radicals are scavenged directly by flavonols and hence this maintains a good concentration of nitric oxide and endothelial prostaglandins<sup>86</sup>. It was reported that flavonoids inhibit the AA pathway by different mechanisms, hence preventing

platelet aggregation indirectly. *In vitro* studies by Van Wauwe *et al.*, showed that to the membrane of platelets, flavonoids bind and hence over a time period have an accumulation effect<sup>87</sup>.

#### **IV. Flavonoids as anti-gastric ulcer agents**

By inhibiting Protein Kinase C (PKC), protein phosphorylation and Cyclooxygenase (COX) formation, flavonoids inhibit gastric ulcer. Investigations also revealed that flavonoids act as favorable substitutes for the usual therapeutic agents. Acylated flavonoids were used as a substitute for aspirin to treat gastric cancer<sup>88</sup>. Gastric acids caused by aspirin were found to decrease in the presence of flavonoids from *Ocimum basilium*. Studies in rats by Tapas *et al.*, showed that intraperitoneally administered rutin, quercetin and kaempferol (25-100mg/kg body weight),

dose dependently inhibited gastric damage caused by ethanol<sup>89</sup>.

#### **V. Flavonoids and Atherosclerosis**

Atherosclerotic changes are promoted when the oxidation of Low Density Lipoproteins (LDLs) takes place in the presence of oxygen radicals. This causes damage to the endothelial walls and hence causes atherosclerosis. Literature, showed that regular intake of flavonoids prevented coronary heart disease<sup>90, 91</sup> and death caused by the disease in older men. Aria *et al.*, described that the intake of flavonoids and the total plasma cholesterol levels are inversely related<sup>92</sup>. Furthermore, an inverse correlation was found between flavonoid intake and incident dementia<sup>93</sup>. The effects of flavonoids on other cardiovascular diseases are tabulated (Table 3).

**Table 3**  
**The effect of flavonoids on cardiovascular diseases**

Sl.no	Cardiovascular diseases	Influence of flavonoids	Reference
1.	Arrhythmia	Oxidative stress is decreased	88
2.	Heart Failure	Scavenging of Reactive Oxygen Species	88
3.	Acute myocardial infarction	Inhibition of platelet aggregation	94
4.	Hypertension	Inhibition of NADPH oxidase	94
5.	Atherosclerosis	Decreased in leucocyte adhesion	89

#### **VI. Flavonoids and Osteoporosis**

Studies by Hegarty *et al* reported that in elderly women who drank regular tea had increased bone density when compared to those who did not consume tea at all. The researchers proposed that the presence of flavonoids in tea could be the reason for its protective effect against osteoporosis<sup>95</sup>.

#### **VII. Flavonoids and Rheumatic Arthritis**

Inflammation and autoimmunity are significant components of rheumatic disease. Painful joints are mainly associated with rheumatitis. Orally consumed flavonoids were found to be useful in the destruction of prostaglandins which are responsible for the pain. Flavonoids were also found to activate T-Lymphocytes that kill cells carrying the antigens and hence prevents inflammation of the joints<sup>95</sup>

#### **VIII. Anti-Allergic Activity of Flavonoids**

Calcium-dependent Adenosine Triphosphate (ATPase) and cyclic Adenosine Mono Phosphate (cAMP) phosphodiesterase are important mediators for the release of histamines from basophils and mast cells<sup>5</sup>. Carlo *et al.*, revealed that quercetin is a very useful anti-allergic flavonoid useful in conditions like hay fever and asthma. It prevents the immune cells and has a negative influence on the release and production of histamines<sup>96</sup>.

#### **IX. Antimicrobial activity of flavonoids**

Wang *et al*<sup>97</sup>, found that flavonoids had an anti-viral activity. They interfered with viruses such as adenovirus, respiratory syncytial virus, herpes simplex virus and para-influenza virus. Kaul *et al.*, described that flavonoids like quercetin interfered with the infective and

replicative abilities of viruses<sup>98</sup>. *In vitro* studies have only been done so far to prove the antiviral effects of flavonoids; still *in vivo* studies are needed to describe the mechanisms in detail. Studies on rotaviruses showed that glycone form of the flavonoids were more effective than the aglycone forms<sup>99</sup>. Flavonoids such as anti-HIV compound have been studied nearly for 20 years now. The studies concentrated on the inhibitory action of reverse transcriptase along with the anti-protease and anti-integrase activities of flavonoids. Different flavonoids were also found to have anti-fungal and anti bacterial activities<sup>100</sup> (Table 7).

### X. Flavonoids and Diabetes mellitus

Diabetes mellitus is a chronic disease that requires the proper control of the glucose levels in blood (101). Diabetes mellitus is mostly of genetic origin; hence all flavonoids cannot cure the disease. Studies by Havsteen *et al.*, have revealed that some flavonoids can control Diabetes mellitus<sup>88</sup> by acting as potential inhibitor of aldose reductase<sup>102</sup>. Many researchers have proved quercetin and chrysin to be a significant anti-diabetic flavonoid that acts by releasing insulin and by regenerating pancreatic islet cells. Studies also revealed that quercetin can stimulate the uptake of calcium ions from the islet cells and thus a potential drug for non-insulin dependent diabetes<sup>89</sup>. Flavonoid fisetin was found to be effective at a concentration of 10 mg/kg body weight to treat Diabetes mellitus<sup>103</sup>.

**Table 4**  
**Anti-microbial activities of Flavonoids<sup>100</sup>**

S.No	Properties	Organism	Flavonoid
1.	Anti-fungal property	<i>Fusarium solani</i> <i>Botrytis cinerea</i> <i>Candida tropicalis</i> <i>Candida albicans</i> <i>Azotobacter vinelandii</i> <i>Verticillium dahlia</i>	Chrysoerol, Quercetin Chlroflavonin, Epicatechin Echinacin, Rutin, Apigenin, Phaseolinisoflavan
2.	Anti-bacterial activity	<i>Staphylococcus aureus</i> <i>Staphylococcus albus</i> <i>Streptococcus baris</i> <i>Escherichia coli</i> <i>Bacillus subtilis</i> <i>Proteus vulgaris</i> <i>Bacillus anthracis</i>	Quercetin, Fisetin, Baicalin, Hesperitin, Apigenin, Rutin, Chrysin, Datisetin, Naringin
3.	Antiviral property	<i>Herpes virus</i> <i>Polio virus</i> <i>Pseudorabies virus</i> <i>Pasrinflenza virus</i> <i>Immuno-deficiency virus</i> <i>Mengo virus</i> <i>Respiratory syncytial virus</i>	Quercetin, Rutin, morin, naringin

### XI. Anti-Toxic Activity of Flavonoids

In the liver, flavonoids have the capacity to bind to RNA polymerase I and hence increase protein production. This helps in the formation and regeneration of hepatocytes<sup>5</sup>. Studies revealed that aqueous fenugreek leaf extracts rich in flavonoids ameliorated pesticide induced hepatotoxicity and nephrotoxicity in male wistar rats<sup>104</sup>. Venoruton and rutin were reported to enhance liver regeneration in experimental cirrhosis<sup>89</sup>. Studies by Kim *et al* showed that

isolated trifolin, isovitexin, quercetin, hirustin and avicularin had hepatoprotective activity in HepG2 cell lines<sup>105</sup>. Gulati *et al.*, investigated and proved quercetin as a potential hepatoprotective agent<sup>106</sup>. In human liver cell lines (Hep 2), different flavonoids such as luteolin, apigenin, quercetin-3-O-glucoside and kaempherol 3-O-glucoside showed hepatoprotective activity<sup>107</sup>. Summary of the effects of flavonoids on various diseases are tabulated in table 8.

**Table 5**  
**Summary of the effect of flavonoids on various diseases.**

SI.NO	DISEASE	FLAVONOID	MECHANISM	REFERENCE
1.	Inflammation	Quercetin, catechin, rutin, apigenin, fisetin, hesperidin, kaempferol, myricetin, fisetin.	Inducible Nitric Oxide Synthase (iNOS) and COX, PG synthesis	89, 108, 109, 110
2.	Thrombosis	Quercetin, rutin, sinesetin, catechol, hesperidin.	Erythrocyte aggregation, vascular permeability.	111
3.	Ulcer- Peptic and Gastric ulcer	Catechins, quercetin, kaempferol	PG synthesis	89, 111
4.	Atherosclerosis	Quercetin	Decrease in LDL oxidation, inhibition of leucocyte adhesion	88
5.	Osteoporosis	Quercetin	Increase in bone density	95
6.	Rheumatoid arthritis	Rutin, apigenin	Inflammation decreased by acting on COX	112, 108
7.	Anti-allergic	Rutin, quercetin, citrin	Act on mast cells	111, 113, 114
8.	Diabetes mellitus	Quercetin, Fisetin	Inhibits production of aldose reductase	112
9.	Hepatoprotective	Luteolin, quercetin, hirustrin	Binds to RNA polymerase I in the liver	115, 116
10.	Cancer-Lung cancer, oral cancer, thyroid cancer, prostate cancer, colon cancer and breast cancer.	Catechins, luteolin, quercetin, kaempferol, apigenin	Enhance the action of catalase, glutathione reductase, glutathione peroxidase, inhibits the action of tyrosine kinase.	111, 88
11.	Memory dysfunction	Quercetin, fisetin, genistein	Inhibits tyrosine kinase	114

## CONCLUSION AND FUTURE IMPLICATIONS

The studies involving flavonoids is complex due to the heterogeneity of various molecular structures. They form a group of biologically active substances that are present in high amounts in plants and consumed in large parts in our daily diet. Flavonoids are gaining more and more importance because of their usefulness and significant roles that they play inside the human body. Future implications involve exploring more potential properties of

flavonoids in the field of immunomodulation and anti-toxic activities.

## ACKNOWLEDGEMENT

The first author would like to acknowledge the financial support provided by DST, New Delhi under the INSPIRE FELLOWSHIP SCHEME Proc No. 8946/PD6/2010.

## REFERENCES

1. Simeonova R, Krasteva I, Kondeva-Burdina M, Benbassat N, Effects of extract from astragalus glycyphylloides on carbon Tetrachloride-induced hepatotoxicity in wistar rats, Int J Pharm Bio Sci 4: 179 – 186 (2013).
2. Shoba G, Shyamala H, Prabhavathi G, Stella,S, Flavonoids - Natural Therapeutic Agents For Polycystic Kidney Disease , Int J Pharm Bio Sci 1: 89 – 105 (2010).
3. Hollman PCH, Katan MB, Dietary Flavonoids: Intake, Health Effects and Bioavailability, Food and Chemical Toxicology, 37: 937-942, (1999).
4. Cushnie TPT, Lamb AJ, Antimicrobial activity of flavonoid,. International Journal Of Antimicrobial Agents,26: 343-356, (2005).
5. Murray MT, Quercetin: Nature's antihistamine, Better Nutrition, (1998).

6. Peterson J, Dwyer MSJ, RD, DSc, Flavonoids: Dietary occurrence and biochemical activity, *Nutrition Research*, 18: 1995-2018, (1998).
7. Tsuchiya H. Structure-dependent membrane interaction of flavonoids associated with their bioactivity, *Food Chemistry* 120: 1089-1096, (2010).
8. Anirudha. R, In-Vitro And In-Vivo Flavonoid Content In *Urgenia Indica* (Roxb.) Kunth, *Int J Pharm Bio Sci* , 4 : 242 - 246 , (2013).
9. Harborne JB, Williams CA, Advances in flavonoid research since 1992, *Phytochemistry*, 55:48–504, (2000).
10. Hollman PC, Katan MB, Dietary flavonoids: Intake, health effects and bioavailability, *Food Chem Toxicol*, 37:937–942, (1999).
11. Di Carlo G, Mascolo N, Izzo AA, Capasso F, Flavonoids: Old and new aspects of a class of natural therapeutic drugs, *Life Sci*, 65:337–353, (1999).
12. Craig WJ, Health-promoting properties of common herbs, *Am J Clin Nutr*, 70:491S–499S , (1999).
13. Kadarian C, Broussalis AM, Mino J, Lopez P, Gorzalczany S, Ferraro G, Acevedo C, Hepatoprotective activity of *Achyrocline satureioides*(Lam) D. C, *Pharmacol Res*, 45:57–61 , (2002).
14. Pascual ME, Slowing K, Carretero E, Sanchez Mata D, Villar A, Lippia: Traditional uses, chemistry and pharmacology: A review, *J Ethnopharmacol*, 76:201–214, (2001).
15. Samuelsen AB, The traditional uses, chemical constituents and biological activities of *Plantago major* L. A review, *J Ethnopharmacol*, 71:1–21, (2000).
16. Middleton E, Jr., Kandaswami C, Theoharides TC, The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease, and cancer, *Pharmacol Rev* , 52:673–751, (2000).
17. Galati G, Teng S, MoridaniMY, Chan TS, O'Brien PJ, Cancer chemoprevention and apoptosis mechanisms induced by dietary polyphenolics, *Drug Metabol Drug Interact*, 17:311–349, (2000).
18. Ren, W, Qiao, Z, Wang, H, Zhu, L, Zhang, L, Flavonoids: Promising Anticancer agents, *Medicinal Research Reviews* , 23: 519-534, (2003).
19. Hertog MG, Feskens EJ, Hollman PC, Katan MB, Kromhout D, Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study, *Lancet* , 342:1007–11, (1993).
20. Loft S, Poulsen HE, Cancer risk and oxidative DNA damage in man, *J Mol Med*, 74:297–312, (1996). [Published erratum appears in *J Mol Med* , 75:67–8, (1997)]
21. Pryor WA, Cigarette smoke radicals and the role of free radicals in chemical carcinogenicity, *Environ Health Perspect* , 105(suppl): 875–82, (1997).
22. Devasena T, Gunasekaran G, Viswanathan P, Menon VP, Chemoprevention of 1,2- dimethyl hydrazine-induced colon carcinogenesis by seeds of *Trigonella foenum graceum* L, *Biologia. Bratislava*, 58:357-364 (2003).
23. Devasena.T, Antitumorigenic action of fenugreek seeds, *Asian J of Bio Science*, 4: 83-87, (2009).
24. Caltagirone S, Rossi C, Poggi A, *et al*, Flavonoids apigenin and quercetin inhibit melanoma growth and metastatic potential, *Int J Cancer* , 87:595–600, (2000).
25. Fan TP, Jaggar R, Bicknell R, Controlling the vasculature: angiogenesis, anti-angiogenesis and vascular targeting of gene therapy, *Trends Pharmacol Sci* , 16:57–66, (1995).
26. Paper DH, Natural products as angiogenesis inhibitors, *Planta Med*, 64:686–95, (1998).
27. Oikawa T, Shimamura M, Ashino H, *et al*, Inhibition of angiogenesis by staurosporine, a potent protein kinase inhibitor, *J Antibiot (Tokyo)*, 45:1155–60, (1992).
28. Bai F, Matsui T, Ohtani-Fujita N, Matsukawa Y, Ding Y, Sakai T, Promoter activation and following induction of the p21/WAF1 gene by flavone is involved in G1 phase arrest in A549 lung



- adenocarcinoma cells, FEBS Lett, 437:61–64, (1998).
29. Caltagirone S, Ranelletti FO, Rinelli A, Maggiano N, Colasante A, Musiani P, Aiello FB, Piantelli M, Interaction with type II estrogen binding sites and antiproliferative activity of tamoxifen and quercetin in human non-small-cell lung cancer, *Am J Respir Cell Mol Biol*, 17:51–59, (1997).
  30. Shi YQ, Fukai T, Sakagami H, Chang WJ, Yang PQ, Wang FP, Nomura T, Cytotoxic flavonoids with isoprenoid groups from *Morus mongolica*, *J Nat Prod*, 64:181–188, (2001).
  31. Fukai T, Sakagami H, Toguchi M, Takayama F, Iwakura I, Atsumi T, Ueha T, Nakashima H, Nomura T, Cytotoxic activity of low molecular weight polyphenols against human oral tumor cell lines, *Anticancer Res*, 20:2525–2536, (2000).
  32. Sakagami H, Jiang Y, Kusama K, Atsumi T, Ueha T, Toguchi M, Iwakura I, Satoh K, Fukai T, Nomura T, Induction of apoptosis by flavones, flavonols (3-hydroxyflavones) and isoprenoid substituted flavonoids in human oral tumor cell lines, *Anticancer Res*, 20:271–277, (2000).
  33. Elattar TM, Virji AS, Effect of tea polyphenols on growth of oral squamous carcinoma cells in vitro, *Anticancer Res*, 20:3459–3465, (2000).
  34. Elattar TM, Virji AS, The inhibitory effect of curcumin, genistein, quercetin and cisplatin on the growth of oral cancer cells in vitro, *Anticancer Res*, 20:1733–1738, (2000).
  35. Yin F, Giuliano AE, Van Herle AJ, Signal pathways involved in apigenin inhibition of growth and induction of apoptosis of human anaplastic thyroid cancer cells (ARO), *Anticancer Res*, 19:4297–4303, (1999).
  36. Yin F, Giuliano AE, Van Herle AJ, Growth inhibitory effects of flavonoids in human thyroid cancer cell lines, *Thyroid*, 9:369–376 (1999).
  37. Knowles LM, Zigrossi DA, Tauber RA, Hightower C, Milner JA, Flavonoids suppress androgen independent human prostate tumor proliferation, *Nutr Cancer*, 38:116–122, (2000).
  38. Bhatia N, Agarwal R, Detrimental effect of cancer preventive phytochemicals silymarin, genistein and epigallocatechin 3-gallate on epigenetic events in human prostate carcinoma DU145 cells, *Prostate*, 46:98–107, (2001).
  39. Kampa M, Hatzoglou A, Notas G, Damianaki A, Bakogeorgou E, Gemetzi C, Kouroumalis E, Martin PM, Castanas E, Wine antioxidant polyphenols inhibit the proliferation of human prostate cancer cell lines, *Nutr Cancer* 37:223–233, (2000).
  40. Agarwal R, Cell signaling and regulators of cell cycle as molecular targets for prostate cancer prevention by dietary agents, *Biochem Pharmacol*, 60:1051–1059, (2000).
  41. Pouget C, Lauthier F, Simon A, Fagnere C, Basly JP, Delage C, Chulia AJ, Flavonoids: Structural requirements for antiproliferative activity on breast cancer cells, *Bioorg Med Chem Lett*, 11:3095–3097, (2001).
  42. Han D, Tachibana H, Yamada K, Inhibition of environmental estrogen-induced proliferation of human breast carcinoma MCF-7 cells by flavonoids, *In Vitro Cell Dev Biol Anim*, 37:275–282, (2001).
  43. Wang IK, Lin-Shiau SY, Lin JK, Induction of apoptosis by apigenin and related flavonoids through cytochrome c release and activation of caspase-9 and caspase-3 in leukaemia HL-60 cells, *Eur J Cancer*, 35:1517–1525, (1999).
  44. Chung HS, Chang LC, Lee SK, Shamon LA, van Breemen RB, Mehta RG, Farnsworth NR, Pezzuto JM, Kinghorn AD, Flavonoid constituents of *Chorizanthe diffusa* with potential cancer chemopreventive activity, *J Agric Food Chem*, 47:36–41, (1999).
  45. Csokay B, Prajda N, Weber G, Olah E, Molecular mechanisms in the antiproliferative action of quercetin, *Life Sci*, 60:2157–2163, (1997).
  46. De Vincenzo R, Ferlini C, Distefano M, Gaggini C, Riva A, Bombardelli E, Morazzoni P, Valenti P, Belluti F, Ranelletti

- FO, Mancuso S, Scambia G, In vitro evaluation of newly developed chalcone analogues in human cancer cells, *Cancer Chemother Pharmacol*, 46:305–312, (2000).
47. Wenzel U, Kuntz S, Brendel MD, Daniel H, Dietary flavone is a potent apoptosis inducer in human colon carcinoma cells, *Cancer Res*, 60:3823–3831, (2000).
  48. Kamei H, Hashimoto Y, Koide T, Kojima T, Hasegawa M, Anti-tumor effect of methanol extracts from red and white wines, *Cancer Biother Radiopharm*, 13:447–452, (1998).
  49. Kuntz S, Wenzel U, Daniel H, Comparative analysis of the effects of flavonoids on proliferation, cytotoxicity, and apoptosis in human colon cancer cell lines, *Eur J Nutr* 38:133–142, (1999).
  50. Kuo SM, Morehouse HF, Jr., Lin CP, Effect of antiproliferative flavonoids on ascorbic acid accumulation in human colon adenocarcinoma cells, *Cancer Lett*, 116:131–137, (1997).
  51. Kuo SM, Antiproliferative potency of structurally distinct dietary flavonoids on human colon cancer cells, *Cancer Lett*, 110:41–48, (1996).
  52. Iwashita K, Kobori M, Yamaki K, Tsushida T, Flavonoids inhibit cell growth and induce apoptosis in B16 melanoma 4A5 cells, *Biosci Biotechnol Biochem* , 64:1813–1820, (2000).
  53. Balasubramanian S, Govindasamy S, Inhibitory effect of dietary flavonol quercetin on 7,12-dimethylbenz[a]anthracene-induced hamster buccal pouch carcinogenesis, *Carcinogenesis* , 17: 877–879, (1996).
  54. Verma AK, Johnson JA, Gould MN, Tanner MA, Inhibition of 7,12 dimethylbenz(a) anthracene and N-nitrosomethylurea induced rat mammary cancer by dietary flavonol quercetin, *Cancer Res*, 48:5754–5758, (1998).
  55. Khanduja KL, Gandhi RK, Pathania V, Syal N, Prevention of N-nitrosodiethylamine-induced lung tumorigenesis by ellagic acid and quercetin in mice, *Food Chem Toxicol*, 37:313–318, (1999).
  56. Akagi K, Hirose M, Hoshiya T, Mizoguchi Y, Ito N, Shirai T, Modulating effects of ellagic acid, vanillin and quercetin in a rat medium term multi-organ carcinogenesis model, *Cancer Lett*, 94:113–121, (1995).
  57. Yang CS, Chung JY, Yang GY, Chhabra SK, Lee MJ, Tea and tea polyphenols in cancer prevention, *J Nutr*, 130:472S–478S, (2000).
  58. Ramos, S, Effects of dietary flavonoids on apoptic pathways related to cancer chemoprevention, *Journal of Nutritional Biochemistry*, 18: 427-442, (2007).
  59. Obrosova IG, Chung SS, Kador PF, Diabetic cataracts: mechanisms and management., *Diabetes Metab Res Rev* , 26: 172–180, (2010).
  60. Sanderson J, McLauchlan WR, Williamson G, Quercetin inhibits hydrogen peroxide-induced oxidation of the rat lens, *Free Rad Biol Med*, 26: 639–645, (1999).
  61. Cornish KM, Williamson G, Sanderson J, Quercetin metabolism in the lens: role in inhibition of hydrogen peroxide induced cataract, *Free Rad Biol Med*, 33: 63–70, (2002).
  62. Gayathri Devi V, Rooban BN, Sasikala V, Sahasranamam V, Abraham A, Isorhamnetin-3-glucoside alleviates oxidative stress and opacification in selenite cataract in vitro, *Toxicol in Vitro*, 24: 1662–1669, (2010).
  63. Rooban BN, Sasikala V, Sahasranamam V, Abraham A, Amelioration of selenite toxicity and cataractogenesis in cultured rat lenses by Vitex negundo, *Graefes Arch Clin Exp Ophthalmol*, (2011).
  64. Kilic F, Bhardwaj R, Trevithick JR, Modelling cortical cataractogenesis XVIII In vitro diabetic cataract reduction by venoruton A flavonoid which prevents lens opacification, *Acta Ophthalmol Scand* 74: 372–378, (1996).
  65. Gupta SK, Selvan VK, Agrawal SS, Saxena R, Advances in pharmacological strategies for the prevention of cataract

- development, *Indian J Ophthalmol*, 57: 175–183, (2009).
66. Kyselova Z, Stefek M, Bauer V, Pharmacological prevention of diabetic cataract, *J Diabetes Complications*, 18: 129–140, (2004).
  67. Varma SD, Mizuno A, Kinoshita JH, Diabetic Cataracts and Flavonoids, *Science*, 195: 205–206, (1977).
  68. Lu M-P, Wang R, Song X, Chibbar R, Wang X, Wu L, Q Meng QH, Dietary soy isoflavones increase insulin secretion and prevent the development of diabetic cataracts in streptozotocin-induced diabetic rats, *Nutr Res*, 28: 464–471, (2008).
  69. Nakano M, Orimo N, Katagiri N, Tsubata M, Takahashi J, Van Chuyen N, Inhibitory effect of astraxanthin combined with Flavangenol on oxidative stress biomarkers in streptozotocin-induced diabetic rats, *Int J Vitam Nutr Res*, 78: 175–182, (2008).
  70. Beyer-Mears A, Farnsworth NP, Diminished sugar cataractogenesis by quercetin, *Exp Eye Res*, 28: 709–716, (1979).
  71. Mohan M, Gupta SK, Agnihotri S, Joshi S, Uppal RK, Anticataract action of topical quercetin and myricetin in galactosemic rats, *Med Sci Res*, 6: 685–686, (1988).
  72. Huang R, Shi F, Lei T, Song Y, Hughes CL, Liu G, Effect of the isoflavone genistein against galactose induced cataracts in rats, *Exp Biol Med (Maywood)* 232: 118–125, (2007).
  73. Isai M, Sakthivel M, Ramesh E, Thomas PA, Geraldine P, Prevention of selenite-induced cataractogenesis by rutin in Wistar rats, *Mol Vis*, 15: 2570–2577, (2009).
  74. Javadzadeh A, Ghorbanihaghjo A, Bonyadi S, Rashidi MR, Mesgari M, Rashtchizadeh N, Argani H, Preventive effect of onion juice on selenite-induced experimental cataract, *Indian J Ophthalmol*, 57: 185–189, (2009).
  75. Gupta SK, Halder N, Srivastava S, Trivedi D, Joshi S, Varma SD, Green tea (*Camellia sinensis*) protects against selenite-induced oxidative stress in experimental cataractogenesis, *Ophthalmic Res*, 34: 258–263, (2002).
  76. Thiagarajan G, Chandani S, Harinarayana Rao S, Samuni AM, Chandrasekaran K, Balasubramanian D, Molecular and cellular assessment of Ginkgo biloba extract as a possible ophthalmic drug, *Exp Eye Res*, 75: 421–430, (2002).
  77. Durukan AH, Evereklioglu C, Hurmeric V, Kerimoglu H, Erdurman C, Bayraktar MZ, Mumcuoglu T, Ingestion of IH636 grape seed proanthocyanidin extract to prevent selenite-induced oxidative stress in experimental cataract, *J Cataract Refract Surg*, 32: 1041–1045, (2006).
  78. Vibin M, Siva Priya SG, N Rooban B, Sasikala V, Sahasranamam V, Abraham A, Broccoli regulates protein alterations and cataractogenesis in selenite models, *Curr Eye Res*, 35: 99–107, (2010).
  79. Rooban BN, Lija Y, Biju PG, Sasikala V, Sahasranamam V, Abrahama A, Vitex negundo attenuates calpain activation and cataractogenesis in selenite models, *Exp Eye Res*, 88: 575–582, (2009).
  80. Ertekin MV, Kocer I, Karslioglu I, Taysi S, Gepdiremen A, Sezen O, Balci E, Bakan N, Effects of oral ginkgo biloba supplementation on cataract formation and oxidative stress occurring in lenses of rats exposed to total cranium radiotherapy, *Jpn J Ophthalmol*, 48: 499–502, (2004).
  81. Lee SM, Ko IG, Kim SE, Kim DH, Kang BN, Protective effect of catechin on apoptosis of the lens epithelium in rats with N-methyl-N-nitrosourea-induced cataracts, *Korean J Ophthalmol* 24: 101–107, (2010).
  82. Jia Z, Song Z, Zhao Y, Wang X, Liu P, Grape seed proanthocyanidin extract protects human lens epithelial cells from oxidative stress via reducing NF- $\kappa$ B and MAPK protein expression, *Mol Vis*, 17: 210–217, (2011).
  83. Yao K, Zhang L, Zhang Y, Ye P, Zhu N, The flavonoid fisetin inhibits UV radiation-induced oxidative stress and the activation of NF-kappaB and MAPK signaling in

- human lens epithelial cells, *Mol Vis*, 14: 1865–1871, (2008).
85. Lou FQ, Zhang MF, Zhang XG, Liu JM, Yuan WL, A study on tea pigment in prevention of atherosclerosis, *Chin Med J (Engl)*, 102:579–83, (2008).
  86. Osman HE, Maalej N, Shanmuganayagam D, Folts JD, Grape juice but not orange or grapefruit juice inhibits platelet activity in dogs and monkeys, *J Nutr*, 128:2307–121 (1998).
  87. Gryglewski RJ, Korbut R, Robak J, Swies J, On the mechanism of antithrombotic action of flavonoids, *Biochem Pharmacol*, 36: 317–22, (1987).
  88. Van Wauwe J, Goossens J, Effects of antioxidants on cyclooxygenase and lipoxygenase activities in intact human platelets: comparison with indomethacin and ETYA, *Prostaglandins*, 26:725–30, (1983).
  89. Havsteen BH, The biochemistry and medical significance of the flavonoids, *Pharmacology and Therapeutics*, 96: 67-202, (2002).
  90. Tapas AR, Sakarkar DM, Kakde RB, Flavonoids as nutraceuticals: A Review, *Tropical Journal of Pharmaceutical Research*, 7: 1089-1099, (2008).
  91. Hertog MG, Kromhout D, Aravanis C, *et al*, Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study, *Arch Intern Med*, 155:381–6, (1995).
  92. Hertog MG, Feskens EJ, Hollman PC, Katan MB, Kromhout D, Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study, *Lancet*, 342:1007–11, (1993).
  93. F Y, Watanabe S, Kimira M, Shimoi K, Mochizuki R, Kinae N, Dietary intakes of flavonols, flavones and isoflavones by Japanese women and the inverse correlation between quercetin intake and plasma LDL cholesterol concentration, *J Nutr*, 130: 2243–50, (2000).
  94. Commenges D, Scotet V, Renaud S, Jacqmin-Gadda H, Barberger- Gateau P, Dartigues JF, Intake of flavonoids and risk of dementia, *Eur J Epidemiol*, 16:357–63, (2000).
  95. Loft S, Poulsen HE, Cancer risk and oxidative DNA damage in man, *J Mol Med*, 74:297–312, (1996). (Published erratum appears in *J Mol Med*, 75:67–8, 1997)
  96. Hegarty VM, May HM, Khaw KT, Tea drinking and bone mineral density in older women, *Am J Clin Nutr*, 71:1003–7(2000),
  97. Carlo, GD, Mascolo, N, Izzo, AA, Capasso, F. Flavonoids, Old and New aspects of a class of Natural Therapeutic Drugs, *Life Sciences*, 65: 337-353, (1999).
  98. Wang HK, Xia Y, Yang ZY, Natschke SL, Lee KH, Recent advances in the discovery and development of flavonoids and their analogues as antitumor and anti-HIV agents, *Adv Exp Med Biol*, 439: 191–225, (1998).
  99. Kaul TN, Middleton E Jr, Ogra PL, Antiviral effect of flavonoids on human viruses, *J Med Virol*, 15:71–9, (1985).
  100. Bae EA, Han MJ, Lee M, Kim DH, In vitro inhibitory effect of some flavonoids on rotavirus infectivity, *Biol Pharm Bull*, 23:1122–4, (2000).
  101. Narayana KR, Reddy SR, Chaluvadi MR, Krishna DR, Bioflavonoids classification, pharmacological, biochemical effects and therapeutic potential, *Indian Journal of Pharmacology*, 33: 2-16, (2001).
  102. Zhou, T, Luo, D, Li, X, Luo, Y, Hypoglycaemic and hypolipidemic effects of flavonoids from lotus (*Nelumbo nuficera*) leaf in diabetic mice, *Journal of medicinal Plants Research*, 3: 290-293, (2009).
  103. Patra JC, Chua BH, Artificial neutral network-based drug design for diabetes mellitus using flavonoids, *Journal of Computational Chemistry*, 32: 555-567, (2010).
  104. Sriram PG, Subramanian S. Fisetin, a bioflavonoid ameliorates hyperglycaemia in streptozotocin-induced experimental diabetes in rats, *International Journal of Pharmaceutical Sciences Review and Research*, 6: 68-74, (2011).
  105. Sushma N and Devasena T, Aqueous extract of *Trigonella foenum graecum*

- (fenugreek) prevents cypermethrin-induced hepatotoxicity and nephrotoxicity, Human and Experimental Toxicology, 29(4) 311–319, (2010).
106. Kim SM, Kang K, Jho EH, Jung Y, Nho CW, Um B, Pan C, Hepatoprotective effect of flavonoids glycoside from *Lespedeza cuneata* against oxidative stress induced by tert-Butyl hyperoxide, Phytotherapy Research, ( 2011).
107. Gulati RK, Agarwal S, Agarwal SS, Hepatoprotective studies on *Phyllanthus emblica* and quercetin, Indian Journal of Experimental Biology, 33: 261-268, (1995).
108. Spencer, JPE, Vanzour, D, Rendeiro, C, Flavonoids and cognition: The molecular mechanisms underlying their behavioural effects, Archives of Biochemistry and Biophysics, 492: 1-9, (2009).
109. Guardia, T, Rotelli, AE, Osvaldo, A, Anti-inflammatory properties of plant flavonoids, Effect of rutin, quercetin and hesperidin on adjuvant arthritis in rats, Il farmaco, 56, (2001).
110. Miller AL, Antioxidant flavonoids: Structure, function and Clinical Usage, Alternative Medicine Review, 1: 103-111, (1996).
111. Marchand LL, Cancer preventive effects of flavonoids- a review, Biomedicine & Pharmacotherapy, 56: 296-301, (2002).
112. Narayana, KR, Reddy, SR, Chaluvadi, MR, Krishna, DR, Bioflavonoids classification, pharmacological, biochemical effects and therapeutic potential, Indian Journal of Pharmacology, 33: 2-16 , (2001).
113. Kang H, Ecklund D, Liu M, Datta SK, Apigenin, a non-mutagenic dietary flavonoids, suppresses lupus by inhibiting autoantigen presentation for expansion of autoreactive Th1 and Th17 cells, Arthritis Research and Therapy, 11: 1-13, (2009).
114. Venugopal SK, Devaraj S, Jialal I, Macrophage conditioned medium induces the expression of C-reactive protein in human aortic endothelial cells: potential for paracrine/autocrine effects, Am J Pathol, 166:1265-1271, (2005).
115. Maher, P, Akalshi, T, Abe, K, Flavonoid fisetin promotes ERK-dependent long-term potentiation and enhances memory, Proceedings of National academy of Sciences United States of America, 103: 16568-16573, (2006).
116. Gulati, RK, Agarwal, S, Agarwal, SS, Hepatoprotective studies on *Phyllanthus emblica* and quercetin, Indian Journal of Experimental Biology, 33: 261-268, (1995).
117. Oh, H, Kim, D, Cho, J, Kim, Y, Hepatoprotective and free radical activities of phenolic petrosins and flavonoids isolated from *Equisetum arvense*, Journal of Ethnopharmacology, 95: 421-424, (2004).