



EFFECT OF HESPERTIN ISOLATED FROM ORANGE PEELS ON CISPLATIN-INDUCED NEPHROTOXICITY

SPANDANA.U*¹, SREEDEVI.A ², HAVILAH ESTHER.S³ AND LAKSHMAIAH .P⁴

^{1,2} Sri padmavathi mahila viswa vidyalayam , Tirupathi, A.P, India.

³ Nirmala college of Pharmacy, Atmakur, Guntur, A.P,India.

⁴ Viswabharathi college of Pharmacy, Perecharla, Guntur, A.P, India.

ABSTRACT

The present study is to determine the effect of hespertin isolated from orange peels on cisplatin-induced toxicity and Isolation of hespertin from the hesperdin by acid degradation to determine its effect on cisplatin – induced nephrotoxicity. In the present study, the hespertin (200 mg & 400 mg/ kg bd.wt) was examined for its protective effect against cisplatin induced renal injury in rats. Thirty healthy male and female albino rats (150- 200 g weight) were chosen and divided into five groups. Vehicle, cisplatin and hespertin were given according to the experimental design. After 8 days of treatment, urinary functional parameters were analysed. Histopathological characters were examined after scarifying and dissecting the rats. Cisplatin (6 mg/kg) alone significantly elevated serum markers level, increased urinary protein excretion, reduced urine to serum creatinine ratio and creatinine clearance. In curative regimen, the extract significantly reduced the elevated serum creatinine and urea levels. Renal antioxidant defense systems, such as superoxide dismutase, catalase, glutathione peroxides activities and reduced glutathione level, depleted by cisplatin therapy were restored to normal by treatment with the extract. The present investigation suggests the hespertin protective effect against cisplatin induced nephrotoxicity.

KEYWORDS: Cisplatin; hespertin; nephrotoxicity; Creatinine; hesperdin.



SPANDANA.U

Sri padmavathi mahila viswa vidyalayam , Tirupathi, A.P, India.

*Corresponding author

INTRODUCTION

Plant

The orange is both a literal and symbolic embodiment of the sun, from whose light it is formed. As a whole food it irradiates us with a spectrum of healing properties, the most prominent of which some call "vitamin C activity," but which is not reducible to the chemical skeleton known as 'ascorbic acid.' Science now confirms the orange has a broad range of medicinal properties, which is why the ancients knew it both as a food and a medicine¹. Oranges is an important medicinal plant of the family Rutaceae. It is cultivated mainly for its alkaloids, which are having anticancer activities and the antibacterial potential in crude extracts of different parts (viz., leaves, stem, root and flower)². The limonene and flavonoids found in orange peel seem to have anti-carcinogenic properties by acting as a blocking agent. Studies have shown that limonin and limonene, both found in high concentrations in the peel, can induce the enzyme activity of glutathione S-transferase, which is an

important detoxifying enzyme. The citric acid found in orange peels also helps starve cancer cells by cutting off their energy supply. A number of studies have shown decreased risk of several cancers, most notably skin, breast and colon cancer, linked with the increased consumption of orange peel. In addition, orange peels and orange peel extract can provide an extra benefit to diabetics and those looking to reduce overeating. This is due to the fact that orange peels are a natural source of pectin, a natural fiber that helps decrease the rise in blood sugar after a meal. It may also be helpful in lowering cholesterol. D-limonene, another biochemical found in orange peels, is helpful in dissolving kidney stones. Orange peel has antiseptic, bactericidal, and fungicidal properties, which are extremely beneficial in eliminating any kidney or urinary tract infections. Orange peels may also help with heartburn. According to a 2007 "Alternative Medicine Review" article by Jidong Sun, Ph.D., Sun, d-limonene possesses the ability to neutralize gastric acid and support normal peristalsis³.




Flavonoids

Phytochemicals are defined as the substances found in edible fruits and vegetables that exhibit a potential for modulating human metabolism in a manner beneficial for the prevention of chronic and degenerative diseases⁴. The most important natural pigments are carotenoids which are tetrapyrrole derivatives of naturally occurring phenolic compounds ubiquitously distributed in plant kingdom⁵. Phenolics are defined as a class of polyphenols which are important secondary metabolites present in plants⁶ and are also responsible for their antioxidant action and various beneficial effects in a multitude of diseases^{7,8}. Among these compounds, flavonoids constitute one of the most ubiquitous groups of all plant phenolics. So far, over 8,000 varieties of flavonoids have been identified⁵. Flavonoids are low molecular weight^{9,10} bioactive polyphenols¹¹ which play a vital role in photosynthesising cells¹⁰. Flavonoids are secondary metabolites characterised by flavan nucleus¹³ and C6-C8-C6 carbon-skeleton^{14,15}. These are group of structurally related compounds with a chromane-type skeleton having phenyl substituent in C2-C3 position¹⁶. They are ubiquitously found in fruits,

vegetables, tea, and wine, and are usually subdivided into six classes including flavonols (e.g., quercetin, kaempferol), flavones (e.g., apigenin, luteolin), flavanones (e.g., hesperidin, naringenin), flavan-3-ols (e.g., catechin, theaflavin, and gallic esters of catechin and theaflavins), anthocyanidins (e.g., pelargonidin, cyanidin), and isoflavones (e.g., genistein, daidzein). Secondary metabolism of flavonoids in plants generates glycosides through linkage to a sugar moiety, which enhances absorption in humans¹⁷. Flavonoids occur as aglycones, glycosides and methylated derivatives¹⁸ flavonoids aglycones (i.e., flavonoids without attached sugar) occur in a variety of structural forms. All contain fifteen carbon atoms in their basic nucleus: two six-membered rings linked with a three carbon unit which may or may not be a part of a third ring¹⁹.

Hesperetin

It is a bioflavonoid and, to be more specific, a flavanone. A trihydroxyflavanone having the three hydroxy groups located at the 3', 5- and 7-positions and an additional methoxy substituent at the 4'-position.



Molecular Formula: $C_{16}H_{14}O_6$

IUPAC Name: (2S)-5,7-Dihydroxy-2-(3-hydroxy-4-methoxyphenyl)-2,3-dihydro-4H-chromen-4-one

The average intake of hesperetin estimate is 28.3 mg/d.

Hesperetin (and orange juice) has been shown to inhibit chemically induced mammary²⁰, urinary bladder²¹, and colon^{22,23} carcinogenesis in laboratory animals. Hesperetin²⁴ as well as the other major citrus flavanone naringenin^{25,26}, also possess some antioxidant activities, although this activity is poorer compared with many other polyphenols. Other possible effects of hesperetin and naringenin are on lipid metabolism. They have been reported to regulate apolipoprotein B secretion by HepG2 cells, possibly through inhibition of cholesterol ester synthesis²⁷, and to inhibit 3-hydroxy-3-methylglutaryl-coenzyme A reductase and acyl coenzyme A: cholesterol O-acyltransferase in rats^{28,29}. Furthermore, a decrease in plasma low-density lipoprotein levels and hepatic cholesterol levels in rabbits fed a high-cholesterol diet has been observed³⁰.

Nephrotoxicity

Nephrotoxicity is an intrinsic adverse effect of certain anticancer drugs. Anticancer drugs have a narrow therapeutic index and therapeutic dose of such drugs usually produces significant nephrotoxicity. The dosage used in clinical trials, mostly, the maximum tolerated dose determined during phase I drug evaluation. But even greater toxicity is expected during curative therapy than during palliative therapy. But cancer patients often exhibit excretory reduced organ function. Modulation of pharmacokinetics and pharmacodynamics of these drugs in cancer patients is therefore necessary in order to improve tolerance³¹. Cancer patients are particularly liable for the development of renal abnormalities. Interestingly, patients with renal abnormalities who have undergone kidney transplantation are at high risk for malignancy. Clinical symptoms of renal involvement are diverse and harmful. In spite of the recent advances in understanding the mechanism of anticancer drug and nephrotoxicity, prevention still relies on drug dosage decrease. Hence an active screening of drugs for renal abnormalities for patients treated with anticancer drugs is still a major field of research³².

METHODS AND MATERIALS

Chemicals

All chemicals used were of analytical grade. Cisplatin, 5-¹ – Dithiobis 2-nitrobenzoic acid (DTNB) purchased

from sigma, St. Louis, M.O. USA. n-Butanol, xylene, silicagel, petroleum ether, ethanol, thiobarbituric acid (TBA), Trichloro acetic acid and (TCA), disodium hydrogen phosphate, Sodium dihydrogen phosphate were obtained from S.d fine chemicals, India. Sodium nitroprusside Sulphanilamide, O-phosphoric acid, naphthylethylene diamedihydro chloride (EDTA), Pyridine, haematoxylene purchased from Merck (INDIA). Methanol, Sulphuric acid (H₂SO₄), Isopropanol, Celite, Acetic acid, Petroleum ether (40-60⁰), Charcoal, formamide purchased from S.d fine chemicals (INDIA).

Instruments

Purity of the compounds was checked by thin layer chromatography (TLC) using glass plates coated with silica gel G, and spots were detected by iodine vapours. Melting points were determined in open capillaries on a Tempo apparatus, and are Uncorrected. Samples were send to IICT, Hyderabad to obtain IR, NMR and MASS IR spectra.

Plant material

The orange peels were collected from local juice centers of tirupathi, hesperetin is isolated from orange peels in Tirupathi.

Isolation of hesperetin

Hesperetin (HTN) was isolated by acid degradation of hesperidin (HDN). 1 gram hesperidin and 20 ml ethylene glycol containing 1 ml sulfuric acid were heated on a steam bath for 40 min. The clear yellow solution is poured into 50 ml water. The precipitated hesperetin was filtered on a Buchner funnel and washed with water. Crystallized from ethanol gives crystals (yield: 350 mg).

Selection of animals

All animal experiments were performed in accordance with our Institutional Animal Ethics Committee and by the animal regulatory body of the government (Regd.No.1629/PO/A/12/CPCSEA). Albino rats of either sex obtained from Ghosh Enterprises, Kolkata were used in the study. Under controlled environment the animals were placed six per cage at a temperature of 22 ± 2°C with 12h light / dark cycle. Rats were fed with standard pellet diet (Ratan Brothers, Hyderabad), and water ad libitum. Animals were kept for twelve days in laboratory for habituation.

Table 1
Preliminary phytochemical screening

S.no	Name of the constituent	Chemical test	Result
1	Carbohydrates	Molisch's test	-ve
2	Proteins and free amino acids	Biuret test	-ve
3	Flavonoids	Shinoda's test	+ve
4	Alkaloids	Dragand roff's test	-ve
5	Saponins	Foam test	-ve

EXPERIMENTAL PROCEDURE

The Effect of Hesperetin on cisplatin induced nephrotoxicity

The animals were divided in to five groups, each group consists of six animals (n=6)

- The Animals of Group – I_b received 1% Carboxy Methyl Cellulose (CMC) in distilled water for 10 days. The animals of group – II_b received Cisplatin (6mg/kg body weight i.p) on 5th day.
- The animals of Group-III_b received the low dose of Hesperetin (20 mg/kg body) suspended in the vehicle for 10 days.

- The animals of group – IV_b received the high dose of Hesperitin (40 mg/kg bd wt) suspended in the vehicle for 10 days.
- The animal of group –V_b received only hesperitin suspended in the vehicle for 10 days. In addition to this the animals of groups III_b, IV_b were co-administered with cisplatin (6mg/kg body weight i.p) on day 5. Group – I_b received normal saline instead of cisplatin.

Assessment of nephro-protector activity

On the day 9, urine was collected with the help of metabolic cages and the urine samples were subjected for estimation of urinary functional parameters. The animals were sacrificed by cervical decapitation and blood samples were collected by cardiac puncture and were used for estimation of Blood urine nitrogen (BUN, Di acetylmonooxime method), Serum creatinine (SC, Alkaline Picrate method). In kidney tissue Malondialdehyde levels were estimated.

Table 2
Effect of Hesperitin on normal rats

S NO	PARAMETER	GROUP I	GROUP IV
	Treatment (mg/kg)	1% carboxy methyl cellulose throught the treatment	Hesperitin given throught the period
1	BUN(mg/dl)	24.56±0.15	26.75±1.5
2	SC(mg/dl)	0.69 ± 0.16	0.71±0.01
3	U _{TP} (mg/24hrs)	6.20±0.33	7.25±0.12
4	Cl _{cr} (ml/hr/100g bd.wt)	18.90±1.3	18.98±1.9
5	LPO(n moles/gm)	73.56±2.16	72.06±2.6

Table 3
Effect of isolated compound of hesperitin on cisplatin induced nephrotoxicity

Group	Treatment (mg/kg)	BUN (mg/dl)	SC (mg/dl)	UTP(mg/24hrs)	Clcr (ml/hr/100g bd.wt)	LPO (n moles/g)
I	Normal (1%carboxy methyl cellulose)	26.76±1.5	0.72±0.2	7.8±0.33	19.80±1.6	11.7±2.6
II	Cisplatin (6mg/kg)	62.2±2.3	2.3±1.6	19.9±1.06	5.13±2.6	14.7±0.3
III	Low dose (200mg/kg)	28.02±2.90	1.6±0.9	6.8±0.6	11.7±1.4	14.1±0.5
IV	High dose (400mg/kg)	29.02±2.9	1.4±0.1	6.3±0.33	10.6±1.56	12.1±0.9

Statistical analysis

The results are expressed as mean±SEM and the data analysed using one way analysis of variance followed by post hoc Student-Keuls test using SPSS computer software for *in vivo* studies. Statistical significance was set at P≤ 0.05.

RESULTS

HESPERTIN

(S)-2,3-dihydro-5,7-dihydroxy-2-(3-hydroxy-4-methoxyphenyl)-2-methylchromen-4-one

It was obtained as white needles from petroleum ether and methanol fraction. M.P ranges between 226 to 228°C. The molecular formula C₁₆H₁₄O₆ was established by Mass spectral data and elemental analysis.

IR data (cm⁻¹)


In IR spectra the characteristic bands were observed. In IR spectra the characteristic bands were observed at 3472.92 cm⁻¹ (phenolic OH), a band at 2917.20 cm⁻¹ for C-H present in OCH₃, C-O-C band was observed at 1203 cm⁻¹, a band at 1647.76 cm⁻¹ (C=O).

¹H NMR

In ¹H NMR spectra signals at δ 4.2 (d, 1H) and δ 5.3 (d, 1H) for phenolic hydroxyl are present at 5 and 3¹ positions respectively. Signal at δ 3.7 (m, 3 H) for OCH₃ group present at 4¹ position. A signal at δ 3.12 (m, 3H) for proton is present at 3 position. Signals at δ 5.8, 5.34, 5.98 (m, 3H) protons present at 5, 2, 8 positions respectively.

Mass spectra

Mass spectra showed characteristic fragmentation peaks 302 (M⁺), 179 (M-133), 150 (M-152), 137 (M-165), 124 (M-188).Based on above physical and spectral data the isolated compound was confirmed as Hesperitin.



3.1 Effect of hesperitin on normal rat kidney

Effect on Serum markers

With the administration of hesperitin to group II_b animals there was no significant changes were observed in BUN and SC levels (Group II 26.75±1.5 and 0.71±0.01) when compared with normal group I_b animals (24.56±0.15 and 0.69 ± 0.16).

Effect on Urinary parameters

Urinary total proteins (U_{TP}) were expressed in mg/24hrs. Animals which received hesperitin showed almost similar values (7.25±0.12) to that of normal control group I_b animals (6.20±0.33). There were no significant changes in Cl_{cr} levels hesperitin treated animals (18.98±1.9) when compared with normal group I_b animals (18.90±1.3).

Effect on LPO activity

The animals which received only hesperitin on oral administration there was no significant changes observed in LPO (72.06±2.6) when compared with normal Group-I_b animals (73.56±2.16)

Effect on antioxidant enzymes

With the oral administration of only hesperitin there was no significant changes observed in GSH, CAT, SOD values (22.01±0.10, 1.66±0.15 and 104.5±5.07 of GSH, CAT and SOD respectively) when compared with normal group I_b animals (22.14±0.13, 1.65±0.12, 105.8±6.48).

3.2 Effect of Hesperitin on cisplatin-induced nephrotoxicity**Effect on Serum markers.**

Animals which received cisplatin (6mg/kg) alone showed significant elevated levels of BUN (62.2±2.3) when compared to group Ib animals (26.76±1.5). Animals which belong to Gr III_b (20 mg/kg) and Gr IV_b (40 mg/kg) has exhibited dose dependent protection (i.e., Gr III_b 28.02±2.90; Gr IV_b 29.02±2.9). The SC levels were increased in Gr II_b which received cisplatin (2.3±1.6) when compared with normal control animals (0.72±0.2). Animals treated with hesperitin of 20 mg/kg Bd.wt showed moderate protection (1.6±0.9) and animals treated with 40mg/kg Bd.wt showed significant protection (1.4±0.1).

Effect on Urinary parameters

Animals administered with cisplatin excreted high amount of U_{TP} (19.9±1.06) when compared with normal group I_b animals (7.8±0.33) where as animals belonging to Gr III_b (20mg/kg) and Gr IV_b (40 mg/kg) reversed the effect caused by cisplatin (i.e., 6.8±0.6, 6.3±0.33). The animals received cisplatin alone exhibited decreased levels of Cl_{cr} (5.13±2.6) when compared with normal animals (19.80±1.6). On oral administration of hesperitin fraction showed significant increase in Cl_{cr} and the values are Gr III_b animals 20mg/kg bd wt (11.7±1.4) and Gr IV_b animals 40 mg/kg bd wt (10.6±1.56) respectively.

3.3 Effect on LPO activity:

The animals received cisplatin alone increased levels of LPO (14.7±0.3) when compared with normal Group-I_b animals (11.7±2.6). On oral administration of hesperitin to the groups of III_b (20mg/bd wt), IV_b (40mg/kg Bd wt) showed decreased levels of LPO (14.1±0.5), (12.1±0.9)

3.4 Effect on antioxidant enzymes

Animals which received cisplatin alone increased levels of decreased levels of GSH (76.5±2.3), CAT (7.10±0.57), SOD (23.1±1.15) when compared with normal group I_b animals (143.14±1.6, 1.25±0.12 and 105.8±6.48, of GSH, CAT and SOD respectively). On oral administration of hesperitin fraction to animals belonging to Gr III_b (20 mg/kg Bd wt) and Gr IV_b (40 mg/kg Bd wt) showed increased levels of GSH, CAT, SOD indicating moderate protection at 20 mg/kg bd wt (85.42±3.7, 12.5±1.50 and 43.1±3.2) and significant protection at 40 mg/kg bd wt (89.13±1.9, 14.0±1.0 and 68.52±9.0)

DISCUSSION

Cisplatin is an antitumor drug; it has been successful in the bladder, lung, head, neck, cervical especially testicular and ovarian cancers. The precise mechanism of cisplatin-induced nephrotoxicity has not been elucidated, but it has been suggested that the oxygen free radicals play an important role. Cisplatin-induced nephrotoxicity is related to increase in lipid peroxide levels in kidney. Reports also suggest that there is an involvement of nitric oxide which induces the nephrotoxicity by cisplatin. Tsulsumishitha and coworkers reported the involvement of H₂O₂ in cisplatin-induced nephrotoxicity in outer medullary cortical tubule (OCMT) cells. Number of studies has shown that reduction of the glomerular filtration rate is common in patients receiving cisplatin treatment. Moreover, cisplatin-induced nephrotoxicity in humans is secondary to the accumulation of this drug in the tubular epithelial cells from segment of the proximal kidney tubules. However the generation of oxygen free radicals in tubular cells has been proposed as an important pathogenic process. Various data indicate that cisplatin induces oxidative stress, lipid peroxide and DNA damage. Flavonoids are the polyphenolic compounds are found in fruits and vegetables, nuts and seeds as well as most types of tea and red wine. Flavonoids show antioxidant and anti inflammatory activity. In present study, the effect of cisplatin is similar to those previously described i.e., elevated serum urea, serum creatinine levels, reduced creatinine clearance, increased urinary protein excretion. Previous reports also evidenced that cisplatin exerted its nephrotoxic effects through LPO. In the present study, cisplatin induced accumulation of lipid peroxides and depletion of glutathione and its related antioxidant enzymes in kidney may play the critical role of oxidative stress in cisplatin nephrotoxicity. A Cisplatin-induced alteration in lipid peroxides is markedly improved by HDN and HTN. The mechanism of cisplatin nephrotoxicity is related to depletion of the antioxidant defence system. In present study, GSH depletion by cisplatin is in agreement with other reports pertaining to cisplatin-induced renal GSH depletion. The results of the present study showed CAT, SOD activities were significantly decreased in the cisplatin treated animals. hesperitin treated animals significantly improved the levels of GSH, CAT and SOD respectively compared to cisplatin treated group The HDN, HTN showed the protective activity against the cisplatin induced nephrotoxicity. The HDN showed more

prominent action when compared with HTN. HDN is a glycoside and HTN is the aglycon of HDN. Presence of sugar moiety in hesperidin may play a critical role in transportation and fast on- set of action. The inhibition of antioxidant enzyme activity in cisplatin nephrotoxicity and prevention of this altered in rats treated with hesperidin and hesperitin support the rationale for the use of antioxidants to ameliorate cisplatin nephrotoxicity. In conclusion, the results provide HDN, HTN are attenuates the nephrotoxicity of cisplatin in rats. The results provide further insight into the mechanisms of cisplatin-induced nephrotoxicity and confirm the antioxidant potential of hesperitin.

CONCLUSION

Hesperitin is a bioflavonoid, to be more specific a flavones. It contains Aglycon moiety and good anti-oxidant properties. It also enhances lipid metabolism and regulate apolipoprotein B secretion by HepG2 cells, possibly through inhibition of cholesterol ester synthesis. Due to its high anti-oxidant properties there is chance of this compound in treating diseases like Diabetes.

CONFLICT OF INTEREST

Conflict of interest declared none.

REFERENCES

- Green Med info LLC.9 Evidence-Based Medicinal Properties of Oranges [Internet]. Sayer ji, founder.2013 [updated 2013 June 15th; cited 2016 August 4th] Available from: <http://www.greenmedinfo.com/blog/9-evidence-based-medicinal-properties-oranges>
- Baseline of health organization. Orange Peels – Eating orange peels/herbal library [Internet] jon barron. Available from: <http://jonbarron.org/herbal-library/foods/eating-orange-peels>.
- Wasnik DD, Tumane PM and Meshram VG, Comparative study of antibacterial activity of peel extracts of Citrus aurantium l. (bitter orange) and Citrus medica l. (lemon) against clinical isolates from wound infection; Int J Pharm Bio Sci 2014 Jan; 5(1):382 - 387
- Tripoli E, Guardia ML, Giammanco S, Majo DD, Giammanco M. Citrus flavonoids: Molecular structure, biological activity and nutritional properties: A review. Food Chemistry 2007; 104: 466-479.
- De Groot H, Raven U. Tissue injury by reactive oxygen species and the protective effects of flavonoids. Fundam Clin Pharma Col 1998; 12:249-255.
- Slade D, Ferreira D, Marais JPJ. Circular dichroism, a powerful tool for the assessment of absolute configuration of flavonoids. Phytochemistry 2005. 66: 2177-2215.
- Hotta H, Nagano S, Ueda M, Tsujino Y, Koyama J, Osakai T. Higher radical scavenging activity of polyphenolic antioxidants ascribed to chemical reactions following oxidation. Biochimica et Biophysica Acta 2002; 1572: 123-132.
- Williams RJ, Spencer JPE, Rice-Evans C. Serial review: Flavonoids and isoflavonones (Phytoestrogens): Absorption, Metabolism and Bioactivity. Free Radical Biology and Medicine 2004; 36: 838-849.
- Fernandez SP, Wasowski C, Loscalzo LM, Granger RE, Johnston GAR, Paladini AC, et al. Central nervous system depressant action of flavonoid glycosides. European Journal of Pharmacology 2006; 539: 168-176.
- Heim KE, Tagliaferro AR, Bobliya DJ. Flavonoids antioxidants: Chemistry, metabolism and structure-activity relationships. The Journal of Nutritional Biochemistry 2002; 13: 572-584.
- Hollman PCH, Katan MB. Dietary Flavonoids: Intake, Health Effects and Bioavailability. Food and Chemical Toxicology 1999; 37: 937-942.
- Cushnie TPT, Lamb AJ. Antimicrobial activity of flavonoids. International Journal of Antimicrobial Agents 2005; 26: 343-356.
- Heim KE, Tagliaferro AR, Bobliya DJ. Flavonoids antioxidants: Chemistry, metabolism and structure-activity relationships. The Journal of Nutritional Biochemistry 2002; 13: 572-584.
- Peterson J, Dwyer MSJ, RD, DSc. Flavonoids: Dietary occurrence and biochemical activity. Nutrition Research 1998; 18: 1995-2018.
- Tsuchiya H. Structure-dependent membrane interaction of flavonoids associated with their bioactivity. Food Chemistry 2010; 120: 1089-1096.
- Rijke, ED, Out P, Niessen WMA, Ariese, F, Goojer C, Brinkman UAT. Analytical separation and detection methods for flavonoids. Journal of Chromatography A 2006; 1112: 31-63
- Ross JA, Kasum CM. Dietary flavonoids: bioavailability, metabolic effects and safety. Annu Rev Nutr. 2002; 22:19–34.
- Harborne JB. The flavonoids- Advances in Research Since 1980. ed 1. London: Chapman and Hall; 1988.
- Middleton E. The flavonoids. Trends Pharmacol Sci 1984; 5:335-338.
- So FV, Guthrie N, Chambers AF, Moussa M, Carroll KK. Inhibition of human breast cancer cell proliferation and delay of mammary tumorigenesis by flavonoids and citrus juices. Nutr Cancer 1996; 262:167 - 81.
- Yang M, Tanaka T, Hirose Y, Deguchi T, Mori H, Kawada Y. Chemopreventive effects of diosmin and hesperidin on N-butyl-N-(4 hydroxybutyl)nitrosamine-induced urinary-bladder carcinogenesis in male ICR mice. Int J Cancer 1997; 73:719 - 24.
- Tanaka T, Makita H, Kawabata K, Mori H, Kakumoto M, Satoh K, et al. Chemoprevention of azoxymethane induced rat colon carcinogenesis by the naturally occurring flavonoids, diosmin and hesperidin. Carcinogenesis 1997;18:957 - 65.
- Miyagi Y, Om AS, Chee KM, Bennink MR. Inhibition of azoxymethane-induced colon cancer by orange juice. Nutr Cancer 2000;36:224 - 9.

24. Miyake Y, Yamamoto K, Tsujihara N, Osawa T. Protective effects of lemon flavonoids on oxidative stress in diabetic rats. *Lipids* 1998;33:689 - 95.
25. Van Acker FA, Tromp MN, Haenen GR, van der Vijgh WJ, Bast A. Flavonoids can replace a-tocopherol as an antioxidant. *FEBS Lett* 2000; 473:145 - 8.
26. Jeon SM, Bok SH, Jang MK, Kim YH, Nam KT, Jeong TS, et al. Comparison of antioxidant effects of naringin and probucol in cholesterol-fed rabbits. *Clin Chim Acta* 2002; 317:181 - 90.
27. Borradaile NM, Carroll KK, Kurowska EM. Regulation of HepG2 cell apolipoprotein B metabolism by the citrus flavanones hesperetin and naringenin. *Lipids* 1999; 34:591 - 8.
28. Lee SH, Jeong TS, Park YB, Kwon YK, and Choi MS, Bok SH. Hypocholesterolemic effect of hesperetin mediated by inhibition of 3-hydroxy-3-methylglutaryl coenzyme A reductase and acyl coenzyme A: cholesterol acyltransferase in rats fed high-cholesterol diet. *Nutr Res* 1999; 19:1245 - 58.
29. Lee SH, Park YB, Bae KH, Kwon YK, Lee ES, Choi MS. Cholesterol-lowering activity of naringenin via inhibition of 3-hydroxy-3-methylglutaryl coenzyme A reductase and acyl coenzyme A: cholesterol acyltransferase in rats. *Ann Nutr Metab* 1999; 43:173 - 80.
30. Kurowska EM, Borradaile NM, Spence JD, Carroll CC. Hypocholesterolemic effects of dietary citrus juices in rabbits. *Nutr Res* 2000;20:121 – 9
31. Zheng X, Wise.S.M, Cristini.V. Nonlinear simulation of tumor necrosis, neo-vascularization and tissue invasion via an adaptive finite-element / level-set method. *A bulletin of Math. Biol.* March (2005). [cited 2015 feb] 67: 211. Available from: <http://dx.doi.org/10.1016/j.nephro.2004.12.001>. DOI: 10.1016/j.bulm.2004.08.001
32. Patricia. A, Kaba.SP, Marilia M Trierweiler, Elio H Shinohara. Osteoblastic metastasis from breast affecting the condyle misinterpreted as temporomandibular joint disorder. *Indian J Cancer* [serial online] 2011 [cited 2016 oct 7]; 48:252-3. Available from: <http://www.indianjcancer.com/text.asp?2011/48/2/252/82877>. DOI:10.4103/0019-509X.82877.