



## COLORIMETRIC DETERMINATION OF VITAMIN C IN FRESH AND DILUTE FRUIT JUICES AND EFFECT OF THERMAL EXPOSURE ON CONCENTRATION AT VARIOUS STAGES

**N.MADHAVI\*<sup>1</sup>, B.JYOTHI<sup>2</sup>**

*<sup>1</sup>P.G department of chemistry, J.K.C College, Guntur*

*<sup>2</sup>Department of Chemistry, Swarna Bharathi Institute of Science and Technology, Khammam, Telangana, India*

### ABSTRACT

Among all vitamins, vitamin C plays a crucial role in maintaining good health. In the present study ten types of fruits are analyzed to compare ascorbic acid levels and the effect of boiling on its quantity is also determined by colorimetric method. It is concluded that its levels are gradually decreased with increase of boiling time. Cheaply and mostly available fruits have higher concentration of ascorbic acid than other fruits. In *Psidium guajana*, *Citrus aurantium* and *Phyllanthus emblica* the concentration of vitamin C is more. It is concluded that the concentration of ascorbic acid is more in semi-ripened stage of the fruit than ripened and unripened stages of *Citrus limon*, *Citrus aurantifolia* and *Psidium guajana*. But it is more in unripened fruits of *Citrus medica* and *Citrus aurantium*. In *Psidium cattleianum* and *Psidium littorale* ascorbic acid concentration is more in fully ripened stage.

**KEYWORDS:** ascorbic acid, fruits, colorimetry, ripening, boiling.



\*corresponding author



**N.MADHAVI\*<sup>1</sup>**

P.G department of chemistry, J.K.C College, Guntur

## INTRODUCTION

Proper health is the basic thing for survival of human life. Fruits protect us from many diseases. Citrus fruits contain major quantity of ascorbic acid<sup>1</sup>, which is an essential nutrient<sup>2</sup> and structurally relates to glucose. It is a weak sugar acid<sup>3,4</sup>, found in many vegetables and fruits. Vitamin C helps to make collagen, absorbs iron<sup>5</sup> and neutralizes free radicals.<sup>6</sup> It is used as food additive<sup>7</sup>, works as a co-factor in several enzymatic reactions<sup>8</sup>, fight against infectious and chronic diseases<sup>9-13</sup> and strengthens immune system.<sup>14</sup> Its quantity decides the nutritional quality of the fruit.<sup>15,16</sup> During thermal exposure its stability is lost.<sup>17,18</sup> Our body is unable to synthesize vitamin C<sup>19</sup> but it has the role in preventing oxidative damage to DNA<sup>20,21</sup> and proteins in our body. Deficiency of this water soluble<sup>22</sup> and antioxidant<sup>23,24</sup> vitamin causes scurvy<sup>25,26</sup> which is identified by bleeding gums, anaemia<sup>27</sup>, fatigue, depression, mal wound healing<sup>28</sup> and scaly skin. Ripening is the process by which fruits attain their desirable flavor, quality, color, palatable nature and other textural properties. It is associated with change in conversion of starch to sugar. During this process the fruits emit ethylene along with increased rate of respiration.

## MATERIALS AND METHODS

### Sample collection

Lime, lemon, citron, sour orange, Indian goose berry, star goose berry, sweet lime, red guava, green guava, white guava fruits of unripened, semi ripened and fully ripened are collected from local fruit gardens of Madhira mandal of Khammam district during winter season.

### Sample preparation

Fruits are washed thoroughly with distilled water, manually sliced and pulped using a blender. Sample is divided into six portions, weighing five grams each and labelled as samples A-E. All these samples are diluted and exposed to heat treatment at 50°C for 5, 10, 15, 20, 25 and 30 minutes respectively. Then absorbance is measured for both fresh samples and diluted samples using colorimeter.

### Analysis

The processed samples are subjected to the following analysis immediately after sample preparation for vitamin C determination. 20ml oxalic acid solution, 0.2ml of 0.01 % methylene blue solution and 1ml of acetate

buffer of pH 4.2 are taken in a 50ml beaker. To this fresh and heated ripened sample at 50°C of 1ml solution is added separately and measured for absorbance periodically after 5, 10, 15, 20, 25 and 30 minutes of boiling. Absorbance of each solution is read using Roy instruments digital photo colorimeter (RI\_506 C) of infra instruments private limited at wavelength 540nm. Repeated the same procedure for semi ripened and unripened fruits of diluted and fresh juices.


## RESULTS AND DISCUSSION

In the calibration curve, the absorbances of standard vitamin C at various concentrations C are plotted. The concentration of vitamin C after boiling times of 5, 10, 15, 20, 25 and 30 minutes at 50°C is plotted and it is found to be more in Psidium guajana, Citrus aurantium and Phyllanthus emblica. The concentration of original sample is prominent than that of boiled samples. The retention of vitamin C is often used as an indicator for the overall nutrient reduction of food products because it is highly sensitive to oxidation and leaching in water soluble media during heating. Higher temperatures does not retain vitamin C, causing it to leach out of the food into the water and also to be oxidized, first to dehydroascorbic acid and then to diketogulonic acid. This last compound has no vitamin C activity at all and is irreversible. It is better to maintain or store samples below the room temperature. Its concentration is decreased during fruit ripening in some fruits. Feeble vitamin C is easily deteriorated during boiling and finally depleted as the boiling time is increased. The present work revealed that the concentration of ascorbic acid is more in semi-ripening stage of the fruit than the ripened and unripened stages of lemon, lime, and green guava. It is noticed to be more in unripened fruits of sour orange (Fig 2) and citron (Fig 1). In fully ripened fruits of red guava (Fig 5) and white guava its quantity is more. So fruits should be consumed before processing to attain more concentration of ascorbic acid. It is diminished more in first 5 and 10 minutes of boiling time, and there is a regular decrease afterwards. In some fruits after periodical heating up to 30 minutes its concentration is completely reduced to zero. Dilution of heated samples causes further decrease of its quantity. The vitamin C concentration in following fruits is as follows: Green Guava > Indian Goose Berry > White Guava = Sour Orange > Red Guava > Lemon > Lime > Citron > Sweet Lime > Star Gooseberry. (Graph 4 to graph 24).

**Table 1**  
**Vitamin-C concentration of fruits in various stages**

S no	Sample	vitamin c concentration in mg/25ml					
		Original			Diluted		
		Unripened	Semi Ripened	Fully Ripened	Unripened	Semi Ripened	Fully Ripened
1	Lemon[Citrus limon]	56	81.1	37.8	8.9	25	15.7
2	Lime[Citrus aurantifolia]	56	65.5	37.8	24	37.8	16
3	Citron[Citrus medica]	47	43.7	33.4	28.4	24.3	22.7
4	Sour orange [Citrus aurantium ]	112	93.7	65.4	81.1	37.8	25.4
5	Indian Goose berry [Phyllanthus emblica]	92	123	104	78	112	91
6	Star goose berry [Phyllanthus acidus]	17	24	20	12	18	14
7	Sweet lime [Citrus sinensis]	37.8	33.4	28.4	18.9	16.2	12.6
8	Red guava [Psidium cattleianum sabine]	85	105.5	107	28.4	43.7	82
9	Green guava [Psidium guajana]	113	127	90	73.4	81.1	33.4
10	white guava [Psidium littorale]	83.2	92.3	112	29.7	57	72

**Graph 1**  
**Vitamin-C concentration of fruits in various stages**



Bars in the graph represent the absorbances of fruits in original and dilute samples in the three ripening stages of fruits

**Table 2**  
**Vitamin-C concentration in original and diluted juices of fruits at various processing times**

FRUIT	FRUIT STAGE	SAMPLE CODE	PROCESSING TIME(min)	mg/25ml	
				ORIGINAL	DILUTED
LEMON	UNRIPENED	A	5	17.5	9
		B	10	15.39	7
		C	15	15.31	6
		D	20	9.8	5
		E	25	6.9	4
		F	30	5	3
	SEMI RIPENED	A	5	56	14.2
		B	10	24	10.3
		C	15	23.5	7.8
		D	20	17.5	6.5
		E	25	16.45	5.8
		F	30	10.78	4
	RIPENED	A	5	14.2	4.9
		B	10	11.1	4.3
		C	15	7.5	4
		D	20	6.7	0
		E	25	5.7	0
		F	30	4.5	0
LIME	UNRIPENED	A	5	33.4	22.7
		B	10	24	16
		C	15	18.9	8.9
		D	20	16.3	6.2
		E	25	14	5.2
		F	30	11.2	4.4
	SEMI RIPENED	A	5	43.7	23.2
		B	10	33.4	17.5
		C	15	25.8	14.5
		D	20	24	12.4
		E	25	22.7	10.33
		F	30	18.9	8
	RIPENED	A	5	24	14.5
		B	10	22.7	11.4
		C	15	18.9	10
		D	20	16.4	6
		E	25	12.6	4.2
		F	30	11.1	3.9
SOUR ORANGE	UNRIPENED	A	5	63	29
		B	10	43.7	22
		C	15	22.7	17

		D	20	16.5	12
		E	25	16.2	9
		F	30	14	6.7
		A	5	43.7	24
		B	10	33.4	20
	SEMI RIPENED	C	15	25.8	15.6
		D	20	23	10.33
		E	25	18.9	7.8
		F	30	14.2	6
		A	5	28.4	20
		B	10	25.8	17.7
	RIPENED	C	15	22.7	16.3
		D	20	19	13
		E	25	16.2	8
		F	30	8.7	4.9
		A	5	43.7	24
		B	10	33.4	23.5
	UNRIPENED	C	15	25.8	14.2
		D	20	23	11.8
		E	25	22.7	10.33
		F	30	18.9	7.9
		A	5	33.4	22.8
		B	10	25.8	16.8
	SEMI RIPENED	C	15	18.2	13.8
		D	20	17.5	9
		E	25	16	7.5
		F	30	10	4.9
		A	5	25	18.2
		B	10	22.7	15.4
	RIPENED	C	15	17.9	13.8
		D	20	16	9
		E	25	12	6.9
		F	30	10.33	4.9
		A	5	72.5	52.4
		B	10	63.4	43.2
	UNRIPENED	C	15	52.3	32
		D	20	42	24
		E	25	35	14
		F	30	23	12
		A	5	97.4	62.3
		B	10	77.4	52.4
	SEMIRIPENED	C	15	65.8	37.8
		D	20	54	25.7
		E	25	45.3	20
		F	30	32	16
		A	5	82.3	60.4
		B	10	75	50.7
	RIPENED	C	15	62	33
		D	20	53	26
		E	25	42.5	16.5
		F	30	29.7	14
		A	5	15.6	10.7
		B	10	14.2	8.4
	UNRIPENED	C	15	12.6	6.8
		D	20	8.7	4.5
		E	25	5.7	0
		F	30	4.5	0
		A	5	20.2	16.4
		B	10	17.2	14.7
	SEMIRIPENED	C	15	13	9
		D	20	9.5	6
		E	25	8.9	3
		F	30	6	0
		A	5	18	12.7
		B	10	16	10.9
	RIPENED	C	15	13	8
		D	20	9.5	5
		E	25	9	0
		F	30	5	0
		A	5	33.4	17.4
		B	10	28	16.3
	UNRIPENED	C	15	24	14.7
		D	20	17	10.33
		E	25	15	8.7
		F	30	9	5.2
		A	5	24	8.7
		B	10	20	7.6
	SEMIRIPENED	C	15	17.2	6
		D	20	13.5	4.9
		E	25	8.9	0
		F	30	8	0
		A	5	30	13.2
	RIPENED	B	10	28	9.5

		C	15	23.5	8.7	
		D	20	16.4	7.6	
		E	25	14	6.3	
		F	30	8.9	4.9	
		A	5	55	38	
		B	10	34	23	
	UNRIPENED	C	15	28.9	18.7	
		D	20	16.2	7.2	
		E	25	13	6	
		F	30	10	5	
		SEMIRIPENED	A	5	68.4	48.3
			B	10	34	25
	C		15	20	19	
	D		20	16.2	8.7	
	E		25	13	7.1	
	F		30	10.33	4.9	
	RIPENED	A	5	81	54	
		B	10	62.7	44.4	
		C	15	48.9	24.8	
		D	20	26.6	17	
		E	25	16.2	11	
		F	30	12	6	
RED GUAVA	UNRIPENED	A	5	56	28.4	
		B	10	32.7	15.2	
		C	15	18	12.6	
		D	20	15.6	8.7	
		E	25	15	7.9	
		F	30	12.6	4.9	
	SEMI RIPENED	A	5	66	33.4	
		B	10	43.7	25	
		C	15	25.8	20	
		D	20	25	17	
		E	25	24	15	
		F	30	22.7	11	
	RIPENED	A	5	43.7	19	
		B	10	24	15.2	
		C	15	17.5	11.4	
		D	20	15	9.2	
		E	25	13.8	6.2	
		F	30	11.2	4.9	
GREEN GUAVA	UNRIPENED	A	5	35	28	
		B	10	22.7	16.3	
		C	15	18.9	14	
		D	20	17.51	11.5	
		E	25	14	7.1	
		F	30	9.5	6	
	SEMI RIPENED	A	5	48.4	25	
		B	10	25.8	12	
		C	15	18.2	10	
		D	20	18	8.7	
		E	25	14.2	8	
		F	30	11.3	7.1	
	RIPENED	A	5	65.8	48.2	
		B	10	42.4	37.51	
		C	15	32.7	24.9	
		D	20	18.9	15	
		E	25	17.51	8.5	
		F	30	14.2	9	
WHITE GUAVA	UNRIPENED	A	5	65.8	48.2	
		B	10	42.4	37.51	
		C	15	32.7	24.9	
		D	20	18.9	15	
		E	25	17.51	8.5	
		F	30	14.2	9	
	SEMI RIPENED	A	5	65.8	48.2	
		B	10	42.4	37.51	
		C	15	32.7	24.9	
		D	20	18.9	15	
		E	25	17.51	8.5	
		F	30	14.2	9	
	RIPENED	A	5	65.8	48.2	
		B	10	42.4	37.51	
		C	15	32.7	24.9	
		D	20	18.9	15	
		E	25	17.51	8.5	
		F	30	14.2	9	



Figure 1  
Citron [*Citrus medica*]



**Figure 2**  
**Sour orange [*Citrus aurantium*]**



**Figure 3**  
**Citron [*Citrus medica*]**



**Figure 4**  
**Sweet lime [*Citrus sinensis*]**




**Figure 5**  
**Red guava [*Psidium cattleianum sabine*] grinded**




**Figure 6**  
*Red guava [Psidium cattleianum sabine] fresh cut pieces*

**Graph 2**  
*Standard curve of vitamin – C at various concentrations*




*Absorbances are plotted at various concentrations of ascorbic acid. As the concentration is decreasing absorbance also decreases.*

**Graph3**  
*Standard curve of vitamin – C at various boiling times*




*Absorbances are plotted at different boiling times. Absorbance of vitamin –c is decreased as the processing time is increased.*


**Graph4**  
*Lemon vitamin – C concentration original*



**Graph5**  
*Lemon vitamin – C concentration diluted*




**Graph6**  
*Lime vitamin – C concentration original*







**Graph7**  
*Lime vitamin – C concentration diluted*




**Graph8**  
*Citron vitamin – C concentration original*




**Graph9**  
*Citron vitamin – C concentration diluted*




**Graph10**  
**Indian goose berry vitamin – C conc. Original**




**Graph11**  
**Indian goose berry vitamin – C conc. diluted**




**Graph12**  
**Star goose berry vitamin – C conc. original**




**Graph13**  
*Star goose berry vitamin – C conc. diluted*




**Graph14**  
*Sour orange vitamin – C conc. Original*




**Graph15**  
*Sour orange vitamin – C conc. Diluted*




**Graph16**  
Sweet lime vitamin – C conc. Original




**Graph17**  
Sweet lime vitamin – C conc. diluted




**Graph18**  
Red guava vitamin – C conc. original




**Graph19**  
*Red guava vitamin – C conc. Diluted*




**Graph20**  
*Green guava vitamin – C conc. Original*




**Graph21**  
*Green guava vitamin – C conc. diluted*



**Graph22**  
**White guava vitamin – C conc. Original**



**Graph23**  
**White guava vitamin – C conc. Diluted**



Among the three stages of ripening, least concentration of ascorbic acid is observed in case of star goose berry whereas maximum concentrations are observed for green guava at unripened and semi ripened stages only. In case of fully ripened white guava maximum concentration is observed. In semi ripened and ripened stages of red guava (Fig 5 and 6) the concentrations are almost similar. In case of dilute samples, the least quantity is observed for lemon at unripened stage. For sweet lime (Fig 3) at semi ripened and fully ripened stages the least concentrations are noticed. Highest concentration at unripened stage is reported for sour orange (Fig 2). In case of Indian gooseberry major concentrations are observed at both semi and fully ripened stages. In citron (Fig 1) more concentration is observed in unripened stage. For lime at semi ripened stage higher concentration is noticed. With increase in processing time also similar decrease in concentration of ascorbic acid is observed. After dilution of lemon juice taken at ripened stage during processing time after 20, 25 and 30 minutes almost the concentration of ascorbic acid is completely reduced to zero. For dilute sample of star goose berry the concentration of ascorbic acid is completely reduced to zero at all the three ripening stages.

## CONCLUSION

From the present investigation it is concluded that cheaply available guava has more concentration of vitamin C than other costly fruits. This method is more appropriate to determine the quantity of ascorbic acid. It is advisable to take fresh fruits than stored pulps and juices. Because processing of a fruit into juice and pulp involves thermal exposure which leads to deterioration of its quality.

## ACKNOWLEDGEMENT

Authors are extremely grateful to the Management, and the Principal of Swarna Bharathi Institute of Science And Technology, Khammam, Telangana, India for providing facilities in the department.

## CONFLICT OF INTEREST

Conflict of interest declared none.

## REFERENCES

1. Harris JR, editor. Subcellular Biochemistry: Ascorbic Acid: Biochemistry and Biomedical Cell Biology. Springer Science & Business Media; 2013 Nov 11.
2. Ariharan VN, Kalirajan K, Meena Devi VN, Nagendra Prasad P. An exotic fruit which forms the new natural source for vitamins-C. *Rasayan J. chemistry.* 2012;5(3):356-9.
3. Tee ES, Young SI, Ho SK, Shahid SM. Determination of vitamin C in fresh fruits and vegetables using the dye-titration and microfluorometric methods. *Pertanika.* 1988;11(1):39-44.
4. Njoku PC, Ayuk AA, Okoye CV. Temperature effects on vitamin C content in citrus fruits. *Pakistan Journal of Nutrition.* 2011;10(12):1168-9.
5. Lee SK, Kader AA. Preharvest and postharvest factors influencing vitamin C content of horticultural crops. *Postharvest biology and technology.* 2000 Nov 30;20(3):207-20.
6. Weber P, Bendich A, Machlin LJ. Vitamin E and human health: rationale for determining recommended intake levels. *Nutrition.* 1997 May 31;13(5):450-60.
7. Mai, T. T. T, Mohammed F. Ultraviolet treatment of orange juice. *Journal of innovative food science and engineering technologies.* 2004; 5: 495-502.
8. Ball GF. Bioavailability and analysis of vitamins in foods. Chapman & Hall; 1998.
9. Willett WC. Diet and health: what should we eat. *Science.* 1994 Apr 22;264(5158):532-7.
10. Heber D. Vegetables, fruits and phytoestrogens in the prevention of diseases. *Journal of postgraduate medicine.* 2004 Jun 1;50(2):145.
11. Cox BD, Whichelow MJ, Prevost AT. Seasonal consumption of salad vegetables and fresh fruit in relation to the development of cardiovascular disease and cancer. *Public health nutrition.* 2000 Mar 1;3(01):19-29.
12. Joshipura KJ, Ascherio A, Manson JE, Stampfer MJ, Rimm EB, Speizer FE, Hennekens CH, Spiegelman D, Willett WC. Fruit and vegetable intake in relation to risk of ischemic stroke. *Jama.* 1999 Oct 6;282(13):1233-9.
13. Lampe JW. Health effects of vegetables and fruit: assessing mechanisms of action in human experimental studies. *The American journal of clinical nutrition.* 1999 Sep 1;70(3):475s-90s.
14. Mapson LW, Isherwood FA. Biological synthesis of ascorbic acid: the conversion of derivatives of D-galacturonic acid into L-ascorbic acid by plant extracts. *Biochemical Journal.* 1956 Sep;64(1):13.
15. Erdurak-Kiliç CS, Uslu B, Dogan B, Ozgen U, Ozkan SA, Coskun M. Anodic voltammetric behavior of ascorbic acid and its selective determination in pharmaceutical dosage forms and some Rosa species of Turkey. *Journal of Analytical Chemistry.* 2006 Nov 1;61(11):1113-20.
16. Aabraha T, Sergawie A. Assessment of Some Selected Beverages and Fresh Edible Vegetables as Nutritional Source of Vitamin C (Ascorbic Acid) by Cyclic and Square Wave Voltammetry. *International journal of science and engineering investigations.* 2014 march;3,( 26):39-49.
17. Dr.D.Gopala Krishna, Asma Alrwahi, Ruqaya Alrhbi, Huda Alwardi, Ch.Kethani Devi. Quantitative estimation of various parameters and physicochemical properties of citrus fruits at variable temperatures. *International journal of analytical and bioanalytical chemistry.* 2015 march; 5(1):24-27.
18. Leahu A, Damian C, Oroian M, Ropciuc S. Physico-chemical parameters of fruit juices-evolution during storage. *University of Agricultural Sciences and Veterinary Medicine Iasi.* 2013 Jun.
19. Li Y, Schellhorn HE. New developments and novel therapeutic perspectives for vitamin C. *The Journal of nutrition.* 2007 Oct 1;137(10):2171-84.
20. Ghafoor S, Arain R, Yasmeen K, Ahmed F, Saeed I. Determination of ascorbic acid content of some capsicum cultivars by cyclic voltammetry performed at GCE by external standard series calibration method. *Int. J. Electrochem. Sci.* 2014 Oct 1;9:5751-62.
21. Trujillo M, Radi R. Peroxynitrite reaction with the reduced and the oxidized forms of lipoic acid: new insights into the reaction of peroxynitrite with thiols. *Archives of biochemistry and biophysics.* 2002 Jan 1;397(1):91-8.
22. Igwemmar NC, Kolawole SA, Imran IA. Effect Of Heating On Vitamin C Content Of Some Selected Vegetables. *International Journal of Scientific and Technology Research.* 2013 Nov 25;2(11):209-12.
23. Blagojević nz, vesna I, brašanac sr. Kinetics of degradation of ascorbic acid by cyclic voltammetry method. *Chem. Ind. Chem. Eng. Q.* 2015. 21 (2) 351–357.
24. Padayatty SJ, Katz A, Wang Y, Eck P, Kwon O, Lee JH, Chen S, Corpe C, Dutta A, Dutta SK, Levine M. Vitamin C as an antioxidant: evaluation of its role in disease prevention. *Journal of the American College of Nutrition.* 2003 Feb 1;22(1):18-35.
25. Esch JR, Friend JR, Kariuki JK. Determination of the vitamin C content of conventionally and organically grown fruits by cyclic voltammetry. *Int. J. Electrochem. Sci.* 2010 Oct 1;5:1464-74.
26. Murcia MA, López-Ayerra B, Martínez-Tomé M, Vera AM, García-Carmona F. Evolution of ascorbic acid and peroxidase during industrial processing of broccoli. *Journal of the Science of Food and Agriculture.* 2000 Oct 1;80(13):1882-6.
27. N.Madhavi,G.Swathi,E.Padmaja.Determination of chemical parameters of various fresh fruit juices.*International Journal of Institutional Pharmacy and Life Sciences.*2015 March 30; 5(2): 181-183.
28. N.Madhavi, D. D Pravallika. Determination of Physico chemical parameters of various fresh fruit juices. *International Journal of Organic and Bioorganic Chemistry.* 2015 March 20; 5(1):13-14.