



ANTIBACTERIAL ACTIVITY OF METHANOL CRUDE EXTRACTS OF MARINE MACRO ALGAE FROM THE THIKKODI CALICUT WEST COAST OF INDIA

Dr.R.VEERAMOHAN^{1*}, G. ADAIKALA RAJ², M.
AND V. VENKATESALU²

^{1*} Associate Professor Department of Botany, Dr. S.R.K Government Arts College, Yanam – 533 464, Pondicherry, India

² Department of Botany, Annamalai University, Annamalainagar - 608 002, Tamil Nadu, India

ABSTRACT

The antibacterial activities of methanol extracts of *Caulerpa racemosa* (Forsk), *Caulerpa sertularioides* (Gmelin) Howe, *Ulva lactuca* Linn, *Padina gymnospora* (Kutzing) Sonder., *Dictyota dichotoma* (Hudson) Lamouroux. and *Gracilaria multipartita* (Clemente) Harvey against bacterial strains viz., *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Proteus mirabilis* and *Salmonella enterica* were studied by using disc diffusion method, determination of MIC and MBC. The methanol extracts of *P. gymnospora* and *D. dichotoma* showed the highest antibacterial activity against all the bacterial strains tested. The mean zones of inhibition produced by the extracts in agar disc diffusion assays were from 6.3 to 19.5 mm. The MIC was between 125 and 500 µg/ml and MBC was between 250 and 1000 µg/ml. The highest mean zone of inhibition (19.5 mm) was observed with methanol extract of *Padina gymnospora* against *S. aureus*. The lowest MIC (125 µg/ml) and MBC (250 µg/ml) values were observed in methanol extract of *P. gymnospora* and *D. dichotoma* against *S. aureus*. These finding suggest that methanol crude extract of *P. gymnospora* have potential antibacterial activity.

KEYWORDS: Marine macroalgae, Antibacterial activity, MIC, MBC



Dr.R.VEERAMOHAN*

Associate Professor Department of Botany, Dr. S.R.K Government Arts
College, Yanam – 533 464, Pondicherry, India

Received on : 20-09-2016

Revised and Accepted on : 29-11-2016

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.1.p136-142>

INTRODUCTION

Infectious diseases represent a critical problem to health and they are one of the main causes of morbidity and mortality worldwide¹. Bacterial infection causes high rate of mortality in human population and aquaculture organisms. Preventing disease outbreaks or treating the disease with drugs or chemicals tackles these problems². During the past several years, there has been an increasing incidence of fungal infections due to a growth in immunocompromised population such as organ transplant recipients, cancer and HIV/AIDS patients. This fact coupled with the resistance to antibiotics and with the toxicity during prolonged treatment with several antifungal drugs³ has been the reason for an extended search for newer drugs to treat opportunistic fungal infections⁴. Nowadays the use of antibiotics increased significantly due to heavy infections and the pathogenic bacteria becoming resistant to drugs is common due to indiscriminate use of antibiotics. Decreased efficiency and resistance of pathogen to antibiotics has necessitated the development of new alteration⁵. Many antifungal drugs, including imidazoles, butenafine and terbinafine, have been used clinically for the topical treatment of dermatophytosis⁶. Triazoles, griseofulvin and terbinafine are used as oral antifungal drugs for systemic therapy of severe dermatophytosis⁷, but the prolonged duration of treatment, drug toxicity and interactions, fungal resistance and high costs are encountered difficulties⁸. These factors render the development of new more efficient and safe antifungal drugs. Secondary metabolites produced by plants constitute a major source of bioactive substances. The scientific interest in these metabolites has increased today with the search of new therapeutic agents from plant source, due to the increasing development of the resistance pattern of microorganisms to most currently used antimicrobial drugs. According to World Health Report of infectious diseases, overcoming antibiotic resistance is one of the major issues of the WHO for the present millennium. Hence the last decade witnessed an increase in the investigation of plants as a source of human disease management⁹. Algae appear to be an interesting source for ethno medicinal and phytochemical studies. The power of algal resources has been sought for thousands of years for their ability to prevent disease and prolong life. Algae contain minerals, an abundance of vitamins, variety of trace elements and have shown high potential in controlling antimicrobial, antitumor, anticoagulant and cytotoxic activity¹⁰. Seaweed extracts were also reported to exhibit antimicrobial activity^{11,12}. Active compounds from seaweeds were found to be effective against human bacterial pathogens, fish bacterial pathogens¹³, leaf spot disease of plants and marine pathogenic microorganisms¹⁴. Many chemically unique compounds of marine algae with biological activities have been isolated and a number of them are under investigation and/or are being developed as new pharmaceuticals such as brominated phenols, sterols, terpenoids, polysaccharides, peptides, proteins, acrylic acid, terpenes, chlorophyllides, phenols and heterocyclic carbons etc.¹⁵. Most identified active antimicrobial compounds are water insoluble and thus organic solvent extracts have been found more potent¹⁶. Hence, the

present study was made to evaluate the antibacterial activity of methanol extracts of *U. lactuca*, *C. racemosa*, *C. sertularioides*, *P. gymnospora*, *D. dichotoma*, and *G. multipartita* against *S. aureus*, *E. coli*, *P. aeruginosa*, *K. pneumoniae*, *P. mirabilis* and *S. enterica* against bacterial strains to develop new antibacterial agent with no side effect.

MATERIALS AND METHODS

Algal sample collection

Ulva lactuca, *Caulerpa racemosa*, *Caulerpa sertularioides* (Chlorophyceae) *Padina gymnospora*, *Dictyota dichotoma* (Phaeophyceae) and *Gracilaria verrucosa* (Rhodophyceae) were collected by hand picking from the submerged marine rocks at Thikkodi village (Lat. 8°30'N; Long. 78°8'E), the Calicut West Coast, Kerala, India. Seaweeds collections were made during the month of August, 2012. The algae were identified by Dr. R. Selvaraj, Former Professor, Department of Botany, Annamalai University and the museum specimens are deposited in the Department of Botany, Annamalai University, Annamalainagar Tamilnadu, India.

Preparation of extracts

The algae were handpicked during low tide and washed thoroughly with sea water to remove all unwanted impurities, epiphytes, animal castings and adhering sand particles etc., morphologically distinct thallus of alga were placed separately in new polythene bags and were kept in a ice box containing slush ice and transported to the laboratory. Then, the samples were blot dried using sterile tissue paper. Then the seaweeds were shade dried under room temperature and kept in a hot air oven for 50 °C for half an hour. Oven dried that the material were ground by using electric blender. The powdered materials were stored in air tight container. Five hundred gram of seaweed materials was packed inside a Soxhlet apparatus and successive extraction was carried out using methanol for 72 hours. The solvent was evaporated under vacuum in a rotary evaporator (Heidolph, Germany) and the dried extracts were stored at 4 °C until further assay.

Microorganisms

The gram positive bacterial strain viz., *Staphylococcus aureus* (MTCC 2063) and gram negative bacterial strains such as *Pseudomonas aeruginosa* (MTCC 741), *Klebsiella pneumoniae* (MTCC 109), *Escherichia coli* (MTCC 443), *Proteus mirabilis* (MTCC 425) and *Salmonella enterica* (MTCC 3231) were used in this study. These standard bacterial strains were obtained from Microbial Type Culture Collection (MTCC) Chandigarh, India. The stock cultures were maintained on Nutrient Agar at 4 °C. *In vitro* antibacterial activity was determined by using Mueller Hinton Agar (MHA) and Mueller Hinton Broth (MHB) and they were obtained from Himedia Ltd., Mumbai.

Disc diffusion method

The disc diffusion method of Bauer *et al.*¹⁷ was followed for antibacterial susceptibility test. Petri plates were prepared by pouring 20 ml of MHA. Then the plates were allowed to solidify and used in susceptibility test. The standard inoculum using bacterial suspensions

containing 10^8 colony forming units (CFU) per ml, were swabbed on the top of the solidified respective media and allowed to dry for 10 minutes. The methanol extract was dissolved in 10 per cent Dimethyl sulfoxide (DMSO) and under aseptic conditions. The sterile discs were impregnated with 20 μ l of different concentrations. The discs with extracts were placed on the surface of the medium with sterile forceps and gently pressed to ensure contact with inoculated agar surface. Methicillin (5 μ g/disc) for *S. aureus* and Ciprofloxacin (10 μ g/disc) for all the bacteria were used as positive control and 10 per cent DMSO was used as blind control in all the assays. Finally, the inoculated plates were incubated at 37 °C for 24 h for all bacterial strains. The zone of inhibitions were observed and measured in millimeters. The assay in this experiment was repeated three times.

Determination of the Minimum Inhibitory Concentration (MIC) for bacteria

To determine the MIC for the algae crude extracts, a modified reazurin microtitre plate assay was carried out according to the method of Sarker *et al.*¹⁸. 50 μ l of Sterile MHB were transferred in to each well of a sterile 96-well micro titer plate. The algae extract was dissolved in 10 per cent DMSO to obtain 1000 μ g/ml stock solution. 50 μ l of crude extract stock solution was added into the first well. After fine mixing of the crude extracts and broth, 50 μ l of the solution was transferred to the second well and in this way, the serial dilution procedure was continued to a twofold dilution to obtain concentrations like 1000 to 15.625 μ g/ml of the extract in each well. To each well, 10 μ l of resazurin indicator solution was added. (The resazurin solution was prepared by dissolving a 270 mg in 40 mL of sterile distilled water. A vortex mixer was used to ensure that it

was a well-dissolved and homogenous solution). Finally, 10 μ l of bacterial suspension was added to each well to achieve a concentration of approximately 5×10^5 CFU/mL. Each plate had a set of controls: a column with all solutions with the exception of the crude extracts; a column with all solutions with the exception of the bacterial solution adding 10 μ l of MHB instead and a column with 10 % DMSO solution as a negative control. The plates were incubated at 37 °C for 24 h for all the bacterial strains. The color change was then assessed visually. The growth was indicated by color changes from purple to pink (or colorless). In this study, the MIC was the lowest concentration of algae extracts that exhibited the growth of the organisms the values by visual reading.

Determination of the Minimum bactericidal concentration (MBC)

MBC was determined by plating loop full samples from each MIC assay well with growth inhibition in to freshly prepared MHA for bacteria strains. The plates were incubated at 37 °C for 24 h for all bacterial strains. The MBC was recorded as the lowest concentration of the extracts that did not permit any visible bacterial growth after the period of incubation.

Statistical Analysis

The experimental data are expressed as the mean \pm SD. All statistical analyses were performed using SPSS version 16.0 statistical software (SPSS Inc., Chicago, IL, USA). Comparison of means for *in vitro* antibacterial assessment was carried out using one-way analysis of variance (ANOVA) and Duncan test. *P* value < 0.05 were considered statistically significant.

RESULTS

Table 1
Antibacterial activity of methanol extracts of *Caulerpa racemosa* and *Caulerpa sertularioides*

S. No.	Name of the microorganisms	Mean zone of inhibition ^a (mm) ^b						Control (Me/Cip)	<i>Caulerpa racemosa</i>		<i>Caulerpa sertularioides</i>	
		Concentration of the disc (μ g/disc)			Concentration of the disc (μ g/disc)				MIC μ g/ml	MBC μ g/ml	MIC μ g/ml	MBC μ g/ml
		<i>Caulerpa racemosa</i>			<i>Caulerpa sertularioides</i>							
		1000	500	250	1000	500	250					
1	<i>Staphylococcus aureus</i>	16.5 \pm 1.0	12.8 \pm 1.3	9.3 \pm 1.3	13.3 \pm 1.3	10.5 \pm 1.0	7.3 \pm 0.8	7.5 \pm 2.4	250	500	500	1000
2	<i>Klebsiella pneumoniae</i>	13.0 \pm 1.2	9.8 \pm 1.0	7.0 \pm 0.8	11.3 \pm 1.3	9.0 \pm 0.8	6.5 \pm 0.6	24.3 \pm 2.1	250	500	500	1000
3	<i>Pseudomonas aeruginosa</i>	12.5 \pm 1.7	9.3 \pm 1.5	6.5 \pm 1.3	11.5 \pm 1.3	9.3 \pm 1.3	6.5 \pm 1.0	30.5 \pm 2.4	500	1000	500	1000
4	<i>Escherichia coli</i>	10.0 \pm 1.4	8.5 \pm 1.3	6.5 \pm 1.0	12.5 \pm 1.3	9.0 \pm 1.2	7.3 \pm 1.0	29.0 \pm 1.8	500	1000	500	1000
5	<i>Proteus mirabilis</i>	12.0 \pm 1.0	9.0 \pm 0.8	7.0 \pm 0.8	10.3 \pm 1.0	8.5 \pm 0.6	6.3 \pm 0.5	27.0 \pm 1.8	500	1000	500	1000
6.	<i>Salmonella enterica</i>	13.8 \pm 1.0	9.0 \pm 0.8	7.0 \pm 0.8	11.3 \pm 1.0	8.5 \pm 0.6	6.3 \pm 0.5	28.5 \pm 1.7	500	1000	500	1000

a- Diameter of zone of inhibition (mm) including disc diameter of 6 mm; b- Mean of four assays; \pm standard deviation; Me – Methicillin for *Staphylococcus aureus* (5 μ g/disc); Cip – Ciprofloxacin for all the bacterial except *S. aureus* strains (10 μ g/disc)

Table 2
Antibacterial activity of methanol extracts of *Ulva lactuca* and *Padina gymnospora*

S. No.	Name of the microorganisms	Mean zone of inhibition ^a (mm) ^b						Control (Me/Cip)	<i>Ulva lactuca</i>		<i>Padina gymnospora</i>	
		Concentration of the disc (µg/disc)			Concentration of the disc (µg/disc)				MIC µg/ml	MBC µg/ml	MIC µg/ml	MBC µg/ml
		<i>Ulva lactuca</i>			<i>Padina gymnospora</i>							
		1000	500	250	1000	500	250					
1	<i>Staphylococcus aureus</i>	13.7 ± 1.6	10.6 ± 1.4	7.1 ± 1.2	19.5 ± 1.3	14.0 ± 1.0	9.3 ± 1.3	9.5 ± 1.5	500	1000	125	250
2	<i>Klebsiella pneumoniae</i>	10.0 ± 1.2	8.5 ± 1.0	6.0 ± 0.3	14.1 ± 1.2	11.0 ± 0.6	8.5 ± 0.5	25.0 ± 1.8	250	500	500	1000
3	<i>Pseudomonas aeruginosa</i>	11.5 ± 1.5	9.3 ± 1.2	6.5 ± 1.8	14.1 ± 1.0	11.6 ± 1.5	9.3 ± 1.0	28.5 ± 1.4	500	1000	250	500
4	<i>Escherichia coli</i>	10.0 ± 1.2	8.1 ± 1.1	6.3 ± 1.0	14.2 ± 1.5	11.0 ± 1.0	9.3 ± 1.5	29.0 ± 1.8	500	1000	500	1000
5	<i>Proteus mirabilis</i>	11.8 ± 1.2	8.0 ± 0.3	7.0 ± 0.8	13.3 ± 1.4	11.5 ± 1.8	7.3 ± 1.7	27.0 ± 1.5	500	1000	250	500
6.	<i>Salmonella enterica</i>	12.4 ± 1.5	9.0 ± 1.8	7.0 ± 1.2	14.3 ± 1.3	12.5 ± 0.6	9.5 ± 1.6	25.5 ± 1.6	500	1000	250	500

a- Diameter of zone of inhibition (mm) including disc diameter of 6 mm; b- Mean of four assays; ± standard deviation; Me – Methicillin for *Staphylococcus aureus* (5 µg/disc); Cip – Ciprofloxacin for all the bacterial except *S. aureus* strains (10 µg/disc)

Table 3
Antibacterial activity of methanol extracts of *Dictyota dichotoma* and *Gracilaria multipartita*

S. No.	Name of the microorganisms	Mean zone of inhibition ^a (mm) ^b						Control (Me/Cip)	<i>Dictyota dichotoma</i>		<i>Gracilaria multipartita</i>	
		Concentration of the disc (µg/disc)			Concentration of the disc (µg/disc)				MIC µg/ml	MBC µg/ml	MIC µg/ml	MBC µg/ml
		<i>Dictyota dichotoma</i>			<i>Gracilaria multipartita</i>							
		1000	500	250	1000	500	250					
1	<i>Staphylococcus aureus</i>	18.3 ± 1.8	13.4 ± 1.5	7.3 ± 1.3	15.1 ± 1.6	12.0 ± 1.3	7.3 ± 1.0	8.6 ± 1.5	125	250	250	500
2	<i>Klebsiella pneumoniae</i>	13.0 ± 1.4	9.5 ± 1.0	7.1 ± 1.6	12.3 ± 1.5	10.0 ± 1.2	7.5 ± 0.6	27.3 ± 1.4	250	500	500	1000
3	<i>Pseudomonas aeruginosa</i>	13.3 ± 1.3	10.3 ± 1.2	8.5 ± 1.1	12.1 ± 1.3	9.3 ± 1.8	7.3 ± 1.5	29.3 ± 1.6	500	1000	250	500
4	<i>Escherichia coli</i>	13.0 ± 1.5	10.3 ± 1.2	8.7 ± 1.6	12.3 ± 1.2	9.0 ± 1.5	7.3 ± 1.6	28.0 ± 1.5	500	1000	500	1000
5	<i>Proteus mirabilis</i>	14.3 ± 1.4	11.0 ± 0.5	9.3 ± 0.2	12.0 ± 1.0	9.5 ± 0.9	7.3 ± 0.2	29.0 ± 1.8	250	500	250	500
6.	<i>Salmonella enterica</i>	16.5 ± 1.4	12.0 ± 0.7	10.0 ± 1.3	12.1 ± 1.0	9.5 ± 0.5	6.9 ± 1.0	27.5 ± 1.0	250	500	500	1000

a- Diameter of zone of inhibition (mm) including disc diameter of 6 mm; b- Mean of four assays; ± standard deviation; Me – Methicillin for *Staphylococcus aureus* (5 µg/disc); Cip – Ciprofloxacin for all the bacterial except *S. aureus* strains (10 µg/disc)

In the present study, methanol extracts of *U. lactuca*, *C. racemosa*, *C. sertularioides*, *P. gymnospora*, *D. dichotoma* and *G. multipartita* were studied against *S. aureus*, *E. coli*, *P. aeruginosa*, *K. pneumoniae*, *P. mirabilis* and *S. enterica*. The highest mean zone of inhibition was observed by methanol extract of *Padina gymnospora* against *Staphylococcus aureus* (the mean zones of inhibition, 19.5 mm). All the extracts of marine macro algae tested possessed significant antibacterial activity against all the bacterial strains tested when compared to the available antibiotics tested. The mean values are presented in Tables 1, 2 & 3. When the different extracts were assayed against the test bacteria by agar disc diffusion assays, the mean zones of inhibition obtained were between 6.3 and 19.5 mm. Methicillin (10 µg/disc), antibacterial positive control produced mean zones of inhibition ranged from 7.5 to 9.5 mm. The Ciprofloxacin (10 µg/disc), antibacterial positive control produced mean zones of inhibition ranged from 24.3 to 30.5 mm. The blind control (10% DMSO) did not produce any zone of inhibition for all the bacterial strains tested. The MIC values of the different extracts of algal extracts ranged between 125 and

500µg/ml, while the MBC values were between 250 and 1000 µg/ml.

DISCUSSION

Marine macro algae are eukaryotic organisms that live in salty water in the ocean and is recognized as a potential source of bioactive natural products¹⁹. They contain compounds ranging from sterols, terpenoids to brominated phenol, which shows bioactivity against microorganisms²⁰. Presently seaweeds constitute commercially important marine renewable resources which are providing valuable ideas for the development of new drugs against cancer, microbial infections and inflammations²¹. In the present investigation, methanol extracts of *U. lactuca*, *C. racemosa*, *C. sertularioides*, *P. gymnospora*, *D. dichotoma*, and *G. multipartita* showed the highest antibacterial activity against *S. aureus*, *E. coli*, *P. aeruginosa*, *K. pneumoniae*, *P. mirabilis* and *S. enterica*. The highest mean zones of inhibition (19.5 mm) was observed in methanol extract of *P. gymnospora* against *S. aureus*. The MIC values of the methanol extracts of *P. gymnospora* ranged between 125 and 500µg/ml, while the MBC values were between

250 and 1000 µg/ml. Salem *et al.*²² reported that higher antibacterial activity was recorded for the ethyl acetate extracts of *C. racemosa*, *Sargassum dentifolium*, *Padina gymnospora*; methanolic extracts of *Sargassum hystrix*, *C. racemosa*, *C. fragile*, *S. dentifolium* and *Cystoseria myrica* against *E. coli*, *S. aureus*, *E. faecalis*, *Salmonella sp.*, *B. cereus* and *P. aeruginosa*. These results were in close agreement with those obtained by Patra *et al.*²³. It was revealed that the chloroform and ethyl acetate extracts of *Enteromorpha compressa*, *Chaetomorpha linum* and *Polysiphonia subtilissima* were active against *Bacillus subtilis*, *B. brevis*, *E. coli*, *S. flexneri*, and *Vibrio cholerae*. Rangaiaha *et al.*²⁴ reported that the *Sargassum ilicifolium*, *Padina tetrastromatica* showed the highest inhibition was noticed with ethanol extracts and lowest with chloroform crude extracts while in case of *Gracilaria corticata*, maximum inhibition was noticed with methanol and minimum with chloroform extracts. Antifungal activity of all the crude extractions of *G. corticata* showed highest activity against *Rhizopus stolonifer*. Mansuya *et al.*²⁵ reported the aqueous and methanolic extract of *U. lactuca*, *U. reticulata*, *Cladophora glomerata*, *G. corticata*, *Kappaphycus alvarezii* and *Sargassum wightii* against *E. coli*, *P. aeruginosa*, *S. typhi*, *Staphylococcus epidermidis* and *S. pyogenes*. The antibacterial activity from methanol, ethanol, dichloromethane and hexane extracts of *Gracilaria fisheri* and *Ulva intestinalis* was tested against *S. aureus*, *Listeria monocytogenes*. Methicillin-resistant *S. aureus*, *Enterobacter faecalis*, *V. alginolyticus*, *V. parahaemolyticus*, *V. harveyi*, *E. coli*, *P. aeruginosa*, *K. pneumoniae*, *Salmonella typhi* and *P. mirabilis*²⁶. Chandrasekaran *et al.*²⁷ showed the antibacterial activity of *U. fasciata* against multi-drug resistant bacterial strains of *B. subtilis*, *S. pyogenes*, *E. coli*, *K. pneumoniae*, *P. aeruginosa*, *S. typhimurium*, *V. cholerae*, *S. flexneri*, *P. mirabilis* and *P. vulgaris*. In the present study, methanol extracts of *U. lactuca*, *C. racemosa*, *C. sertularioides*, *P. gymnospora*, *D. dichotoma*, and *G. multipartita* have possessed antibacterial activity against all the clinical and standard bacterial strains tested. Hediati *et al.*²⁸ reported that different solvents have been reported to have the capacity to extract different phytoconstituents depending on their solubility or polarity in the solvent. In this present study also supported that optimizes their antibacterial activity by methanol extract the active compound from seaweeds. Seaweed extracts in different solvents exhibited different antimicrobial activities²⁴. The high and low activity of organic extracts against microorganisms could be related to the presence of bioactive metabolites, which can be soluble in solvents²⁹. In the present study, methanol extracts of marine macro algae possessed antibacterial activity against all the bacterial strains tested. Chandrasekaran *et al.*¹² methanol extracts of *Sargassum wightii* showed highest antibacterial activity against multi-drug resistant strains of *B. subtilis*, *S. pyogenes*, *E. coli*, *K. pneumoniae*, *P. aeruginosa*, *S. typhimurium*, *V. cholerae*, *S. flexneri*, *S. dysenteriae*, *P. mirabilis* and *P. vulgaris*. The methanol crude extracts of *Sargassum longifolium* and *G. corticata* showed highest antibacterial activity against *Aeromonas hydrophila*, *P. aeruginosa*,

V. cholerae, *V. harveyi* and *V. parahaemolyticus*³⁰. The methanol extracts of *G. verrucosa*, *G. ferugunoi*, *G. verrucosa* var *Hypnea musciformis*, *Enatiocladia prolifera*, and *Gelidium* species showed highest antibacterial activity against *B. subtilis*, *E. coli*, *P. aeruginosa*, *S. typhi*, *Streptococcus aureus* and *Candida albicans*³¹. Radhika *et al.*³² reported the antibacterial activity of seaweeds *Ulva fasciata*, *S. wightii* and *G. corticata* against *Bacillus cereus*, *Vibrio cholerae* classical, *V. cholerae* 0139, *E. coli*, *Pseudomonas aeruginosa*, *Aeromonas hydrophila*, *Salmonella typhi* and *Shigella flexneri*. Phytochemicals are compounds from food and medicine to protect and maintain human health. These have antioxidant or hormone-like effect which helps to fight against diseases like cancer, heart disease, diabetes, high blood pressure and preventing the formation of carcinogens on their target tissues. It is reported earlier that seaweeds are also rich in polysaccharides such as alginates, fucans, and laminarans which possess medicinal values³³. Krishnaveni and Johnson³⁴ reported that phytochemical analysis of various solvents extracts revealed that the presence of alkaloids, glycosides, saponins, steroids, phenol and tannins in *G. corticata*. Glycosides serve as defence mechanisms against predation by many microorganisms, insects and herbivores³⁵. Phytochemicals such as saponins, terpenoids, flavonoids, tannins, steroids and alkaloids are reported to have antimicrobial effects³⁶. Tannins play a major role as antihaemorrhagic agent and showed to have immense significance as antihypercholesterol, hypotensive and cardiac depressant properties³⁷. Glycosides, flavonoids, tannins and alkaloids have hypoglycemic activities³⁸. Steroids, saponins and triterpenoids showed the analgesic properties³⁹. Phenolic compounds may affect growth and metabolism of bacteria. They could have an activating or inhibiting effect on microbial growth according to their constitution and concentration⁴⁰. In the present study, Gram-positive bacteria were found to be more susceptible than the Gram-negative bacteria. The resistance of Gram negative bacteria towards antibacterial substances is related to the hydrophilic surface of their outer membrane which is rich in lipopolysaccharide molecules, presenting a barrier to the penetration of numerous antibiotic molecules. The membrane is also associated with the enzymes in the periplasmic space which are capable of breaking down the molecules introduced from outside⁴¹. However, the Gram positive bacteria do not possess such outer membrane and cell wall structures⁴¹. In the present study, almost all crude extracts tested have shown strong antibacterial potential against pathogenic bacteria.

CONCLUSION

Finally, it can be conclude that study proved the efficiency of antibacterial properties of seaweeds and advocates the potentiality of the plant as a source of alternative medicine. So, use of natural products, especially *P. gymnospora* and *D. dichotoma* may be considered as a new source of natural antibacterial agents.

ACKNOWLEDGMENT

We thank the Professor and Head Department of Botany, Annamalai University, for having provided laboratory facilities.

CONFLICT OF INTEREST

Conflict of interest declared none.

REFERENCES

- Mandal FB. The Changing Ecology of Infectious Diseases in Human. World Environ. 2011 Jan; 1(1): 14-9.
- Kandhasamy M, Arunachalam KD. Evaluation of *in vitro* antibacterial property of seaweeds of southeast coast of India. African J Biotech. 2008 Jun 17; 7(12): 1958–61.
- Giordani R, Trebaux J, Masi M, Regli P. Enhanced antifungal activity of ketoconazole by *Euphorbia characias* latex against *Candida albicans*. J Ethnopharmacol. 2001 Nov 78 (1); 1–5.
- Fostel J, Lartey P. Emerging novel antifungal agents. Drug Discovery Today. 2000 Jan; 5(1): 25–32.
- Claudio D and Miranda. Raul Zemelman Bacterial resistance to oxytetracycline in Chilean salmon farming. Aquacul. 2002 Mar 11; 212: 31–47.
- Watanabe S. Present state and future direction of topical antifungals Japanese. J Med Mycol. 1999 Jan 28; 40: 151–5.
- Leshner JL. Oral therapy of common superficial fungal infections of the skin. J Am Acad Dermatol. 1999 Jun 40; 40: 31–4.
- Bennett ML, Fleischer Jr AB, Loveless JW, Fledman SR. Oral griseofulvin remains the treatment of choice for *tinea capitis* in children. Pediatric Dermatol. 2000 Aug; 17 (4): 304–9.
- Prashanth D, Asha MK, Amit A. Antibacterial activity of *Punica granatum*. Fitoterapia. 2001 Feb; 72: 171–3.
- Kolanjinathan K, Ganesh P, Saranraj P. Pharmacological Importance of Seaweeds: A Review World J Fish Mar Sci. 2014 Jun 6; 6 (1): 01-15.
- Chandrasekaran M, Venkatesalu V, Adaikala Raj G. Anti-MRSA activity of Brown and Red algae from Gulf of Mannar Coast, South India. Int J Life Sci Technol. 2014a Dec 4; 7(14): 22-31.
- Chandrasekaran M, Venkatesalu V, Adaikala Raj G, Krishnamoorthy S. Antibacterial properties of various extracts of *Sargassum wightii* against Multidrug Resistant Bacterial Strains. Phykos. 2014b Jan 2; 44(2): 17 – 28.
- Kolanjinathan K, Ganesh P, Govindarajan M. Antibacterial activity of ethanol extracts of seaweeds against fish bacterial pathogens. European Rev Med Pharmacol Sci. 2009 Jun; 13 (3): 173-7.
- Engel S, Puglisi MP, Jensen PR, Fenical W. Antimicrobial activities of extracts from tropical Atlantic marine plants against marine pathogens and saprophytes. Marine Biol. 2006 Aug; 149 (5): 991-1002.
- Priyadharshini S, Bragadeeswaran S, Prabhu K, Ran SS. Antimicrobial and hemolytic activity of seaweed extracts *Ulva fasciata* (Delile 1813) from Mandapam, Southeast coast of India. Asian Pac J Trop Biomed. 2011 Sep 10; 1 (1): 38–9.
- Villanueva RD, Sousa AMM, Gonçalves MP, Nilsson M, Hilliou L. Production and properties of agar from the invasive marine alga, *Gracilaria vermiculophylla* (Gracilariales, Rhodophyta). J Appl Phycol. 2010 Apr 9; 22 (2): 211–20.
- Bauer AW, Kirby WMM, Scherris J, Turck C. Antibiotic susceptibility testing by a standardized single disk method. Am J Clin Pathol. 1966 Apr; 45 (4): 493-6.
- Sarker SD, Nahar L, Kumarasamy Y. Microtitre plate-based antibacterial assay incorporating resazurin as an indicator of cell growth, and its application in the *in vitro* antibacterial screening of phytochemicals. Methods. 2007 Aug; 42(4): 321-4.
- Michael TM, John MM, Jack P. Brock Microbiology of Microorganisms. 11th Edition, New Jersey; 2005.
- Perry NB, Blunt JW, Munro, MHA. Cytotoxic and antifungal 1,4-naphthoquinone and related compounds from a New Zealand brown algae. *Landsburgia quercifolia*. J Nat Prod. 1991 Aug; 54 (4): 978-85.
- Elena M, Francisco Y, Erickson KL. "Mailiohydrin, a Cytotoxic *Chamigrene Dibromohydrin* from a Phillipine *Laurencia* Species," J Nat Prod. 2003 Jan; 64 (6): 790-91.
- Patra JK, Rath SK, Jena K, Rathod VK, Thatoi H. Evaluation of Antioxidant and Antimicrobial Activity of Seaweed (*Sargassum* sp.) Extract: A Study on Inhibition of Glutathione-S-Transferase Activity. Turk J Biol. 2008 Jun; 32: 119-25.
- Salem WM, Galal H, Nasr El-deen F. Screening for antibacterial activities in some marine algae from the red sea (Hurgada, Egypt). Afr J Microbiol Res. 2011 Aug 4; 5(15): 2160-67.
- Rangaiaha GS, Lakshmi PA, Sruthikeerthia K. The antimicrobial activity of the crude extracts of Chlorophycean seaweeds *Ulva*, *Caulerpa* and *Spongomorpha* spp. against clinical and phytopathogens. Drug Invent Today. 2010 Apr 4; 2 (6): 311-4.
- Mansuya P, Aruna P, Sridhar S, Suresh Kumar J, Babu S. Antibacterial activity and qualitative phytochemical analysis of selected seaweeds from Gulf of Mannar Region. J Exper Sci. 2010 Jun; 1(8): 23-6.
- Srikong W, Mittraparp-arthorn P, Rattanaporn O, Bovornreungroj N, Bovornreungroj P. Antimicrobial activity of seaweed extracts from

- Pattani, Southeast coast of Thailand. Food Appl Biosci J. 2015 Jan; 3 (1): 39–49.
27. Chandrasekaran M, Venkatesalu V, Adaikala Raj G, Krishnamoorthy S. Antibacterial activity of *Ulva fasciata* against Multidrug Resistant Bacterial Strains. Int Lett Nat Sci. 2014c Jun; 14: 40 – 51.
 28. Hediath MH, Salama, Najat M. Antimicrobial activity and phytochemical analyses of *Polygonum aviculare* L. (Polygonaceae), naturally growing in Egypt. Saudi J Biol Sci. 2010 Jan 17; 17: 57-63.
 29. Elakkia SA, Venkatesalu V. Antimicrobial activity of different solvent extracts of some *Cassia* species. Int J Pharm Bio Sci. 2013 July; 4(3): 728 – 36.
 30. Sangeetha S, Dhayanithi NB, Sivakumar N. Antibacterial activity of *Sargassum longifolium* and *Gracilaria corticata* From Gulf of Mannar against selected Common Shrimp Pathogens. Int J Pharm Bio Sci. 2014 Apr; 5(2): 76 – 82.
 31. Adaikala Raj G, Patric Raja D, Johnson M, Janakiraman N, Babu A. Antibacterial potential of selected red seaweeds from Manapad coastal areas, Thoothukudi, Tamil Nadu, India. Asian Pacific J Trop Biomed. 2012 July; 1077-80.
 32. Radhika D, Veerabahu C, Vijayalakshmi M, Priya R. Antibacterial Effect of some species of seaweed from different station of the gulf of mannar coast. Int J Biol Pharmace Res. 2013 Jan; 4(11): 783-7.
 33. Smit AJ. "Medicinal and pharmaceutical uses of seaweed natural products: a review," J Appl Phycol. Apr 21; 16: 245– 62. (2004).
 34. Krishnaveni E, Johnson M. Preliminary Phytochemical, UV-VIS, HPLC and Anti-bacterial Studies on *Gracilaria corticata* J. Ag. Asian Pac J Trop Biomed. 2012 Jan; 568-74.
 35. Dhar ML, Dhar MM, Dhawan BN, Ray C. Screening of Indian plants for biological activity. Part-I, Indian J Biol. 1979 July 19; 6: 232-4.
 36. Vinoth B. Manivasagaperumal R. Antimicrobial activity of different extracts of *Azima tetracantha* root. Int J Pharm Bio Sci. 2015 Apr; 6(2): 613 – 20.
 37. Cherian S, Augusti KT Insulin sparing action of leucopelargonidin derivative isolated from *Ficus bengalensis* L, Indian J Exper Biol. 1995 Sep 16; 33: 608-11.
 38. Miliauskasa G, Venskutonisa PR. Beekb van TA. Screening of radical scavenging activity of some medicinal and aromatic plant extracts. Food Chem. 2004 Dec; 85: 231–7.
 39. Reguant C, Bordons Arola A, Roze LN. Influence of phenolic compounds on the physiology of *Oenococcus oeni*. J Appl Microbiol. 2000 Jun; 88(6): 1065-71.
 40. Shan B, Cai Y-Z, Brooks JD, Corke H. The *in vitro* antibacterial activity of dietary spice and medicinal herb extract. Int J Food Microbiol. 2007 Jun 10; 117: 112-9.
 41. Tajkarimi MM, Ibrahima SA, Cliver DO. Antimicrobial herb and spice compounds in food. Food Control. 2010 Sep; 21(9): 1199–218.