



EFFICIENT PROTOCOL FOR REGENERATION OF TRANSFORMED *ARTEMISIA ANNUA* PLANTS FROM HAIRY ROOTS

SUGANDHA TIWARI*

Department of Botany, DG College, CSJM University, Kanpur – 208001, UP, India.

ABSTRACT

Elite genotypes from high yielding variety of *Artemisia annua* L. (Jeevanraksha, CIMAP Lucknow) were regenerated from nodal explants and maintained in MS culture medium with BAP and NAA. The leaves of tissue culture grown plants were inoculated with *Agrobacterium rhizogenes* strain LBA9402 containing the selective marker gene NPT II. Hairy roots emerged from the leaves, 8-10 days after inoculation and were maintained in MSO liquid/ solid medium with kanamycin as the selective agent. Hairy root line was selected on the basis of high artemisinin content. Regeneration from hairy roots was obtained in half strength MS solid medium containing a combination of cytokinins and auxins (BAP, Kinetin, NAA and IBA). Transgenic plants showed typical transgenic morphology and profuse growth of plagiotropic roots. Transformed hairy root cultures and transgenic plants can be used as an experimental system for studies related to induction of secondary metabolite synthesis.

KEYWORDS: Artemisinin, sesquiterpene lactone, antimalarial, *Agrobacterium rhizogenes*, Benzylaminopurine (BAP), Naphthaleneacetic acid (NAA), Indolebutyric acid (IBA)



*SUGANDHA TIWARI

Department of Botany, DG College, CSJM University, Kanpur – 208001, UP, India.

Received on : 26-09-2016

Revised and Accepted on 20-12-2016

DOI: <http://dx.doi.org/10.22376/ijpbs.b396-401>

INTRODUCTION

Artemisinin also known as qinghaosu is a sesquiterpene lactone, naturally formed in the shoots of *Artemisia annua* L. Artemisinin has proved to be an effective and safe antimalarial agent, exhibiting excellent activities against *Plasmodium falciparum* parasites that are resistant to commonly used antimalarial drugs.¹ Artemisinin and its derivatives are also found to be effective against Schistosomiasis caused by parasite *Schistosoma japonica*, *S. mansoni* and *S. haematobia* causing 1-1.5 million disabilities each year.² Oral administration of artemisinin has also been found to prevent and delay the development of 7, 12-dimethylbenz (a) anthracene (DMBA) induced breast cancer in rat.³ Artemisinin reacts with iron and forms cytotoxic free radical. It is selectively more toxic to cancer than normal cells because cancer cells contain more intracellular free iron. Covalent tagging of artemisinin to transferrin is reported to increase its selectivity and toxicity towards cancer cells in vitro. Artemisinin- transferrin conjugate significantly retarded the growth rate of breast tumors in rat and could be developed into a potent therapeutic agent of cancer in humans.⁴ Artemisinin is also reported to be effective against human leukemia, colon cancer and lung carcinomas.^{5,6} According to WHO, artemisinin based combination therapy (ACT) is the only reliable treatment option for multidrug resistant *Plasmodium falciparum* malaria.⁷ Low yield of artemisinin in *A. annua* (0.01-0.8%, in some genotypes 1.5%) is a serious limitation to meet the annual targets of the drug for ACT treatments. Research studies are conducted in many laboratories to enhance artemisinin content in cell or tissue cultures of *A. annua* L. Biotechnological approaches are also being explored to increase artemisinin content. Transformed root and shoot cultures of *A. annua* were established by infection with *Agrobacterium rhizogenes* and *A. tumifaciens*. Hairy roots were induced from *A. annua* L. by transformation with *Agrobacterium rhizogenes*.^{8,9} Transformed hairy root cultures of *A. annua* L were produced by co-culture method using leaf segments of

A. annua and *A. rhizogenes* strain NCIB 8196 or MAFF 03-01724.¹⁰ Hairy roots from *A. annua* L. leaf blade pieces and petiole segments were induced by *A. rhizogenes* strain 1602 and a clone was obtained with high content of artemisinin (1.195 mg/g dry weight).¹¹ *Agrobacterium tumifaciens* mediated transformation system was developed for *A. annua* L. plants with high transformation rates (75% regenerants harbouring foreign genes).¹² Studies have been conducted in different laboratories to elucidate the biochemical pathway of artemisinin and its regulation with an aim to improve artemisinin content of *A. annua* L.^{13,14,15} Conversion of farnesyl diphosphate to amorpha 4-11-diene is the first committed step in the biosynthesis of artemisinin, but little is known about the intermediates and enzymes that catalyze the pathway of artemisinin biosynthesis beyond amorpha- diene to artemisinic acid.¹³ Twelve genes related to artemisinin biosynthetic pathway have been cloned from *A. annua* L. Researches in the field of transgenics aim to increase artemisinin content in the plants and to increase the supply of artemisinin and its derivatives for ACT treatments. Increase in the yield of artemisinin would significantly bring down the cost of malaria treatment and would help in reducing the incidence of malaria.

MATERIALS & METHODS

Tissue culture of field grown *Artemisia annua* plants

Seeds of high yielding variety of *A. annua* L. (Jeevanraksha, CIMAP Lucknow)¹⁶ were sown and plants grown to maturity in the field. Field grown plants (fig 1) were screened for artemisinin content at preflowering and flowering stage. Nodal explants from field grown *A. annua* plants, having high artemisinin content (0.5% and above) were selected for tissue culture. *Artemisia annua* plants were surface sterilized with 0.1% (w/v) mercuric chloride for 2 min, rinsed thoroughly with sterile distilled water 7-8 times and cultured on MS basal medium (table 1, fig 2,3,4) with BAP (0.2mg/l) and NAA (0.05mg/l).

Table 1
Murashige and Skoog Medium for tissue culture of *Artemisia annua*

Composition	Concentration
MS A	10 ml/l
MS B	10 ml/l
MS C	5 ml/l
MS D	5ml/l
MS E	5ml/l
MS F	5 ml/l
Sucrose	10 g/l
Myo Inositol	100 mg/l
Casein hydrolysate	0.5 mg/l
RT Vitamins	2 ml/l
Biotin	5 ml/l
Agar agar	8 g/l
BAP	0.2 mg/l
NAA	0.05 mg/l
pH	5.7

Agrobacterium rhizogenes culture and infection of *Artemisia annua* leaves

Agrobacterium rhizogenes strain LBA 9402 containing NPT II gene for kanamycin resistance was maintained in

YMB medium (table 2) with 50µg/ ml kanamycin. For transformation single cell colony of *A. rhizogenes* LBA 9402 was inoculated in YMB liquid medium and kept on rotary shaker. The bacterial culture was used after 24

hours for inoculation in axenically grown *A. annua* leaves. The leaf tissue was wounded with a syringe loaded with bacterial culture and incubated in MSO medium in dark at 5°C for 24 hours. The leaves were then transferred to MS basal medium containing the selective agent kanamycin (50µg/ ml) and an antibiotic sporidex (1000 µg/ ml) to eliminate excess agrobacterial growth. Hairy roots emerged from the leaves 8-10 days

after inoculation (fig 5). These hairy root cultures were maintained in MSO medium with kanamycin (50µg/ ml) as the selective agent and kept at 25 ± 2°C in dark (fig 6). The neoplastic (cancerous) roots produced by *A. rhizogenes* infection are characterized by high growth rate, genetic stability and growth in hormone free media. Root emergence from non transformed explants (control) was not observed.

Table 2
YMB medium for culture of *Agrobacterium rhizogenes*

Composition	Concentration
K ₂ HPO ₄	0.5 g/l
MgSO ₄	2.0 g/l
NaCl	0.1 g/l
Mannitol	10.0 g/l
Yeast extract	0.4 g/l
Kanamycin	50 µg/ ml
Agar agar	15.0 g/l
pH	7.0

Regeneration of transgenic *Artemisia annua* plants from hairy roots

MS media with varying sucrose levels and different combinations and concentrations of auxin and cytokinin were tested for regeneration of transgenic plants. Single combination of cytokinin and auxin were not able to induce regeneration. Regeneration of transgenic *A. annua* plants from hairy roots was obtained in a media containing a combination of cytokinins and auxins (BAP

+ Kinetin + NAA + IBA). The selective agent kanamycin (50µg/ ml) was maintained in all the culture media throughout plant development (table 3, fig 7,8). Transformed cells and hairy root lines containing NPT II gene encode enzyme neomycin phosphotransferase (NPT) and thus inactivate the antibiotic kanamycin present in the medium, but normal plant cells are very sensitive to kanamycin and are killed in a medium containing kanamycin.

Table 3
Media composition (MS Media) for regeneration of transformed *Artemisia annua* plants from hairy roots

Composition	Concentration
MS A	5.0 ml/l
MS B	5.0 ml/l
MS C	2.5 ml/l
MS D	2.5 ml/l
MS E	2.5 ml/l
MS F	2.5 ml/l
Sucrose	20 g/l
Myo inositol	100 mg/l
Casein hydrolysate	0.5 g/l
RT vitamins	2 ml/l
Biotin	5 ml/l
Kanamycin	50µg/ ml
BAP	2 mg/l
Kn	3 mg/l
NAA	0.05 mg/l
IBA	0.05 mg/l
pH	5.7

RESULTS & DISCUSSION

The study reports for the first time the hormone combination and concentration for the production of transgenic *A. annua* plants from hairy roots. Regeneration from hairy roots was obtained in half strength MS solid medium containing a combination of cytokinins and auxins (BAP 2mg/l, Kinetin 3mg/l, NAA 0.05mg/l & IBA 0.05mg/l). Transgenic plants showed typical transgenic morphology and profuse growth of

plageotropic roots (fig 8). Conversion of farnesyl diphosphate to amorpha 4-11- diene is the first committed step in the biosynthesis of artemisinin.⁷ Two genes of artemisinin biosynthetic pathway have now been identified – amorpha 4-11-diene synthase and sesquiterpene cyclase or synthase.⁹ Transformed hairy root cultures and transgenic plants can be used as an experimental system for studies related to induction of secondary metabolite synthesis.



Figure 1
Field grown plant of A. annua




Figure 2
Internodal explant in medium




Figure 3
Growth of tissue culture explant



Figure 4
Tissue culture raised plants




Figure 5
Induction of hairy roots




Figure 6
Hairy root culture



Figure 7
Regeneration of plant from hairy root




Figure 8
Transgenic A. annua plant

CONCLUSION

It would be economically interesting to produce transgenic plants of *Artemisia annua* which ensure a constant high production of artemisinin by over-expressing a key enzyme in the biosynthetic pathway of artemisinin or by inhibiting an enzyme of another pathway, competitive for the precursors. The 'Hairy root' system can be used as a tool for plant breeding and improvement, via the transfer of desirable genes and for studies of gene expression with respect to changes in phenotype and secondary metabolism.

REFERENCES

1. Woerdenbag HJ, Lugt CB, Pras N. *Artemisia annua* L: a source of novel antimalarial drugs. *Pharmaceut. Weekbl. (Sci)*. 1990; 12 : 169-181.
2. Shuhua X, Tanner MN, Goran EK, Utzinger J, Chollet J, Bergquist R, Minggang C, Jiang Z. Recent investigations of artemether, a novel agent for the prevention of *Schistosomiasis japonica, mansoni and haematobia*. *Acta. Trop.* 2002; 82:175-181.
3. Lai H, Singh NP. Oral artemisinin prevents and delays the development of 7, 12- dimethylbenz 9a) anthracene (DMBA) induced breast cancer in the rat. *Cancer Lett.* 2006; 231(1): 43-48.
4. Lai H, Nakase I, Lacoste E, Singh NP, Sasaki T. Artemisinin transferrin conjugate retards growth of breast tumors in the rat. *Anticancer Res.* 2009; 29 : 3807-3810.
5. Efferth T, Dunstan H, Sauerbrey A, Miyachi H, Chitambar GR. The antimalarial artesunate is also effective against cancer. *Int J of Oncology.* 2001;18: 767- 773.
6. Singh NP, Lai H. Selective toxicity of dihydroartemisinin and holotransferrin towards human breast cancer cells. *Life Sci.* 2001; 70: 49-56.
7. WHO. Antimalarial drug combination therapy: Report of a Technical Consultation. WHO, Geneva 1-36. (4-5 April 2001)
8. Weathers PJ, Cheetham RD, Follansbee E, Teoh T. Artemisinin production by transformed roots of *Artemisia annua* L. *Biotechnol. Lett.* 1994; 16: 1281-1286.
9. Qin MB, Li GZ, Yun Y, Ye HC, Li GF. Induction of hairy root from *Artemisia annua* with *Agrobacterium rhizogenes* and its culture in vitro. *Acta Bot. Sin.*1994; 36: 165-170.
10. Jaziri M, Shimomura K, Yoshimatsu K, Fauconnier ML, Marlier M, Homes J. Establishment of normal and transformed root cultures of *Artemisia annua* L. for artemisinin production. *J. of Plant Physiol.* 1995; 145: 175-177.
11. Xie DY, Zou ZR, Ye HC, Li GF, Guo ZC. Selection of hairy root clones of *Artemisia annua* L. for artemisinin production. *Isr. J. Plant Sci.* 2001; 49: 129-134.
12. Vergauwe A, Cammaert R, Vandenberghe D, Genetello C, Van Montagu M, Vanden Eeckhout E. *Agrobacterium tumefaciens* mediated transformation of *Artemisia annua* L. *Plant Cell Rep.* 1996; 15: 929-937.
13. Berteau CM, Freije JR, Van der Woud H, Verstappen FW, Perk L, Marquez V, De Kraker JW, Posthumus MA, Jansen BJM, De Groot AE, Franssen MCR, Bouwmeester HJ. Identification of intermediates and enzymes involved in the early steps of artemisinin biosynthesis in *Artemisia annua*. *Planta Medica.* 2005; 71 (1): 40-47.
14. Weathers PJ, Elkholy S, Wobbe KK. Artemisinin: The biosynthetic pathway and its regulation in *Artemisia annua*, a terpenoid rich species. *In Vitro Cell. Dev. Biol.* 2006; 42 (4): 309-317.
15. Tiwari S. Recent advances in optimization and maximization of artemisinin yield for artemisinin based combination therapy. *Int. J. Pharm Bio Sci.* 2012; 3(4): 686-696.
16. Kumar S , Banerjee S , Dwivedi S, Gupta MM, Verma RK, Jain DC, Khanuja SPS, Mathur AK, Bagchi GD, Zehra M, Mehta VK, Naqvi AA, Paul S, Ram G, Ram M, Saikia D, Sangwan RS, Kumar TRS, Shasany AK, Darokar MP, Singh AK, Singh A. Registration of Jeevanraksha and Suraksha varieties of antimalarial medicinal plant *Artemisia annua*. *J. Med. Arom. Plant Sci.*1999; 21: 47-48.

ACKNOWLEDGEMENT

I gratefully acknowledge the guidance and support of Dr. Suchitra Banerjee, Scientist CIMAP, Lucknow and Prof. Sushil Kumar, Former Director CIMAP, Lucknow.

CONFLICT OF INTEREST

Conflict of interest declared none.