



CORRELATION BETWEEN LEVEL OF ALBUMIN AND GLYCOSYLATED HAEMOGLOBIN IN SOUTH INDIAN PATIENTS WITH TYPE 2 DIABETES MELLITUS

S. SAKTHI DASAN¹, C. SUTHAKARAN^{2*}

¹ Associate Professor, Department of Biochemistry, Melmaruvathur Adhi Parasakthi Institute of Medical Sciences & Research, Melmaruvathur 603319, Kancheepuram Dist, Tamilnadu, India

² Associate Professor Department of Pharmacology Melmaruvathur Adhi Parasakthi Institute of Medical Sciences & Research, Melmaruvathur 603319, Kancheepuram Dist, Tamilnadu, India Affiliated to Tamilnadu Dr MGR Medical University

ABSTRACT

Type 2 diabetes mellitus in Asian Indian people is characterized by a young age of onset and occurrence at low levels of BMI. Available data also suggest that the susceptibility of Asian Indian people to the complications of diabetes mellitus differs from that of white populations. To investigate whether a correlation exists between HbA1c and serum albumin levels in South Indian patients with type 2 diabetes mellitus, a prospective study of electronic case records of type 2 diabetes patients presenting to the institution as part of the routine care was undertaken, between June 2015 and July 2016. Those who attended the diabetes clinic for screening but were not found to be diabetic (HbA1c <6.5%) served as the reference group. Fasting and post-prandial blood glucose, HbA1c levels and serum albumin levels were measured. Spearman correlation and multiple linear regression analysis was done to test the dependence of albumin on HbA1c, fasting blood sugar as well as post-prandial blood sugar levels. The values of HbA1c were higher when the levels of albumin were low and lower when the level of albumin level was high. Serum albumin level was significantly lower in the reference group of patients when compared with control group and correlated negatively with HbA1C levels. Although a negative correlation was found between the levels of HbA1c and albumin in control and reference groups, the relationship between the variables was weak. Since HbA1c has become the standard tool for monitoring glycaemic control in patients with diabetes, the values should be interpreted with caution in our population.

KEYWORDS: Type 2 diabetes mellitus - serum albumin – HbA1c – blood glucose - South India



C. SUTHAKARAN*

Associate Professor Department of Pharmacology Melmaruvathur Adhi Parasakthi Institute of Medical Sciences & Research, Melmaruvathur 603319, Kancheepuram Dist, Tamilnadu, India

Received on: 02.12.2016

Revised and Accepted on: 23.01.2017

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.1.b638-640>

INTRODUCTION

India has emerged as one of the epicentres of the global diabetes mellitus pandemic. Rapid socioeconomic development and demographic changes, along with increased susceptibility for Indian individuals, have led to a dangerous increase in the prevalence of diabetes mellitus in India over the past four decades.¹ Type 2 diabetes mellitus in Asian Indian people is characterized by a young age of onset and occurrence at low levels of BMI. Available data also suggest that the susceptibility of Asian Indian people to the complications of diabetes mellitus differs from that of white populations.² Glycated haemoglobin (HbA1c) is a commonly employed and extensively validated marker for the assessment of glycaemic control in the routine management of diabetes since it provides an accurate, precise measure of chronic glycaemic levels and correlates with the risk of diabetes complications.³ Although HbA1c has established itself as a valuable tool, it is expensive and time consuming to perform. Further, there are a number of clinical conditions which may affect the accuracy of the test, resulting in falsely high or low readings.⁴ Despite a significant negative correlation existing between plasma albumin levels and HbA1c in type 2 diabetes, levels of serum albumin are not routinely monitored in patients with diabetes⁵. Hence, the present study was designed to investigate whether a correlation exists between HbA1c and serum albumin levels in South Indian patients with type 2 diabetes mellitus.

PATIENTS & METHODS

Study design

This study was a prospective study of case records undertaken at a rural tertiary care hospital in southern India, between June 2015 and July 2016. Studies were performed at no extra cost to the participants and patients were not compensated.

Patients

The electronic records of type 2 diabetes patients presenting to the institution as part of the routine care of our centre were reviewed. Patients with type 2 diabetes formed the study group, whereas those who attended

the diabetes clinic for screening but were not found to be diabetic (HbA1c <6.5%) served as the reference group. The inclusion was as broad as possible in order to maximize generalization and to reflect the 'real-world' conditions. The data were extracted using a standardized data extraction form. The study was approved by the Institutional Ethics Committee and written informed consent was obtained from all participants.

Assessments

A fasting blood sample was taken after ensuring a minimum of 8 h of overnight fasting for the estimation of glucose (FPG), HbA1c and albumin. Blood sample was also collected for measurement of postprandial glucose (PPG) concentration. HbA1c was estimated by high-performance liquid chromatography using a Variant machine (Bio-Rad, Hercules, CA). Albumin was estimated by colorimetric assay by dye-binding method, with bromocresol green.

STATISTICAL ANALYSIS

Continuous variables were expressed as mean \pm SD, and categorical variables as the number and percentage. Student's t-test and chi square-test were used for comparisons between sex and groups. Analysis of variance (ANOVA) was applied for comparisons. The correlations among A1C, fasting plasma glucose (FPG), and 2-h post-prandial plasma glucose were determined using Spearman correlations. Multiple linear regression analysis was done to test the dependence of albumin on HbA1c, fasting blood sugar as well as post-prandial blood sugar levels. $P < 0.05$ was considered as statistically significant.

RESULTS

The demographic and clinical characteristics of the Reference and Study groups are shown in Table 1. Both groups were comparable at baseline, and there were no significant differences with respect to age, sex, weight, and BMI or waist circumference except for duration of diabetes.

Table 1
Demographic and Clinical Characteristics

Characteristic	Reference Group (n= 26)	Study Group (n=91)
Age (years)	55.3 \pm 8.9	57.5 \pm 9.7(NS)
Women (%)	15 (58)	55 (60)(NS)
Body weight (kg)	66.8 \pm 9.1	70.1 \pm 12.3(NS)
Body mass index kg/m ²)	24.7 \pm 3.0	26.4 \pm 4.4(NS)
Waist Circumference (cm)	90.7 \pm 8.8	93.0 \pm 11.9(NS)
Duration of diabetes (years)	--	10.9 \pm (4.1)
More than 5 years (%)		47 (52)
Less than 5 years (%)		44 (48)

NS = Not significant

Correlation among A1c albumin FPG and PPG

The values of HbA1c values were higher when the levels of albumin were low and lower when the albumin levels were high (Table 2). Serum albumin level was significantly lower in the reference group of patients when compared with control group and correlated

negatively with HbA1C levels. Although a negative correlation was found, the relationship between the variables was weak ($r = -0.1051$ and -0.1274). The value of R^2 the co-efficient determination was 0.011 and the results were not significant. Stepwise multiple

regression analysis showed no significant correlation between HbA1c and albumin or post prandial blood

sugar or duration of diabetes.

Table 2
Biochemical Characteristics of Study and Reference Groups

Characteristic	Reference Group (n= 26)	Study Group (n=91)
Fasting Blood sugar (mmol/L)	05.21 ± 0.7	10.35 ± 3.91***
Postprandial Blood Sugar (mmol/L)	10.35 ± 1.25	15.48 ± 5.30***
HbA1c (%)	4.4 ± 1.1	9.4 ± 1.9***
Albumin (g/dL)	5.3 ± 0.6	4.1 ± 0.9***

ANOVA $F = 10.99$ $p < 0.001$

*** $P < 0.001$

DISCUSSION

The present study designed to investigate whether a correlation exists between HbA1c and serum albumin levels in South Indian patients with type 2 diabetes mellitus found serum albumin levels to be low. The finding of low albumin levels could be attributed to predominantly vegetarian food which is low in proteins. Fall in HbA1c levels associated with increase in albumin levels implies that that the glucose control of patients with albumin levels significantly above or below average may not be properly reflected by the standard classification in terms of HbA1c measurements alone. Since HbA1c has become the standard tool for monitoring glycaemic control in patients with diabetes⁶, the values should be interpreted with caution in our population. There was no significant correlation between HbA1c and serum albumin or post prandial blood sugar levels. However a positive correlation was observed between HbA1c and serum albumin levels and age as well as duration of diabetes in studies conducted by Arnetz and Kilpatrick^{7,8} while Kabadi found no significant relationship between age, duration of

diabetes and fasting blood glucose, glycated haemoglobin, glycated protein or glycated albumin.⁹ In contradiction to this there was a significant negative correlation between HbA1c and serum albumin.^{5,10}

CONCLUSION

In conclusion our study shows that HbA1c values should be interpreted with caution in our population. To our knowledge, the current study is the first study to report the relationship between HbA1c and albumin levels in South Indian predominantly Tamil speaking patients with type 2 diabetes mellitus. Since the number studied is small, it may not be possible to draw generalised conclusions; hence studies carried out at multiple centres across India, are warranted and may throw light on diverse Indian population.

CONFLICT OF INTEREST

Conflict of Interest declared none.

REFERENCES

- Krishnan V, Kannan R. Physical and psychological determinants of diabetes associated depression among patients receiving oral Hypoglycemic agents versus insulin regimen. Int J Pharm Bio Sci 2016 Oct ; 7(4): (P) 11 - 14
- Unnikrishnan R, Anjana RM, Mohan V. Diabetes mellitus and its complications in India. Nat Rev Endocrinol. 2016 Jun;12(6):357-70.
- Incani M, Sentinelli F, Perra L, Pani MG, Porcu M, Lenzi A et.al. Glycated hemoglobin for the diagnosis of diabetes and prediabetes: Diagnostic impact on obese and lean subjects, and phenotypic characterization. J Diabetes Investig. 2015 Jan;6(1):44-50.
- Welsh KJ, Kirkman MS, Sacks DB. Role of Glycated Proteins in the Diagnosis and Management of Diabetes: Research Gaps and Future Directions. Diabetes Care. 2016 Aug; 39(8): 1299-306.
- Tiwari S, Bothale M, Hasan I, Kulkarni MJ, Sayyad MG, Basu R, et al. Association between serum albumin and glycated haemoglobin in Asian Indian subjects. Indian J Endocr Metab 2015 Jan-Feb; 19(1):52-5.
- Radin MS. Pitfalls in hemoglobin A1c measurement: when results may be misleading. J Gen Intern Med. 2014 Feb;29(2):388-94.
- Arnetz BB, Kallner A, Theorell T. The influence of aging on hemoglobin A1c (HbA1c). J Gerontol. 1982 Nov;37(6):648-50.
- Kilpatrick ES, Dominiczak MH, Small M. The effects of aging on glycation and the interpretation of glycaemic control in type 2 diabetes. (1996). Q J Med. 1996 Apr;89(4): 307-312.
- Kabadi UM. Glycosylation of proteins. Lack of influence of aging. Diabetes Care. 1988 May;11(5):429-32.
- Rodríguez-Segade S, Rodríguez J, Mayan D, Camiña F. Plasma albumin concentration is a predictor of HbA1c among type 2 diabetic patients, independently of fasting plasma glucose and fr. Diabetes Care. 2005Feb;28(2):437-9.