



CHEMICALS AS ADDITIVES IN FOOD PROCESSING -A REVIEW

KUMARESAN D¹, NITHYA SERMUGAPANDIAN*, HEMASHREE S², RUBINI K R²

School of Pharmaceutical Sciences, Vels University (VISTAS), Velan Nagar, P.V. Vaithyalingam Road, Pallavaram, Chennai - 600 117. Tamil Nadu, India.

ABSTRACT

Food additives are edible substances added to food products to enhance their taste, color, appearance and texture. Besides improving quality, these additives are added to prevent microbial growth and increase longevity of processed food products. This review have the aim in giving the outline of common food additives and their devastating effects to our body as a public concern. "Food additive" is any substance their intended use of which results, directly or indirectly, in it's becoming a component of or otherwise affecting the characteristic of any food, and which is safe under the condition of its use. Many literature shows about the problems faced with the use of food additives. Hence in this review the effects of various additives are discussed.

KEYWORDS: Food Additives, Edible Substances, Food Processing, Devasting effects



NITHYA SERMUGAPANDIAN*

School of Pharmaceutical Sciences, Vels University (VISTAS), Velan Nagar, P.V. Vaithyalingam Road, Pallavaram, Chennai - 600 117. Tamil Nadu, India.

Received on: 06-10-2016

Revised and Accepted on : 30-01-2017

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.1.p276-282>

INTRODUCTION

Most food that comes in a packet or container contains at least some food additives¹⁻². Especially when it comes to the names and numbers of food additives, which are often completely unidentifiable and unpronounceable. Being one of the most populous countries in the world, India has been witnessing significant rise in food consumption level³, prevalence of processed and packaged food, convenience food, alcoholic and non-alcoholic beverages, bakery items, ready-to-eat food products and cereals^{4,2}. These trends are expected to drive remarkable growth in the demand for food additives in the country over the coming years⁵. According to the recently published Tech Sci Research report "India Food Additives Market Forecast and Opportunities, 2020", the market for food additives in India is anticipated to witness double digit growth at a Compound annual growth Rate CAGR over 12% during 2015-2020⁴. Growth from the capital invested disposable income and rising urban middle class population are among the key market contributors. Due to cultural and regional diversities, food consumption pattern is very not uniform in various parts of the country. As a result, western region generates the highest demand for food additives in the country. The region is expected to continue spearheading the market due to the presence of a large number of food and beverage manufacturing companies in the states of Gujarat and Maharashtra.⁶ Firmenich Aromatics India Private Limited, Zydus, Cadila, Danisco and Adani Wilmer Limited are some of the major food additives suppliers operating in India. Global food giants such as Mars, Mondelez, PepsiCo, Inc. and Kraft Foods have also been supplying their food items and are among the major food additives consumers in India^{7,8}. "Indian food additives market is dominated by the flavors segment due to their substantial use in beverages, savory items and cereals⁹⁻¹¹. With rising incidences of health problems like diabetes, consumers are also shifting towards all-natural, low calorie sweeteners such as Stevia¹²⁻¹³. As a result, sweeteners are most likely to witness faster growth over the next five years and emerge as the leading segment in India food additives market through 2020¹⁴⁻¹⁵."

Needs of Food Additives

Additive ingredients helps to preserve and improve the taste of food for centuries and without certain additives many processed foods would be unsafe to eat, and if they weren't we wouldn't want to eat them anyway¹⁶⁻¹⁸. Some of the functions of food additives include:

- adding or restoring colour to foods (artificial colours have code numbers in 100s)
- preventing food from 'going off' (preservatives have code numbers in 200s)
- slowing or preventing the oxidative deterioration of foods (antioxidants have code numbers in 300s)
- Improving the flavour of food (food enhancers have code numbers in the 600s).

"Consumers expect certain types of food to be certain types of colour so for example, if there are no colours we'd be eating clear margarine."

Problems of Food additives

Preservatives have been associated with intolerances, particularly among people with asthma. Sulfites (including sodium bisulphite (222), sodium metabisulphite (223) and potassium bisulphite (228)) found in wine, beer and dried fruit, are known to trigger asthmatic episodes and cause migraines in people who are sensitive to them¹⁹. Also sodium nitrate (251) and sodium nitrite (250), which are used in processed meats, have been classified as 'probably carcinogenic to humans' by the International Agency for Research of Cancer (IARC)²⁰⁻²¹. Flavour enhancer monosodium glutamate (MSG) (621) is often used in Asian cooking and has been associated with 'Chinese Restaurant Syndrome' (a collection of symptoms including headache, numbness and tingling that some people experienced after eating foods containing MSG)²²⁻²⁴. While numerous studies have found MSG is safe for most of us, some people experience symptoms if they eat a large amount of MSG in a single meal²⁵. Food colourings, such as tartrazine (102), allura red (129) and ponceau 4R (124), are often credited as the cause of hyperactivity in children²⁶.

Natural additives

The distinction between 'artificial' and 'natural' food additives, is misguided because almost everyone who is sensitive to an artificial additive will also be sensitive to one or more natural substances²⁷⁻²⁸. "Sometimes the additives and the natural substance are chemically identical, sometimes they're chemically closely related and the distinction between natural and artificial is completely artificial." Swain says the main difference between these compounds in 'natural' versus processed foods is concentration. However, she stresses that for the majority of people, the nature and levels of additives in our food supply are not an issue as long as they are consumed in moderation²⁹.

Commonly used chemicals as additives

Chemical food additives preserve the shelf life by reducing or eliminating the growth of microorganisms that causes food decay. These added chemicals are controversial as some may have a detrimental effect on health. Eating more fresh, whole foods may be safe alternative for use of these chemicals. The commonly used additives in food products includes^{8,26}

- a. Sweeteners-Aspartame, Saccharine
- b. Artificial colours- Bixin, Carotene, Saffron
- c. Flavours and Flavour Enhancers-MSG
- d. Antioxidants- BHA & BHT
- e. Preservatives- Nitrites, nitrates, Sulfites, Benzoic acid and sodium benzoate
- f. Flouring agents- Potassium bromate

EFFECTS OF ADDITIVES

Sweetners- Aspartame, Saccharine

Aspartames are sweeteners they add sweetness with or without the extra calories. The artificial sweetener aspartame (L-aspartyl-L-phenylalanyl-methyl ester), is consumed, primarily in beverages, causing significant elevations in plasma and brain phenylalanine levels few reports suggest that some people suffer neurologic or behavioral reactions in association with aspartame

consumption³⁰⁻³¹. Since phenylalanine can be neurotoxic and can affect the synthesis of inhibitory monoamine neurotransmitters, the phenylalanine in aspartame could conceivably mediate neurologic effects. The metabolites of aspartame include aspartate, methanol, phenylalanine and phenylketoneuria³²⁻³³. Strong link between intake of sugary beverages and overweight in children and the Sweeteners also found to increase appetite thereby by causing weight gain in adults³⁴.

Artificial Colours- Bixin, Carotene, Saffron

Food coloring, or color additive, is any dye, pigment or substance that imparts color when it is added to food or drink. They come in many forms consisting of liquids, powders, gels, and pastes. Offset color loss due to exposure to light, air, temperature extremes, moisture and storage conditions; correct natural variations in color; enhance colors that occur naturally; provide color to colorless and "fun" foods³⁵. These include colour stabilizers, colour fixatives, colour retention agents, etc. They consist of synthetic colours, and Natural Colours colours from . Even though colours add nothing to the nutritive value of foods, without certain colours most consumers will not feel them attractive to buy or eat some foods without colours¹⁵. Thus, colours are frequently added to restore the natural ones lost in food processing or to give the preparations the natural colour we expect. A number of natural food colours extracted from seeds, flowers, insects, and foods, are also used as food additives. One of the best known and most widespread red pigment is bixin, derived from the seed coat of *Bixa orellana*, the lipstick pod plant of South American origin. Bixin is not considered to be carcinogenic. The major use of this plant on a worldwide basis, however, is for the annatto dye, a yellow to red colouring material extracted from the orange red pulp of the seeds. Annatto has been used as colouring matter in butter, cheese, margarine, and other foods. Another yellow colour, a carotene derived from carrot, is used in margarine. Saffron has both flavouring and colouring properties and has been used for colouring foods. Turmeric is a spice that gives the characteristic colour of curries and some meat products and salad dressings. A natural red colour, cochineal (or carnum) obtained by extraction from the female insect (*Coccus cacti*), grape skin extract, and caramel, the brown colour obtained from burnt sugar, are some natural colours that are used as food additives. Food colorings may cause mental disorder of the neurodevelopmental type in childrens^{36,37}. Examples Many processed foods, (candies, snack foods, margarine, cheese, soft drinks, jams/jellies, gelatins, pudding and pie fillings)

Artificial Flavouring- MSG

Add specific flavors (natural and synthetic). Flavour enhancers are not flavours themselves but they amplify the flavours of other substance through a synergistic effect³⁸. Flavour and flavor enhancers constitute the largest class of food additives. Natural flavor are substances, such as spices, herbs, roots, essences, and essential oils, have been used in the past as flavor additives³⁹. The flavours are in short supply and the amount of flavor substances in them is very tiny. It would take about tonne of many spices to produce 1 g of the flavor substances, and in some cases only 0.1 g

can be extracted^{40,41}. Natural food flavours are thus being replaced by synthetic flavor materials. The agents responsible for flavor are esters, aldehydes, ketones, alcohols, and ethers. These substances are easily synthesized and can be easily substituted for natural ones. Typical of the synthetic flavor additives are amyl acetate for banana, aldehyde, hydroxyl and ether for vanillin, methyl anthranilate for grapes, ethyl butyrate for pineapple, etc⁴²⁻⁴³. Generally, most synthetic flavours are mixtures of a number of different substances. For example, one imitation cherry flavor contains fifteen different esters, alcohols, and aldehydes. One of the best known, most widely used and somewhat controversial flavor enhancers is monosodium glutamate (MSG), the sodium salt of the naturally occurring amino acid glutamic acid. This is added to over 10,000 different processed foods. This has been extracted from seaweeds and soya bean⁴⁴. MSG is found to produce attractive meat like flavor. MSG is now manufactured on a large scale all over the world, and especially in Japan. MSG is generally recognized as safe⁴⁵. Few reports suggests for some individuals experience symptoms often comparable to those of heart attack, when served with food containing large amounts of MSG. Yeast extract has the same flavor enhancing property as MSG. It is found that, in this case, the flavor enhancing substances are the ribonucleotides. These are ten times more powerful than MSG⁴⁶. Examples of foods rich in flavouring agents includes Pudding and pie fillings, gelatin dessert mixes, cakemixes, salad dressings, candies, soft drinks, ice cream, BBQ sauce²⁵

Antioxidants- Butylated hydroxyl anisole

These are the antioxidants used in food processing. An antioxidant is a substance added to fats and fat containing substances to retard oxidation and thereby prolong their wholesomeness, palatability, and, sometimes, keeping time⁴⁷. An antioxidant should not contribute an objectionable odour, flavor, or colour, to the fat or to the food in which it is present⁴⁸. It should be effective in low concentrations, and be fat soluble. Also, it should not have a harmful physiological effect. Some antioxidants used in foods are butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), propyl gallate (PG), and tertiarybutyl hydroquinone (TBHQ), which are all phenolic substances. Thioldipropionic acid and dilauryl thioldipropionate are also used as food antioxidants⁴⁹. The Joint FAO/WHO Expert Committee on Food Additives has recently considered the Acceptable Daily Intakes (ADIs) of BHA and BHT and set them at 0-0.5 mg/kg body weight for BHA and 0-0.3 mg/kg body weight for BHT. Naturally occurring substances that act as anti-oxidants are tocopherols. The tocopherols act as biological anti-oxidants in plant and animal tissues, but they are rarely used as additives because they are more expensive than synthetic antioxidants⁵⁰. However when large amounts of antioxidants nutrients are taken, they can also act as pro-oxidants by inducing oxidative stress⁵¹

Preservatives - Benzoic acid and sodium benzoate

A preservative are any substance which is capable of inhibiting, retarding, or arresting, the growth of microorganisms, of any deterioration of food due to microorganisms, or of masking the evidence of any such

deterioration. It is estimated that nearly 1/5 of the world's food is lost by microbial spoilage⁵². Chemical preservatives interfere with the cell membrane of microorganisms, their enzymes, or their genetic mechanisms. The compounds used as preservatives include natural preservatives, such as sugar, salt, acids, etc, as well as synthetic preservative Prevent food spoilage from bacteria, molds, fungi, or yeast (antimicrobials); slow or prevent changes in color, flavor, or texture and delay rancidity (antioxidants); maintain freshness Examples- Fruit sauces and jellies, beverages, baked goods, cured meats, oils and margarines, cereals, dressings, snack foods, fruits and vegetables⁵³. There are certain harmful effects of using chemicals for preservation such as ;Sulfites are common preservatives used in various fruits, may have side effects in form of headaches, palpitations, allergies, and even cancer. Nitrates and Nitrites: These additives are used as curing agents in meat products⁵⁴. It gets converted into nitrous acid when consumed and is suspected of causing stomach cancer. Benzoates are used in foods as antimicrobial preservatives, and have been suspected to cause allergies, asthma and skin rashes. Sorbates/sorbic acid are added to foods⁵⁵ as antimicrobial preservatives. Reactions to sorbates are rare, but have included reports of urticaria and contact dermatitis⁵⁶. Also a nuclear radiation when used for preservation does not make foods radioactive, but may cause changes in food color or texture^{53,57}.

Flour improvers- Potassium bromate

These are bleaching and maturing agents; usually, they both bleach and "mature" the flour. These are important in the flour milling and bread baking industries. Freshly milled flour has a yellowish tint and yields a weak dough that produces poor bread. Both the colour and baking properties improve by storing the flour for several months before making bread. Chemical agents used as flour improvers are oxidizing agents, which may participate in bleaching only, in both bleaching and dough improvement, or in dough improvement only⁵⁸. The agent that is used only for flour bleaching is benzoyl peroxide. This does not influence the quality of dough. Materials used both for bleaching and improving are chlorine gas, chlorine dioxide, nitrosyl chloride, and nitrogen di and tetraoxides⁵⁹. Oxidizing agents used only

for dough improvement are potassium bromate, potassium iodate, Calcium iodate and calcium peroxide. Potassium bromate has been banned in food products because they are potent carcinogens⁶⁰.

Nutrient Supplements – Vitamin C

Nutrient supplements restore values lost in processing or storage, or ensure higher nutritional value than what nature may have provided. When foods are processed, there may be loss of Some nutrients and additives may be added to restore the original value. For example, to produce white flour, wheat is milled in such a way as to remove the brown coloured part of the grain, which is rich in vitamins and minerals⁶¹. To restore the nutritive value, thiamine, nicotinic acid, iron and calcium, are added to the flour. Similarly, vitamin C is added to canned citrus fruits to make up the loss of the vitamin during processing. The added vitamin c will increase the acidic levels in the canned citrus juice⁶².

CONCLUSION

Additive chemicals are used to increase the shelf life of food and to maintain the quality and colour of the finished product for the long period of time.the chemicals that are used as additives have side effects. Hence as a consumer we have to check the label of the consumable products and know about the preservatives colouring agents and other additives added to the product and then select for consuming purpose. . Side reactions of these additives can be immediate or build up in the body over time. Thus it is always best to eat a additive free diet if at all possible as pure as organic from Nature.

ACKNOWLEDGEMENT

We wish to thank, Dr. Ishari K Ganesh, Chancellor, Vels University Chennai, India for providing the facilities necessary to carry out the research work

CONFLICT OF INTEREST

Conflict of interest declared none.

REFERENCES

- Okano K, Ose A, Takai M, Kaneko M, Nishioka C, Ohzu Y, Odano M, Sekiyama Y, Mizukami Y, Nakamura N, Ichinoe M. Inhibition of aflatoxin production and fungal growth on stored corn by allyl isothiocyanate vapor. *Shokuhin eiseigaku zasshi. Journal of the Food Hygienic Society of Japan.* 2014 Dec;56(1):1-7.
- Gardini F, Özogul Y, Suzzi G, Tabanelli G, Özogul F. Technological factors affecting biogenic amine content in foods: a review. *Frontiers in microbiology.* 2016;7.
- Shamansurova Z, Tan P, Ahmed B, Pepin E, Seda O, Lavoie JL. Adipose tissue (P) RR regulates insulin sensitivity, fat mass and body weight. *Molecular Metabolism.* 2016 Oct 31;5(10):959-69.
- Roberts-Gray C, Briley ME, Ranjit N, Byrd-Williams CE, Sweitzer SJ, Sharma SV, Palafox MR, Hoelscher DM. Efficacy of the Lunch is in the Bag intervention to increase parents' packing of healthy bag lunches for young children: a cluster-randomized trial in early care and education centers. *International Journal of Behavioral Nutrition and Physical Activity.* 2016 Jan 8;13(1):3.
- Lee. The Food Hygiene Institutionalization of Park Junghee Government in 1960s. *Korean Journal of Medical History.* 2016 Aug 30;25(2):205-40.
- Lee J, Jeong S, Ko G, Park H, Ko Y. Development of a Food Safety and Nutrition Education Program for Adolescents by Applying Social Cognitive Theory. *Osong Public Health and*

- Research Perspectives. 2016 Aug 31;7(4):248-60.
7. Chunhabundit R. Cadmium Exposure and Potential Health Risk from Foods in Contaminated Area, Thailand. *Toxicological research*. 2016 Jan;32(1):65.
 8. Stern D, Piernas C, Barquera S, Rivera JA, Popkin BM. Caloric beverages were major sources of energy among children and adults in Mexico, 1999–2012. *The Journal of nutrition*. 2014 Jun 1;144(6):949-56.
 9. Serra S. Recent advances in the synthesis of carotenoid-derived flavours and fragrances. *Molecules*. 2015 Jul 15;20(7):12817-40.
 10. de Billerbeck GM. Bioflavour: natural flavours and fragrances by yeasts and other microorganisms—the holistic approach of the BIOFLAVOUR Consortium. *Yeast*. 2015 Jan 1;32:1.
 11. Ozarda O, Demirköz AB, Özdemir M. Sensory characteristics and antioxidant capacity of red raspberry extract as a preservative in fruity flavoured beverages. *Journal of food science and technology*. 2015 Oct 1;52(10):6687-94.
 12. Winkler HC, Suter M, Naegeli H. Critical review of the safety assessment of nano-structured silica additives in food. *Journal of nanobiotechnology*. 2016 Jun 10;14(1):44.
 13. Stobnicka A, Górny RL. Exposure to flour dust in the occupational environment. *International journal of occupational safety and ergonomics*. 2015 Jul 3;21(3):241-9.
 14. Di Ianni ME, del Valle ME, Enrique AV, Rosella MA, Bruno F, Bruno-Blanch LE, Talevi A. Computer-aided identification of anticonvulsant effect of natural nonnutritive sweeteners stevioside and rebaudioside A. *Assay and drug development technologies*. 2015 Jul 1;13(6):313-8.
 15. Di Ianni ME, del Valle ME, Enrique AV, Rosella MA, Bruno F, Bruno-Blanch LE, Talevi A. Computer-aided identification of anticonvulsant effect of natural nonnutritive sweeteners stevioside and rebaudioside A. *Assay and drug development technologies*. 2015 Jul 1;13(6):313-8.
 16. Keraka M, Ochieng C, Engelbrecht J, Hongoro C. Association between the use of biomass fuels on respiratory health of workers in food catering enterprises in Nairobi Kenya. *The Pan African medical journal*. 2013;15.
 17. Tassew, H., Abdissa, A., Beyene, G. & Gebre-Selassie, S. Microbial flora and food borne pathogens on minced meat and their susceptibility to antimicrobial agents. *Ethiop. J. Heal. Sci.* 2010 **20**, 137–143 .
 18. Burt BM, Volel C, Finkel M. Safety of vendor-prepared foods: evaluation of 10 processing mobile food vendors in Manhattan. *Public Health Reports*. 2003 Sep 1;118(5):470-6.
 19. Bezerra RM, Veiga LF, Caetano AC, Rosalen PL, Amaral ME, Palanch AC, de Alencar SM. Caffeic acid phenethyl ester reduces the activation of the nuclear factor κB pathway by high-fat diet-induced obesity in mice. *Metabolism*. 2012 Nov 30;61(11):1606-14.
 20. Mania M, Szynal T, Rebeniak M, Wojciechowska-Mazurek M, Starska K, Strzelecka A. Human exposure assessment to different arsenic species in tea. *Roczniki Państwowego Zakładu Higieny*. 2014;65(4).
 21. Jones JA, Hopper AO, Power GG, Blood AB. Dietary intake and bio-activation of nitrite and nitrate in newborn infants. *Pediatric research*. 2014 Oct 14;77(1-2):173-81.
 22. Kadeerbai HS, Liu M, Dai XQ, Li XG, Li YL, Yang XS, Lu BH, Wang ZH, Zhu FX, Wu H, Zhao X. Analysis of disease types of outpatients with intestinal diseases from two grade-three general hospitals in Beijing. *Beijing da xue xue bao. Yi xue ban= Journal of Peking University. Health sciences*. 2010 Jun;42(3):308-13.
 23. Gaüzère BA, Djardem Y, Bourdé A, Blanc P. A new chinese restaurant syndrome: the 'chinese fondue' carbon monoxide mass intoxication. *Crit Care*. 1999 Mar 15;3:55.
 24. Chow EP, Lau JT, Zhuang X, Zhang X, Wang Y, Zhang L. HIV prevalence trends, risky behaviours, and governmental and community responses to the epidemic among men who have sex with men in China. *BioMed research international*. 2014 Apr 14;2014.
 25. Ostrosky-Zeichner L, Shoham S, Vazquez J, Reboli A, Betts R, Barron MA, Schuster M, Judson MA, Revankar SG, Caeiro JP, Mangino JE. MSG-01: a randomized, double-blind, placebo-controlled trial of caspofungin prophylaxis followed by preemptive therapy for invasive candidiasis in high-risk adults in the critical care setting. *Clinical infectious diseases*. 2014 May 1;58(9):1219-26.
 26. Arnold LE, Lofthouse N, Hurt E. Artificial food colors and attention-deficit/hyperactivity symptoms: conclusions to dye for. *Neurotherapeutics*. 2012 Jul 1;9(3):599-609.
 27. Darwish HW, Attia MI, Abdelhameed AS, Alanazi AM, Bakheit AH. Comparative ANNs with different input layers and GA-PLS study for simultaneous spectrofluorimetric determination of melatonin and pyridoxine HCl in the presence of melatonin's main impurity. *Molecules*. 2013 Jan 14;18(1):974-96.
 28. Hahn A, Fuhlrott J, Loos A, Barcikowski S. Cytotoxicity and ion release of alloy nanoparticles. *Journal of Nanoparticle Research*. 2012 Jan 1;14(1):686.
 29. Shawle, K., Urge, M. & Animut, G. Effect of different levels of *Lepidium sativum* L. on growth performance, carcass characteristics, hematology and serum biochemical parameters of broilers. *Springerplus* 2016 **5**, 1441 .
 30. Bhattacharyya S, Feferman L, Unterman T, Tobacman JK. Exposure to common food additive carrageenan alone leads to fasting hyperglycemia and in combination with high fat diet exacerbates glucose intolerance and hyperlipidemia without effect on weight. *Journal of diabetes research*. 2015 Mar 25;2015.
 31. Elson KM, Fox N, Tipper JL, Kirkham J, Hall RM, Fisher J, Ingham E. Non-destructive monitoring of viability in an ex vivo organ culture model of

- osteochondral tissue. *eCells and Materials Journal*. 2015 Jun 30;29:356-69.
32. Ashok I, Sheeladevi R, Wankhar D. Acute effect of aspartame-induced oxidative stress in Wistar albino rat brain. *Journal of biomedical research*. 2015 Sep;29(5):390.
 33. Palmnäs MS, Cowan TE, Bomhof MR, Su J, Reimer RA, Vogel HJ, Hittel DS, Shearer J. Low-dose aspartame consumption differentially affects gut microbiota-host metabolic interactions in the diet-induced obese rat. *PLoS one*. 2014 Oct 14;9(10):e109841.
 34. Chan TF, Lin WT, Huang HL, Lee CY, Wu PW, Chiu YW, Huang CC, Tsai S, Lin CL, Lee CH. Consumption of sugar-sweetened beverages is associated with components of the metabolic syndrome in adolescents. *Nutrients*. 2014 May 23;6(5):2088-103.
 35. Cao F, Liu T, Xu Y, Xu D, Feng S. Curcumin inhibits cell proliferation and promotes apoptosis in human osteoclastoma cell through MMP-9, NF- κ B and JNK signaling pathways. *International journal of clinical and experimental pathology*. 2015;8(6):6037.
 36. Adusumilli H, Avula JS, Kakarla P, Bandi S, Mallela GM, Vallabhaneni K. Color stability of esthetic restorative materials used in pediatric dentistry: An in vitro study. *Journal of Indian Society of Pedodontics and Preventive Dentistry*. 2016 Jul 1;34(3):233.
 37. Ruiz R, Peinado MJ, Aranda-Olmedo I, Abecia L, Suárez-Pereira E, Ortiz Mellet C, García Fernández JM, Rubio LA. Effects of feed additives on ileal mucosa-associated microbiota composition of broiler chickens. *Journal of animal science*. 2015 Jul 1;93(7):3410-20.
 38. Pogue AI, Lukiw WJ. Natural and Synthetic Neurotoxins in Our Environment: From Alzheimer's Disease (AD) to Autism Spectrum Disorder (ASD). *Journal of Alzheimer's disease & Parkinsonism*. 2016 Aug;6(4).
 39. Wang ZM, Lu ZM, Shi JS, Xu ZH. Exploring flavour-producing core microbiota in multispecies solid-state fermentation of traditional Chinese vinegar. *Scientific reports*. 2016;6.
 40. Li C, Sun J, Fu C, Yu B, Liu SQ, Li T, Huang D. Synthesis and evaluation of odour-active methionyl esters of fatty acids via esterification and transesterification of butter oil. *Food chemistry*. 2014 Feb 15;145:796-801.
 41. Schlegel, W. Nutrition and flavour-legislation. *Nahr*. 1976 **20**, 253–258
 42. Faber F, Tran L, Byndloss MX, Lopez CA, Velazquez EM, Kerrinnes T, Nuccio SP, Wangdi T, Fiehn O, Tsohis RM, Bäumlner AJ. Host-mediated sugar oxidation promotes post-antibiotic pathogen expansion. *Nature*. 2016 Jun 15.
 43. Stadlbauer V, Haselgrübler R, Lanzerstorfer P, Plochberger B, Borgmann D, Jacak J, Winkler SM, Schröder K, Höglinger O, Weghuber J. Biomolecular Characterization of Putative Antidiabetic Herbal Extracts. *PLoS one*. 2016 Jan 28;11(1):e0148109.
 44. Faucitano L, Berthiaume R, D'Amours M, Pellerin D, Ouellet DR. Effects of corn grain particle size and treated soybean meal on carcass and meat quality characteristics of beef steers finished on a corn silage diet. *Meat science*. 2011 Aug 31;88(4):750-4.
 45. Brosnan JT, Drewnowski A, Friedman MI. Is there a relationship between dietary MSG obesity in animals or humans?. *Amino acids*. 2014 Sep 1;46(9):2075-87.
 46. Sobrevals L, Müller P, Fabra A, Castro S. Role of glutathione in the growth of *Bradyrhizobium* sp.(peanut microsymbiont) under different environmental stresses and in symbiosis with the host plant. *Canadian journal of microbiology*. 2006 Jul 1;52(7):609-16.
 47. Garcimartín A, Santos-López JA, Bastida S, Benedí J, Sánchez-Muniz FJ. Silicon-Enriched Restructured Pork Affects the Lipoprotein Profile, VLDL Oxidation, and LDL Receptor Gene Expression in Aged Rats Fed an Atherogenic Diet. *The Journal of nutrition*. 2015 Sep 1;145(9):2039-45.
 48. Kumar D, Kumar S, Singh J, Vashistha BD, Singh N. Free radical scavenging and analgesic activities of *Cucumis sativus* L. fruit extract. *Journal of Young Pharmacists*. 2010 Oct 1;2(4):365-8.
 49. Bekele RT, Venkatraman G, Liu RZ, Tang X, Mi S, Benesch MG, Mackey JR, Godbout R, Curtis JM, McMullen TP, Brindley DN. Oxidative stress contributes to the tamoxifen-induced killing of breast cancer cells: implications for tamoxifen therapy and resistance. *Scientific reports*. 2016;6.
 50. Chen YF, Shibu MA, Fan MJ, Chen MC, Viswanadha VP, Lin YL, Lai CH, Lin KH, Ho TJ, Kuo WW, Huang CY. Purple rice anthocyanin extract protects cardiac function in STZ-induced diabetes rat hearts by inhibiting cardiac hypertrophy and fibrosis. *The Journal of nutritional biochemistry*. 2016 May 31;31:98-105.
 51. Weaver AC, See MT, Kim SW. Protective effect of two yeast based feed additives on pigs chronically exposed to deoxynivalenol and zearalenone. *Toxins*. 2014 Dec 12;6(12):3336-53.
 52. Yu HS, Lee NK, Jeon HL, Eom SJ, Yoo MY, Lim SD, Paik HD. Benzoic Acid Production with Respect to Starter Culture and Incubation Temperature during Yogurt Fermentation using Response Surface Methodology. *Korean Journal for Food Science of Animal Resources*. 2016;36(3):427.
 53. di Gesso JL, Kerr JS, Zhang Q, Raheem S, Yalamanchili SK, O'hagan D, Kay CD, O'connell MA. Flavonoid metabolites reduce tumor necrosis factor- α secretion to a greater extent than their precursor compounds in human THP-1 monocytes. *Molecular nutrition & food research*. 2015 Jun 1;59(6):1143-54.
 54. Silva F, Domingues FC, Nerín C. Trends in microbial control techniques for poultry products. *Critical Reviews in Food Science and Nutrition*. 2016 Jul 12(just-accepted):00-.
 55. Janjarasskul T, Suppakul P. Active and Intelligent Packaging: The Indication of Quality and Safety. *Critical Reviews in Food Science and Nutrition*. 2016 Sep 10(just-accepted):00-.

56. Kieć-Świerczyńska M, Świerczyńska-Machura D, Chomiczewska-Skóra D, Nowakowska-Świrta E, Kręcisz B. Occupational allergic and irritant contact dermatitis in workers exposed to polyurethane foam. *International journal of occupational medicine and environmental health*. 2014 Apr 1;27(2):196-205.
57. Pongsavee M. Effect of sodium benzoate preservative on micronucleus induction, chromosome break, and Ala40Thr superoxide dismutase gene mutation in lymphocytes. *BioMed research international*. 2015 Feb 17;2015.
58. Leboda R, Skubiszewska-Zięba J, Charmas B, Chodorowski S, Pokrovskiy VA. Carbon–mineral adsorbents from waste materials: case study. *Journal of colloid and interface science*. 2003 Mar 1;259(1):1-2.
59. Andrade-Mahecha MM, Pelissari FM, Tapia-Blácido DR, Menegalli FC. Achira as a source of biodegradable materials: Isolation and characterization of nanofibers. *Carbohydrate polymers*. 2015 Jun 5;123:406-15.
60. Kurokawa Y, Maekawa A, Takahashi M, Hayashi Y. Toxicity and carcinogenicity of potassium bromate--a new renal carcinogen. *Environmental health perspectives*. 1990 Jul;87:309.
61. Parnell JA, Wiens KP, Erdman KA. Dietary Intakes and Supplement Use in Pre-Adolescent and Adolescent Canadian Athletes. *Nutrients*. 2016 Aug 26;8(9):526.
62. Ghorbani, Z., Hajizadeh, M. & Hekmatdoost, A. Dietary supplementation in patients with alcoholic liver disease: a review on current evidence. *Hepatobiliary Pancreat. Dis. Int. Hbpd Int* 2016 15; 348–360