



OPTIMIZATION OF BIOSORPTIVE REMOVAL OF COBALT(II) ONTO *EUCALYPTUS GLOBULUS L.*

D. HYMAVATHI¹ AND DR G. PRABHAKAR^{*2}

^{1, *2}Department of Chemical Engineering, S. V. University College of Engineering, S. V. University,
Tirupati – 517502, Andhra Pradesh, India

ABSTRACT

The objective of this study is to evaluate the efficacy of *Eucalyptus globulus* leaf powder in the removal of Cobalt(II) from aqueous solutions. Detailed batch mode experimentation using Response Surface Methodology (RSM) is undertaken and at optimum conditions, the maximum biosorption capacity of *E. globulus* leaf powder for Cobalt(II) loading is $q_{\max} = 7.194$ mg/g. Freundlich equilibrium model is identified for adsorption process, after trying three standard isotherm models - Langmuir, Freundlich and Temkin isotherms. Pseudo-second-order kinetic model describes the adsorption process. Surface morphology studies through Fourier Transform Infrared Spectroscopy[FTIR] and Scanning Electron Microscopy [SEM] with Energy Dispersive X-ray spectroscopy(EDX) confirms the metal loading on the adsorbent surface.

KEYWORDS: Cobalt(II) removal, Biosorption, *E. globulus L.*, Equilibrium, Kinetics, RSM



DR G. PRABHAKAR

Department of Chemical Engineering, S. V. University College of Engineering, S. V. University,
Tirupati – 517502, Andhra Pradesh, India

Received on: 29.11.2016

Revised and Accepted on 13-02-2017

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.2.b107-115>

INTRODUCTION

Quality water is essential for the well being of all living systems. Unfortunately, its supply is diminishing due to continuous, unwanted interference with the ecosystem. The problem is compounded by the discharge of toxic exit streams of different processing industries, making water unfit for reuse. One of the major water pollutants is heavy metals- Co, Pb, Cr, Zn, Cd, Cu, Hg, As, Al and Ni, discharged by various industrial units. Levels of these ions in water are high enough to cause several health issues. Hence, there is a need to treat the water; to make it potable and adsorption is a good treatment option, because of its efficacy and efficiency^{1, 2}. Numerous techniques available for removal of heavy metals from industrial waste water are chemical precipitation, reverse osmosis, ion exchange, coagulation, electro dialysis, ultrafiltration³⁻⁸, but they are costlier and inadequate compared to adsorption. Biosorption of Cobalt(II) using different forms of agriculture waste material were tried earlier and the sorbent materials studied are functional ligand anchored nanomaterial (facial adsorbent)⁹, Chrysanthemum indicum¹⁰, leaves of *Acacia nilotica*¹¹, fruit wastes of *Blighia sapida* (Akee apple) pod¹², *ficus religiosa*¹³ lemon¹⁴⁻¹⁶, lichen *Evernia prunastri*¹⁷, rose waste biomass¹⁸, crab and arca shell¹⁹, Black carrot residues²⁰, crab²¹, hazelnut shell²², fruit peels of banana and orange²³. Eucalyptus leaves contain essential oils like α -pinene, 1, 8-cineol and pinocarveol-trans at a level of 1.2% to 3% (w/w). The oil contains the highest antibacterial activity against microbes of *Listeria ivanovii* and *Bacillus cereus*²⁴⁻²⁶. In the present study *E. globulus* leaf powder is the sorbent chosen for the removal of Cobalt(II) and parametric dependence, equilibrium and kinetic modeling and surface modification characteristics are reported.

MATERIALS AND METHODS

Chemicals

All used chemical compounds are of analytic grade (Merck, Germany). A stock solution of Cobalt(II) (0.254mg/100mL) is prepared by dissolving 0.254 mg of 97% $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ in 100 mL of distilled water. The stock solution is appropriately diluted to obtain desired concentrations of Cobalt(II) ion. pH of the metal solution is adjusted with 0.1 N NaOH and 0.1N H_2SO_4 solutions.

Preparation and Treatment of biosorbent

E. globulus leaves are collected from Sri Venkateswara University Campus, Tirupati, washed several times to remove impurities and then dried under sunlight. Dried leaves are ground to fine powder, using a mechanical grinder, to 240(63 μm) mesh B.S.S. These particles are again washed with distilled water, dried at room temperature and stored in air tight bottles for further experimental work.

Biosorption procedure

Adsorption of cobalt with *E. globulus L.* is analyzed in a batch mode. 50 mL of solution having 20 mg/L of cobalt

ion is taken into a 250 mL Erlenmeyer flask. 10 g/L of bio sorbent of 63 μm size are added to the flasks. Initial solution pH is adjusted by adding 0.1 N NaOH / 0.1 N sulfuric acid. Samples are agitated on an orbital shaker at constant speed and at room temperature. At different time periods (1, 3, 5, 10..... 120 & 150 min), samples are collected, filtered and the filtrate is analyzed for Cobalt(II) concentration, using inductively coupled plasma optical emission spectroscopy (Optima 8000). It is observed that 60-minute contact time will take the procedure to equilibrium. The % Removal of Cobalt(II) is calculated using equation 1.

$$\frac{c_0 - c_e}{c_0} * 100 \quad (1)$$

Effects of four factors - Cobalt(II) concentration, initial solution pH, adsorbent dosage and temperature of Cobalt(II) solution are then studied, keeping the particle size and processing time fixed. The equilibrium metal uptake capacity (q_e) is estimated by using the following equation 2.

$$q_e = \frac{c_0 - c_e}{m} * V \quad (2)$$

Where c_0 is initial Cobalt(II) concentration(mg/L), c_e is equilibrium Cobalt(II) concentration(mg/L), m is mass of adsorbent(g), V is volume of Cobalt(II) solution(ml)

Characterization of biosorbent

FT-IR (ALPHA interferometer (ECO-ATR)) Germany Spectroscopy, is used to record the IR spectrum before and after adsorption process in the range of 4000–500 cm^{-1} and to identify the functional groups involved in retention of Cobalt(II) ion. The structure and morphology of *E. globulus L.* before and after adsorption of Cobalt(II) is observed using a scanning electron microscope (SEM–EVO MA 15).

RESULTS AND DISCUSSIONS

Optimization using Response Surface Methodology (RSM)

The contribution and inter relationship among the four independent variables and on the percentage removal of Cobalt(II) ions is studied and is fitted with a quadratic model using Design expert 10.0.3.0. Results from RSM (CCD) for Cobalt(II) biosorption by *E. globulus L.* are as shown in Table 1 and Levels of different process variables in coded and un coded form for biosorption of Cobalt(II) using *E. globulus L.* leaf powder as shown in Table 2. ANOVA and estimated regression coefficients for the Cobalt(II) biosorption onto *E. globulus L.* are in Table 3. A comparison of optimal value (predicted) and actual percentage (experimental) values of Cobalt(II) and the normal plot for residuals are given in the form of Fig.1 and Surface contour plots for sorption of Cobalt (II) onto *E. globulus* leaf powder are grouped in to Fig.2. When a solution of 20 mg/L Co(II) is treated with 25 g/L *E. globulus L.* at a pH of 5.0 and temperature of 303 K, 80.849 % of Cobalt(II) present is removed, the following quadratic expression (equation 3) indicates the relative contributions of individual factors and their quadratic products to the percentage removal Cobalt(II) (Y).

$$Y = 70.26 - 1.56*A + 1.6*B + 6.05*C + 1.31*D + 0.59*AB + 0.027*AC + 0.86*AD - 0.12*BC - 0.40 * BD + 0.25*CD + 0.65*A^2 + 0.46*B^2 - 0.29*C^2 - 2.42*D^2 \dots\dots\dots (3)$$

Y is the estimated response is in equation 3 and suggests that maximum percentage of biosorption of Cobalt(II) is 79.519 % for the set of Cobalt(II)

concentration(Co) = 15.126 mg/L, pH = 6.0, sorbent dosage (w) = 20.0 g/L, and absolute temperature(T) = 301.705K.

Table 1
Results from CCD for Cobalt(II) biosorption by E. globulus L.

Run	A(Co)mg/L	B(pH)	C(w), g/L	D(T), K	Biosorption of Cobalt(II) % Removal	Predicted
1	0.000	0.000	0.000	0.000	70.2648	70.2648
2	1.000	-1.000	1.000	-1.000	68.5800	68.3089
3	0.000	0.000	0.000	-2.000	57.2800	57.9842
4	1.000	1.000	-1.000	1.000	65.3000	64.8751
5	-1.000	-1.000	-1.000	-1.000	63.1000	62.4838
6	-1.000	-1.000	-1.000	1.000	63.5800	63.6681
7	-1.000	1.000	1.000	1.000	77.8900	77.4460
8	-1.000	-1.000	1.000	-1.000	74.2290	74.2671
9	0.000	2.000	0.000	0.000	74.3650	75.2950
10	-1.000	-1.000	1.000	1.000	76.8000	76.4453
11	1.000	-1.000	1.000	1.000	73.8100	73.9362
12	0.000	0.000	0.000	0.000	70.2648	70.2648
13	1.000	1.000	1.000	1.000	77.3050	77.2902
14	-1.000	1.000	-1.000	-1.000	66.0500	65.5370
15	1.000	1.000	-1.000	-1.000	62.1000	61.8237
16	1.000	1.000	1.000	-1.000	73.7200	73.2451
17	0.000	0.000	0.000	2.000	62.9000	63.2137
18	0.000	0.000	0.000	0.000	70.2648	70.2648
19	0.000	0.000	-2.000	0.000	56.3800	57.0242
20	1.000	-1.000	-1.000	1.000	61.3120	61.0509
21	2.000	0.000	0.000	0.000	69.5100	69.7710
22	1.000	-1.000	-1.000	-1.000	56.3600	56.4172
23	0.000	0.000	0.000	0.000	70.2648	70.2648
24	-1.000	1.000	1.000	-1.000	77.2200	76.8501
25	0.000	0.000	0.000	0.000	70.2648	70.2648
26	-2.000	0.000	0.000	0.000	75.2364	70.2648
27	0.000	0.000	2.000	0.000	80.8490	81.2227
28	-1.000	1.000	-1.000	1.000	65.4990	65.1391
29	0.000	0.000	0.000	0.000	70.2648	70.2648
30	0.000	-2.000	0.000	0.000	68.8000	68.8878

A-Initial concentration(C_o) mg/L, B -pH of the solution, C- Biosorbent dosage(w) g/L, D(T) K- Temperature in kelvins CCD-Central Composite Design

Table 2
Levels of different process variables in coded and uncoded form for biosorption of Cobalt(II) using E.globulus L. leaf powder.

Variable	Name	Range and level				
		-2	-1	0	1	2
A	Initial concentration(C _o) mg/L	10	15	20	25	30
B	pH of the solution	3	4	5	6	7
C	Biosorbent dosage(w), g/L	5	10	15	20	25
D	Temperature(T), K	283	293	303	313	323

Table 3
ANOVA and estimated regression coefficients for the Cobalt(II) biosorption onto E.globulus L.

Source	Sum of Squares	df	Mean Square	F-Value	p-value	
					Prob > F	
Model	1261.92	14	90.14	297.31	< 0.0001	Significant
A	58.08	1	58.08	191.56	< 0.0001	
B	61.58	1	61.58	203.11	< 0.0001	
C	878.35	1	878.35	2897.18	< 0.0001	
D	41.02	1	41.02	135.31	< 0.0001	
A*B	5.54	1	5.54	18.27	0.00070	
A*C	0.012	1	0.012	0.039	0.84680*	
A*D	11.90	1	11.90	39.24	< 0.0001	
B*C	0.22	1	0.22	0.73	0.40650*	
B*D	2.50	1	2.50	8.26	0.01160	

C*D	0.99	1	0.99	3.26	0.09120*
A ²	11.74	1	11.74	38.74	< 0.0001
B ²	5.72	1	5.72	18.87	0.00060
C ²	2.23	1	2.23	7.37	0.01600
D ²	160.16	1	160.16	528.29	< 0.0001
Residual	4.55	15	0.30		
Lack of Fit	4.55	10	0.45		
Pure Error	0.000	5	0.000		
Cor Total	1266.47	29			

R^2 (Adj) = 0.9931 and R^2 (Pred) = 0.9793, *insignificant ANOVA-Analysis of variance

Model F-value of 297.31 implies that the model is significant. There is only a 0.01% chance that an F-value this large could occur due to noise. Values of "Prob> F" and < 0.05 indicates the model terms are

significant. In this case A, B, C, D, A*B, A*D, B*D, A², B², C², D² are significant model terms and A*C, B*C and C*D are not significant.




Figure 1

A comparison of predicted values and the percentage removal (actual values) and the normal plot for residual for Cobalt(II) onto *E.globulus L*




Figure 2

Surface contour plots, (a). pH -Initial conc -% Removal, (b). Initial conc- dosage - % Removal, (c). Temp.- Initial conc -% Removal, (d). pH - dosage -% Removal, (e). pH - Temp. - % Removal, (f). Temp. - dosage- % Removal.

Equilibrium Isotherms

Equilibrium isotherm is used to know the interactions between adsorption capacity and equilibrium concentration for metal removal. The equilibrium isotherms of Langmuir, Freundlich and Temkin models are adopted. Langmuir isotherm²⁷ model relates to coverage of molecules on a solid surface to concentration of a medium above the solid surface at a fixed temperature and is expressed by the following equation 4 and 5.

$$\frac{q_e}{q_{max}} = \frac{bC_e}{1+bC_e} \tag{4}$$

And the linear form of the above equation is

$$\frac{C_e}{q_e} = \frac{1}{bq_{max}} + \frac{C_e}{q_{max}} \tag{5}$$

A plot of (c_e / q_e) against (c_e) for *E. globulus* L. obtained for the experimental data indicates a slope and intercept. Where q_e is the amount of metal adsorbed at equilibrium (mg/g), q_{max} is the maximum adsorption capacity (mg/g). The separation factor, R_L of 0.93, which describes the essential characteristics of the Langmuir model and R_L can be defined in equation 5i and Langmuir isotherm for adsorption of Cobalt(II) onto *E. globulus* L. as shown in Fig.3.

$$R_L = \frac{1}{(1+bC_e)} \tag{5i}$$

$0 < R_L < 1$ indicates favorable adsorption, $R_L > 1$ indicates unfavorable adsorption, $R_L = 1$ indicates linear adsorption
 $R_L = 0$ indicates irrepressible adsorption




Figure 3
Langmuir isotherm for adsorption of Cobalt(II)

Freundlich isotherm²⁸ assumes that the adsorbent has heterogeneous surfaces on the sorbent and different functional groups are active in adsorption and is

$$q_e = K_f C_e^{1/n} \tag{6}$$

$$\ln q_e = \ln K_f + \frac{1}{n} \ln C_e \tag{7}$$

expressed by the following equation 6 and the linear form of above equation is in equation 7.

The value of slope, n (1.444) in the equation satisfies the condition of $0 < n < 1$ indicating suitability for sorption and Freundlich constant from slope and intercept of the

equation from Freundlich isotherm for biosorption of Cobalt(II) in




Figure 4
Freundlich isotherm for biosorption of Cobalt(II)

Temkin and Pyzhev isotherm²⁹ equation assumes low interactions between the adsorbent and adsorbate and the adsorption energy of all the molecules in the surface layer decrease at the cover surface and the linear form is in equation 8.

$$q_e = \frac{RT}{b_T} \ln A_T + \frac{RT}{b_T} \ln C_e \tag{8}$$

Temkin constants can be calculated from slope and intercepts for the plot of q_e against $\ln c_e$. Where R is universal gas constant (8.314 J/mol K) and T is the

absolute temperature (K), A_T is the Temkin isotherm constant (L/g), C_e is the maximum binding energy and b_T is the Temkin constant related to the heat of sorption (J/mol). Temkin isotherm for biosorption of Cobalt(II)

onto *E. globulus* L. as shown in Fig.5. All isotherm constants evaluated at optimum conditions are provided in Table 4.




Figure 5
Temkin isotherm for biosorption of Cobalt(II) onto *E. globulus* L.

Table 4
Equilibrium Isotherm constants

Langmuir	Freundlich	Temkin
$q_{max} = 7.194 \text{ mg/g}$	$K_f = 0.23 \text{ mg/g}$	$b_T = 2094.04$
$K_L = b = 0.0193 \text{ L/g}$	$n = 1.44$	$A_T = 0.3157$
$R^2 = 0.8499$	$R^2 = 0.987$	$R^2 = 0.914$

q_{max} - Maximum adsorption capacity (mg/g), K_L - b - Langmuir constant, R^2 - Regression coefficient, K_f and n are Freundlich coefficients, A_T - Temkin isotherm constant (L/g), b_T - Temkin constant related to the heat of sorption (J/mol)

Kinetics of biosorption

Lagergren pseudo first kinetic model and pseudo second order kinetic models are in equation 9 and 10, applied to the current study and the rate constants and equilibrium metal uptake capacity are evaluated from plot of $\log(q_e - q_t)$ against time (t) for pseudo first order and the intercept value should be equals the experimental q_e then the reaction can be categorized as pseudo first order rate equation¹¹ and from plot of t/q_t against t for pseudo second order model considers the rate-limiting step as the formation of chemisorptive bond

involving electron sharing between the solute and sorbent. The rate constants, correlation coefficients for biosorption of Cobalt(II) onto *E. globulus* are tabulated in Table 5.

$$\log(q_e - q_t) = \log q_e - \frac{k_1}{2.303} t \quad (9)$$

$$\frac{t}{q_t} = \frac{1}{k_2 q_e^2} + \frac{1}{q_e} t \quad (10)$$

Table 5
The rate equations and coefficients for biosorption of Cobalt(II) onto *E. globulus* L.

Kinetic model	Obtained equation	Rate constant,	R^2
Pseudo 1 st order	$\log(q_e - q_t) = -0.0111t - 0.4141$	$k_1 = 0.0253 \text{ min}^{-1}$	0.9582
2 nd order	$(t/q_t) = 0.771 t + 1.6387$	$k_2 = 0.363 \text{ g/mg.min}$	0.9974

Characterization of the *E. globulus* L. FT-IR analysis, SEM and EDX analysis

FTIR (ALPHA interferometer (ECO-ATR), Germany), Spectra are as shown in Fig.6(a1 -a2). Peaks at 3327 and 3249 cm^{-1} indicate strong O-H stretch, those at 2911 cm^{-1} C-H stretch, peak at 2354 cm^{-1} indicates alkyne triple bonds and nitrile triple bonds. Peaks at 1628 cm^{-1} C=C stretch, Peak at 1441 cm^{-1} -C-H bending, Wave number at 1317, 1221, 1153 cm^{-1} C-N stretch and 1046 cm^{-1} are due to the skeletal vibration of the C-O stretch, Peak at and 850-550 C-Cl stretching vibrations. The Spectra of adsorbent before and after

adsorption showed similar band characteristics, but the intensity of bands is either stronger or weaker. The peaks in Fig.6(a1) are shifted from 3327, 3249, 2911, 2354, 1628, 1441, 1317, 1221, 1153, 1046, 771, 662 and 582 cm^{-1} to 3338, 3244, 2906, 2348, 1631, 1531, 1441, 1324, 1224, 1018, 655 and 540 cm^{-1} in Fig.6(a2) respectively. The change in band strongly indicates that adsorption has happened and Fig.6 (b1- b2) shows SEM images and Fig.6(c1- c2) shows EDAX spectra of the *E. globulus* L. before and after Cobalt(II) loading.




Figure 6
(a1- a2). FTIR spectra, (b1- b2). SEM images and (c1- c2). EDAX spectra of the E.globulus L. before and after Cobalt(II) loading.

Comparative study

A comparative picture of various studies, on removal of Cobalt(II) by adsorption process, is provided in Table 6.

Table 6
Comparison of maximum adsorption capacity, based on the Langmuir isotherm by different adsorbents for Cobalt(II) from aqueous solutions.

Adsorbent	Concentration range, mg/L	Time, min	pH	Metal uptake, mg/g
Acacia nilotica ¹²	5-200	120	5.0	35.461
PET-TSE fibers ³⁰	10-400	180	5.0	78.08
Natural hemp fibers ³¹ (batch & fixed bed)	25-200	25	4.5	13.58& 15.44
Apricot stone activated carbon ³² (ASAC)	40-80	90	9.0	111.11
Chrysanthemum indicum flower raw(CIFR) and biochar(CIFBC) ¹⁰	25-200	60&45	5.0	14.84 45.44
Cystoseira indicia (C. indicia) ³³	25 and 300	4h	5.0	59.524
Ficus carica ³⁴ (MgCl ₂ treated)	20-200	120	6.0	82.64
Facial adsorbent ⁹	2-75.10	120	9.5	157.73
Diplotaxis harra & G. coronaria L ³⁵	25-200	60 & 45	6.5 & 7.5	33.02 & 24.52
E. globulus L.	5 -150	60	5.0	7.192

PET-TSE- Poly ethylene terephthalate thiosemicarbazide MgCl₂-Magnesium dichloride, CIFR- Chrysanthemum indicum flower raw, CIFBC- Chrysanthemum indicum flower biochar, G. coronaria L- Glebionis coronaria leaf.

CONCLUSION

The study conclusively establishes the suitability of *E. globulus L.* to remove Cobalt(II) after thorough investigation.

REFERENCES

1. Sud D, Mahajan G, Kaur MP. Agricultural waste material as potential adsorbent for sequestering heavy metal ions from aqueous solutions–A review. Bioresour Technol. 2008 Sep 30; 99(14):6017-27.

CONFLICT OF INTEREST

Conflict of interest declared none.

2. Dursun AY. A comparative study on determination of the equilibrium, kinetic and thermodynamic parameters of biosorption of copper (II) and lead (II) ions onto pretreated Aspergillus niger. Biochem Engg J. 2006 Feb 15; 28(2):187-95.
 3. Gray NF. Water technology: an introduction for environmental scientists and engineers. 2nd ed.

- Ireland: Elsevier Sci and Technol Books; 2005.p.551-52.
4. Shahalam AM, Al-Harthy A, Al-Zawhry A. Feed water pretreatment in RO systems: unit processes in the Middle East. *Desal.* 2002 Nov 10; 150(3):235-45.
 5. Kang SY, Lee JU, Moon SH, Kim KW. Competitive adsorption characteristics of Co²⁺, Ni²⁺, and Cr³⁺ by IRN-77 cation exchange resin in synthesized wastewater. *Chemosphere.* 2004 Jul 31; 56(2):141-7.
 6. El Samrani AG, Lartiges BS, Villiéras F. Chemical coagulation of combined sewer overflow: Heavy metal removal and treatment optimization. *Water research.* 2008 Feb 29; 42(4):951-60.
 7. Sadrzadeh M, Mohammadi T, Ivakpour J, Kasiri N. Neural network modeling of Pb²⁺ removal from wastewater using electrodialysis. *Chem Engg and Processing: Process Intensification.* 2009 Aug 31; 48(8):1371-81.
 8. Landaburu-Aguirre J, García V, Pongrácz E, Keiski RL. The removal of zinc from synthetic wastewaters by micellar-enhanced ultrafiltration: statistical design of experiments. *Desal.* 2009 May 15; 240(1):262-9.
 9. Shahat A, Awual MR, Naushad M. Functional ligand anchored nanomaterial based facial adsorbent for cobalt (II) detection and removal from water samples. *Chem Engg J.* 2015 Jul 1; 271:155-63.
 10. Vilvanathan S, Shanthakumar S. Biosorption of Co (II) ions from aqueous solution using *Chrysanthemum indicum*: Kinetics, equilibrium and thermodynamics. *Process Safety and Environ Protection.* 2015 Jul 31; 96:98-110.
 11. Thilagavathy P, Santhi T. Kinetics, isotherms and equilibrium study of Co (II) adsorption from single and binary aqueous solutions by *Acacia nilotica* leaf carbon. *Chinese J of Chem Engg.* 2014 Nov 30; 22(11):1193-8.
 12. Jimoh TO, Buoro AT, Muriana M. Utilization of *Blighia sapida* (Akee apple) pod in the removal of lead, cadmium and cobalt ions from aqueous solution. *J of Environ Chemi and Ecotoxi.* Vol. 2012 Jul; 4(10):178-87.
 13. Krishna B, Venkateswarlu P. Influence of *Ficus religiosa* leaf powder on bisorption of cobalt. *Indian J of Chem Technol.* 2011 Sep 1; 18(5):381-90.
 14. Bhatnagar A, Minocha AK, Sillanpää M. Adsorptive removal of cobalt from aqueous solution by utilizing lemon peel as biosorbent. *Biochem Engg J.* 2010 Jan 15;48(2):181-6.
 15. Bhatti HN, Bajwa II, Hanif MA, Bukhari IH. Removal of lead and cobalt using lignocellulosic fiber derived from *Citrus reticulata* waste biomass. *Korean J of Chem Engg.* 2010 Jan 1; 27(1):218-27.
 16. Li X, Tang Y, Cao X, Lu D, Luo F, Shao W. Preparation and evaluation of orange peel cellulose adsorbents for effective removal of cadmium, zinc, cobalt and nickel. *Colloids and Surfaces A: Physicochem and Engg Aspects.* 2008 Mar 20; 317(1):512-21.
 17. Pipíška M, Horník M, Vrtoch L, Augustín J, Lesný J. Biosorption of Zn and Co ions by *Evernia prunastri* from single and binary metal solutions. *Chemi and Ecol.* 2008 Jun 1;24(3):181-90.
 18. Javed MA, Bhatti HN, Hanif MA, Nadeem R. Kinetic and equilibrium modeling of Pb (II) and Co (II) sorption onto rose waste biomass. *Sep Sci and Technol.* 2007 Dec 1; 42(16):3641-56.
 19. Dahiya S, Tripathi RM, Hegde AG. Biosorption of lead and copper from aqueous solutions by pre-treated crab and arca shell biomass. *Bioresour Technol.* 2008 Jan 31; 99(1):179-87.
 20. Güzel F, Yakut H, Topal G. Determination of kinetic and equilibrium parameters of the batch adsorption of Mn (II), Co (II), Ni (II) and Cu (II) from aqueous solution by black carrot (*Daucus carota* L.) residues. *J of Hazard Mater.* 2008 May 30; 153(3):1275-87.
 21. Vijayaraghavan K, Palanivelu K, Velan M. Biosorption of copper (II) and cobalt (II) from aqueous solutions by crab shell particles. *Bioresour Technol.* 2006 Aug 31; 97(12):1411-9.
 22. Demirbaş E. Adsorption of cobalt (II) ions from aqueous solution onto activated carbon prepared from hazelnut shells. *Adsorp Sci & Technol.* 2003 Dec 1; 21(10):951-63.
 23. Annadurai G, Juang RS, Lee DJ. Adsorption of heavy metals from water using banana and orange peels. *Water Sci and Technol.* 2003 Jan 1; 47(1):185-90.
 24. Tyagi AK, Malik A. Antimicrobial potential and chemical composition of *Eucalyptus globulus* oil in liquid and vapour phase against food spoilage microorganisms. *Food Chemi* 2011 May 1; 126(1):228-35.
 25. Sebei K, Sakouhi F, Herchi W, Khouja ML, Boukhchina S. Chemical composition and antibacterial activities of seven *Eucalyptus* species essential oils leaves. *Biologi Research.* 2015 Jan 19; 48(1):1.
 26. Pereira V, Dias C, Vasconcelos MC, Rosa E, Saavedra MJ. Antibacterial activity and synergistic effects between *Eucalyptus globulus* leaf residues (essential oils and extracts) and antibiotics against several isolates of respiratory tract infections (*Pseudomonas aeruginosa*). *Indust Crops and Products.* 2014 Jan 31; 52:1-7.
 27. Langmuir I. The constitution and fundamental properties of solids and liquids. Part I. solids. *J of the Ameri Chemi Soc.* 1916 Nov;38(11):2221-95.
 28. Freundlich H. *Kapillarchemie, eine Darstellung der Chemie der Kolloide und verwandter Gebiete*, von Dr. Herbert Freundlich, akademische Verlagsgesellschaft; 1909.
 29. Boparai HK, Joseph M, O'Carroll DM. Kinetics and thermodynamics of cadmium ion removal by adsorption onto nano zerovalent iron particles. *J of Hazard Mater.* 2011 Feb 15; 186(1):458-65.
 30. Monier M, Abdel-Latif DA. Modification and characterization of PET fibers for fast removal of Hg (II), Cu (II) and Co (II) metal ions from aqueous solutions. *J of Hazard Mater.* 2013 Apr 15; 250:122-30.
 31. Tofan L, Teodosiu C, Paduraru C, Wenkert R. Cobalt (II) removal from aqueous solutions by

- natural hemp fibers: batch and fixed-bed column studies. *Applied Surf Sci.* 2013 Nov 15; 285:33-9.
32. Abbas M, Kaddour S, Trari M. Kinetic and equilibrium studies of cobalt adsorption on apricot stone activated carbon. *J. of Indust. and Eng. Chemi.* 2014 May 25; 20(3):745-51.
33. Akbari M, Hallajisani A, Keshtkar AR, Shahbeig H, Ghorbanian SA. Equilibrium and kinetic study and modeling of Cu (II) and Co (II) synergistic biosorption from Cu (II)-Co (II) single and binary mixtures on brown algae *C. indica*. *J of Environ Chem Engg.* 2015 Mar 31; 3(1):140-9.
34. Dabbagh R, Ashtiani Moghaddam Z, Ghafourian H. Removal of cobalt (II) ion from water by adsorption using intact and modified *Ficus carica* leaves as low-cost natural sorbent. *Desal and Water Treat.* 2015 Oct 20:1-3.
35. Tounsadi H, Khalidi A, Abdennouri M, Barka N. Biosorption potential of *Diplotaxis harra* and *Glebionis coronaria* L. biomasses for the removal of Cd (II) and Co (II) from aqueous solutions. *J of Environ Chem Engg.* 2015 Jun 30; 3(2):822-30.

Reviewers of this article

P. Rajendra Prasad

Professor, Department of Chemical
Engineering,
A.U. College of Engineering (A)
Andhra University, Visakhapatnam, India



Prof. Srawan Kumar G.Y

Associate Professor, Nalanda Institute of
Pharmaceutical Sciences, Sattenapalli,
Guntur, Andrapradesh, India



Prof. Dr. K. Suriaprabha

Asst. Editor, International Journal
of Pharma and Bio sciences.



Prof. P. Muthuprasanna

Managing Editor, International
Journal of Pharma and Bio sciences.

We sincerely thank the above reviewers for peer reviewing the manuscript