



QUANTITATIVE ANALYSIS OF FLAVONOID ‘NARINGIN’ A NATURAL PRODUCT AND ITS CORRELATION WITH ANTIOXIDANT ACTIVITY IN DIFFERENT FRUIT FRACTIONS OF GRAPEFRUIT, *CITRUS DECUMANA* VAR. *PARADISI* (MACFAD.) H.H.A NICHOLLS]: USE OF INDUSTRIAL WASTE

DR. BABITA RANA* AND DR. HARSHAD M. PANDIT

Botany Department, G.N. Khalsa College, Matunga, Mumbai-Maharashtra, India 400019.

ABSTRACT

Grapefruit [*Citrus decumana* var. *paradisi* (Macfad.) H.H.A Nicholls] is a subtropical citrus tree. It is an 18th century hybrid of *Citrus maxima* × *Citrus sinensis*, was first bred in Barbados (West Indies). Among the various phytochemicals detected in grapefruit, the flavonoid, naringin, which contributes to the distinctive bitterness of citrus fruits, exerts a variety of pharmacological effects. The methanolic extract of different parts of the fruit such as flavedo, albedo, segment membranes and seeds which are the main by products in citrus processing industry were subjected to HPLC analysis for their naringin content with mobile phase containing acetonitrile and 2% acetic acid (25:75) at wavelength 280 nm while antioxidant activity was determined by DPPH free radical scavenging assay. Highest naringin content was detected in albedo, flavedo and juice followed by segment membranes and seeds. Antioxidant activity determined by DPPH free radical scavenging activity revealed the fruit peel containing albedo and flavedo parts with highest antioxidant activity which could be correlated with their flavonoid content.

KEY WORDS: *Grapefruit, naringin, flavedo, albedo, acetonitrile, antioxidant, DPPH.*



DR. BABITA RANA*

Botany Department, G.N. Khalsa College, Matunga, Mumbai-Maharashtra, India 400019.

*Corresponding Author

Received on: 05-01-2017

Revised and Accepted on: 11-04-2017

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.2.b715-721>

INTRODUCTION

The beneficial effect of a diet supplemented with fruits and vegetables have enhanced interest in their bioactive compounds. It has been shown that positive effects of these natural products is usually linked with their antioxidant compounds such as carotenoids, flavonoids, vitamin C etc. Until the last 2 decades, it was assumed that vitamin C and carotenoids were the only chemopreventive agents in citrus because of its free radical scavenging ability.¹ Recent accumulative evidences suggest that citrus contains several possible anti-cancer agents such as flavonoids and limonoids.² Citrus flavonoids have shown a wide range of therapeutic properties for medical and clinical applications such as anti-inflammatory, antihypertensive, diuretic, analgesic, and hypolipidemic effects^{2,3}. Citrus juice flavonoids have also shown activity against acute myeloid leukemia.⁴ A flavonoid glycoside, naringin, which contributes to the distinctive bitterness of citrus fruits, was first found in grapefruit exerts a variety of pharmacological effects such as antioxidant, anticancer, anti-atherogenic and blood lipid lowering activity.⁵ To increase the amount of nutrients absorbed by the body the activity of several enzymes within the digestive tract is decreased by naringin. This prevents the breakdown of nutrients allowing more to be absorbed into the blood stream. Naringin supplements are consumed along with other nutritional supplements

to increase their absorption. Naringin, when treated with potassium hydroxide or another strong base, then catalytically hydrogenated, it becomes naringin dihydrochalcone, a compound roughly 300-1800 times sweeter than sugar at threshold concentrations^{5,6}. Solid residues consisting of grapefruit peel, segment membranes and seeds of grapefruit processing industry cause great environment havoc in term of its disposal. It was opportune if this residue can be exploited to produce higher value added products for use in food, pharmaceutical and cosmetic industry. Commercial brands of grapefruit juice have been analyzed for their flavonoid content by HPLC and naringin was identified in all grapefruit juices^{7,8}. In this study different parts of grapefruit fruit such as albedo, flavedo, segment membranes and seeds in addition to juice were subjected to HPLC analysis to determine their naringin content and antioxidant activity was determined by DPPH (2,2-diphenyl-1-picrylhydrazyl) free radical scavenging assay.

Naringin profile

Chemical name of naringin: Naringenine-7-rhamnoside-glucoside; 4',5,7-Trihydroxyflavone-7-rhamnoglucoside


Chemical formula of naringin: $C_{27}H_{32}O_{14}$

Molecular weight: 580.53 in its purest form.

Molar mass: 580.54 g/mol

Melting point: 166°C

Molecular structure of naringin



MATERIAL AND METHODS

Material

Fully mature fruits of grapefruit were obtained in the month of January from the plant growing in the college campus. The fruit was identified by the Head Department of Botany of our college (specimen #: br p 190111) and deposited in the college herbarium.

HPLC analysis of flavonoid naringin

Reagents and chemicals

Glacial acetic acid was purchased from S.D. Fine Chemicals, Mumbai and methanol, acetonitrile, deionised water from Merck Limited, Mumbai. All HPLC grade, standard naringin were obtained from Hi Media Laboratories Pvt. Limited, Mumbai.

Instrumentation

HPLC system consists of Shimadzu Prominence HPLC

(gradient) system with two hydraulic pumps, 6-valved Rheodyne injector, Thermoscientific Synchronis C18 column (250 × 4.6 mm, pore size 5µm) and a diode array detector. Data was processed by acquisition and integration LC software solutions.

Preparation of samples

Fruit was thoroughly washed with running tap water and divided into five different parts: flavedo, albedo, segment membranes, fruit juice and seeds as shown in Fig.1. All these tissue except for the juice were homogenously blended with the help of methanol by using a blender to make fine slurry. The juice was typically obtained by use of mechanical juice maker. Both juice and slurry of different tissues were stored frozen at -80°C for future analysis.




Figure 1
Cut fruit of grapefruit to show different parts used for naringin analysis

Extraction of naringin from fruit juice

A sample of 1-2 ml of the juice was extracted with 4 ml of methanol by shaking for 1 min using a Vortex Mixer and then followed by centrifuging at 3000 rpm for 10 min. The residue was re-extracted twice under the same condition to ensure complete extraction. The extract was passed through a 0.22 μm nylon filter prior to the HPLC analysis⁹.

Extraction of naringin from fruit parts

Sample of 5 gm slurry of each was weighed and then extracted with 20 ml of methanol by shaking for 1 min with a Vortex Mixer, followed by centrifuging at 3000 rpm for 10 min. The extract was passed through 0.22 μm nylon filter prior to the HPLC analysis. Albedo, flavedo and segment membrane samples were diluted with methanol to 1:50, 1:25 and 1:25 respectively. Seed and juice samples were used without any dilution¹⁰.

Standard naringin solution

100 mg / 100 mL (1 mg / mL). 100 mg or 0.1 gm standard naringin was dissolved in 100 ml of mobile phase consisting of 2% acetic acid: acetonitrile (75:30) to prepare 1000 ppm stock solution of naringin. Standard solutions of naringin were prepared by diluting the stock solution to 60, 80, 100, 120 and 160 ppm with the mobile phase.

Determination of naringin by HPLC

For determination of naringin mobile phase consisted of (2% acetic acid:acetonitrile, 75:30) with flow rate of 1 ml/min. The injection volume of sample and detection wavelength was 20 μl . and 280 nm respectively. Experiment was carried out at room temperature and pressure 134 kgf (Kilogram force).

Antioxidant activity by DPPH free radical scavenging assay

The radical scavenging ability of flavedo, albedo, segment membranes, seeds and juice was assessed on the basis of the radical scavenging effect on the DPPH free radical. The fruit juice and fruit fractions sample as prepared for naringin analysis were prepared in methanol (10-200 μg /ml). In clean and labeled test tubes, 2 mL of DPPH solution (0.002% in methanol) was mixed with 2 ml of each sample extract. The tubes were incubated at room temperature in dark for 30 minutes and the optical density was measured at 517 nm using

UV-Vis Spectrophotometer. The absorbance of the DPPH control was also noted. The scavenging activity of all fruit fractions and juice was calculated using the formula: Scavenging activity (%) = $[(A - B) / A] \times 100$, where A is absorbance of DPPH control and B is absorbance of DPPH and sample combination^{10, 11}. Experiment was repeated thrice for each sample and results were expressed as mean \pm SD.

Hydrogen peroxide scavenging activity

Hydrogen peroxide scavenging activity of various samples prepared from different fruit parts (albedo, flavedo, segment membranes, seeds and fruit juice) was determined according to the method of Yen and Chen (1995)¹¹. A solution of hydrogen peroxide (45 mM) was prepared in 0.1M phosphate buffer (pH 7.4). Different concentrations (10–200 $\mu\text{g}/\text{ml}$) of different samples prepared in methanol were added to 0.6 ml of hydrogen peroxide solution (45 mM). Absorbance of reaction mixture was recorded at 230 nm in a spectrophotometer (Shimadzu UV- 1201; Shimadzu, Kyoto, Japan) against a blank solution containing phosphate buffer without hydrogen peroxide. The percentage of scavenging of hydrogen peroxide of different samples was calculated using the following equation: Scavenging activity (%) = $[(A - B) / A] \times 100$, where A is absorbance of hydrogen peroxide control and B is absorbance of hydrogen peroxide and sample combination^{12, 13, 14}.

RESULTS AND DISCUSSION

HPLC analysis of flavonoid naringin

Each fruit part - flavedo, albedo, segment membranes, seeds and juice was analyzed for naringin content. Chromatogram of standard naringin (Fig. 2) showed the RT (retention time) equal to 6.646 min. and the peak areas detected at wavelength 280 nm of all standard concentrations of naringin were taken to construct a calibration curve¹². Calibration curve (Fig.7) obtained was linear with regression coefficient $R^2 = 0.9953$. By simply determining the peak area for naringin from the chromatograms of all samples (Fig.,3,4,5 and 6) a line was drawn on the standard naringin graph at that area. When the calibration curve is reached, a line was dropped down to the x-axis which gave the concentration of the naringin in sample (Fig.8)^{15, 16, 17}.




Figure 2
Chromatogram of standard naringin (100 ppm)




Figure 3
Chromatogram of naringin in grapefruit albedo




Figure 4
Chromatogram of naringin in grapefruit flavedo




Figure 5
Chromatogram of naringin in grapefruit juice




Figure 6
Chromatogram of naringin in grapefruit segment membranes




Figure 7
Calibration graph of naringin




Figure 8
Concentration of naringin in grapefruit juice shown with drop line

Flavonoid composition appears to vary greatly in the different parts of the fruit used. The major source of naringin was the albedo, while seed and segment membranes contained a minor amount. Fruit parts that contained naringin in a decreasing order were: albedo (3924.50 ± 385.43 ppm), flavedo (2575.50 ± 318.67 ppm), juice (1872.34 ± 259.33 ppm), seeds (1478.53 ± 93.78 ppm) and segment membranes (524.70 ± 87.90 ppm).

Antioxidant activity by DPPH free radical scavenging assay

The antioxidant activity increases with increase in concentration of sample solution. The highest antioxidant activity in term of % inhibition (Table 1) was

detected in juice (98.85 ± 1.78), followed by flavedo (79.87 ± 0.24), albedo (69.80 ± 0.45), segment membranes (39.55 ± 1.15) and seeds (35.42 ± 1.25)¹⁸.

Antioxidant activity by hydrogen peroxide scavenging assay

Antioxidant activity of methanolic fruit fraction samples is also dose dependent as highest antioxidant activity was shown at concentration 200 μ g/ml for all samples. The same results were obtained by hydrogen scavenging assay although the overall percentage inhibition is less as compared to DPPH scavenging assay but again highest antioxidant activity is detected in fruit peel (albedo 59.89 ± 0.75 , flavedo 67.31 ± 0.65) next to fruit juice (89.90 ± 0.90) as in Table 2.

Table 1
DPPH free radical scavenging assay of different fruit fractions of *Citrus paradisi*

Concentration µg/ml	% inhibition of albedo	% inhibition of flavedo	% inhibition of segment membranes	% inhibition of seeds	% inhibition of fruit juice
10	23.31±0.23	30.23 ±0.42	7.25±0.21	5.66±0.19	57.72±0.33
20	25.45±1.2	32.57 ± 0.35	8.12±0.23	6.65±0.20	60.50±0.45
30	28.18±1.34	39.65 ±0.26	12.87±0.35	8.43±0.32	68.34±0.35
40	35.76±0.89	45.67 ± 0.41	15.48±0.67	11.30±0.12	76.71±1.62
50	46.32±0.34	56.12 ±0.32	19.40±0.78	15.40±0.15	80.15±1.32
100	54.15±0.32	64.67 ± 0.13	24.12±0.43	20.42±0.89	92.50±1.85
150	63.78±1.45	72.73 ±0.32	28.16±1.30	25.86±0.56	97.72±2.30
200	69.80±0.45	79.87 ±0.24	39.55±1.15	35.42±1.25	98.85±1.78

Table 2
Hydrogen peroxide scavenging activity of different fruit fractions of *Citrus paradisi*

Concentration µg/ml	% inhibition of albedo	% inhibition of flavedo	% inhibition of segment membranes	% inhibition of seeds	% inhibition of fruit juice
10	20.21±0.29	24.35±0.40	6.21±0.12	4.21±0.13	43.67±0.67
20	23.37±0.34	27.21±0.21	7.19±0.15	4.79±0.14	50.40±0.29
30	25.17±0.31	31.75±0.32	10.15±0.13	5.80±0.09	55.32±0.90
40	30.32±0.54	39.45± 0.29	12.34±0.42	8.40±0.12	61.78±0.47
50	39.10±0.89	44.15±0.90	15.43±0.48	10.42±0.12	69.50±1.39
100	45.12±1.38	57.50±1.49	17.23±0.31	14.12±0.36	77.70±1.35
150	54.76±1.95	61.45±1.29	22.15±0.21	19.40±0.48	82.56±1.85
200	59.89±0.75	67.31±0.65	26.30±1.15	28.15±0.25	89.90±0.90

The results indicate that there is no direct correlation between flavonoid naringin content of grapefruit fruit fractions and antioxidant activity except the naringin content and antioxidant activity of peel (albedo and flavedo) and segment membranes. Thus many other phytochemicals such as carotenoids, phenolics and vitamin C reported in grapefruit may also playing a pivotal role in scavenging free radicals^{19,20}. Study conducted by AK Hadi Al- Anbari et al, on antioxidant activity of leaves and seeds of *Citrus paradisi* Macfad showed scavenging of Hydrogen peroxide 95.1%²¹. The high phenol and flavonoids contents of peels and tissue of *C. reticulata* var. Ponkan has been attributed to high antioxidant activity of this plant²². Toh JJ et al. (2013) reported that the peels of pommelo, *Citrus grandis* (L) Osbeck] fruit had higher antioxidant content and was positively high correlated with total phenolic content ($r = 0.978$) and total flavonoid content ($r = 0.959$)²³. The model of scavenging the stable DPPH radical is a widely used method to evaluate the free radical scavenging ability of various samples (Lee et al., 2003)²⁴. It was found that the radical- scavenging activities of all the extracts increased with increasing concentration. Phenols and polyphenolic compounds, such as flavonoids, are widely found in food products derived from plant sources, and they have been shown to possess significant antioxidant activities (Van Acker et al., 1996)²⁵. In the present study, the correlation between total flavonoids contents and radical scavenging activity of grapefruit fruit fractions were analyzed. In general, extracts or fractions with a high radical scavenging activity showed a high flavonoid content as well, but good correlations could not be found among them. Lack of relationship is in agreement with

other literature (Anagnostopoulou et al., 2006; Nickavar et al., 2007)^{26,27}. It is known that only flavonoids with a certain structure and particularly hydroxyl position in the molecule can act as proton donating and show radical scavenging activity (Mensor et al., 2001; Hou et al., 2003)^{28,29}. Furthermore, the extracts are very complex mixtures of many different compounds with distinct activities (Mensor et al., 2001; Hou et al., 2003).

CONCLUSION

The higher amount of flavonoid naringin in grapefruit peel, flavedo, albedo and segment membranes makes these fruit parts a valuable secondary raw material. The higher antioxidant activity of fruit peel and segment membranes next to fruit juice offers insight into how to explore further the benefits of by- products of citrus juice industry for use as a rich source of flavonoids which are the compounds of considerable importance to the food, pharmaceuticals and cosmetic industry.

ACKNOWLEDGEMENT

Authors are thankful to Head of Department of Guru Nanak Institute of Research and Development for sparing necessary facilities and guidance required carrying out HPLC analysis.

CONFLICT OF INTEREST

Conflict of interest declared none.

REFERENCES

1. Sies H and Stahl W. Vitamins E and C, beta-carotene, and other carotenoids as antioxidants. *Am. J. Clin. Nutr.* 1995 Dec; 62 (1):1315S–1321S.

2. Crespo ME, Galvez J, Cruz T, Ocete MA and Zarzuelo A. Anti-inflammatory activity of diosmin and hesperidin in rat colitis induced by TNBS. *Planta Medica* 1999 Oct; 65:651–653.

3. Gil-Izquierdo A, Gil MI, Ferreres F, Tomas-Barberan FA. *In vitro* availability of flavonoids and other phenolics in orange juice. *J. Agric. Food Chem* . 2001 Feb 19; 49 (2):1035-1041.
4. Tomasik P. Chemical and functional properties of food saccharides. Boca Raton London, CRC Press LLC. 2004.
5. G.P. Phondke. The Wealth of India-An Encyclopedia of India's Raw Materials Resources Ca-Ci. New Delhi: Publication and Information Directorate, CSIR, New Delhi; 1992.
6. Benavente-Garcia O, Castello J, Marin FR, Ortuno A and Del Rio JA. Use and properties of Citrus Flavonoids. In: *J. Agric. Food Chem*. 1997; 45:4506-4515.
7. FASEB (Federation of American Society for Experimental Biology) Evaluation of the health aspects of hesperidin, naringin and citrus bioflavonoids extracts as food ingredients. Life Sciences Research Office. FASEB, Bethesda, MD. 1982; 1-34.
8. Middleton E Jr., Kandaswami C. The impact of plant flavonoids on mammalian biology: Implications for immunity, inflammation and cancer. in: J.B. Harborne (Ed.) *The Flavonoids: Advances in Research Since 1986*. London UK: Chapman and Hall; 1993. p.619–652.
9. Harborne JB. *Phytochemical Methods; A guide to modern techniques of plant Analysis*. 2nd ed. New York: Chapman and Hall; 1973; 33-73.
10. Micali G, Coppolino R, Lo Coco F and Lanuzza F. Flavonoids in the pulp obtained from refining citrus fruit juices. *Forum ware Int*. 2003; 2: 54-61.
11. Gorinstein S, Leontowicz H, Leontowicz M Krzemiński R, Gralak M, Delgado-Licon E, Martinez Ayala AL, Katrich E and Trakhtenberg S. Changes in plasma lipid and antioxidant activity in rats as a result of naringin and red grapefruit supplementation. *J. Agric. Food Chem*. 2005 Apr 20; 53(8): 3223-8.
12. Kumar A, Dogra S, Prakash A. Productive effects of naringin, a citrus flavonoid against cochicine-induced cognitive dysfunction and oxidative damage in rats. *J. Med. Food*. 2010Aug; 13(4): 976-984.
13. Belajova E and Suhaj M. Determination of phenolic constituents in citrus juices: Method of high performance liquid chromatography. *Food Chem*. 2004 Jul; 86(3): 339-343.
14. Bronner WE and Beecher GR. Extraction and measurement of prominent flavonoids in orange and grapefruit juice concentrates. *J. Chromatogr. A*. 1995; 705: 247– 256.
15. Gorinstein S, Huang D, and Leontowicz H. Determination of naringin and hesperidin in citrus fruit by high performance liquid chromatography. Antioxidant potential of citrus fruit. *ACTA CHROMATOGR*. 2006; 17:108-124.
16. Jairam Vanamala, Lavanya Reddivari, Kil Sun Yoo, Leonard M. Pike, Bhimanagouda S. Patil. Variation in the content of bioactive flavonoids in different brands of orange and grapefruit juices. *J. Food Comp. Anal*. 2006; 19(2-3): 157-166.
17. Rouseff, RL. HPLC determination of Naringin and Neohesperidin to Detect Grapefruit Juice in Orange Juice. *J. Assoc. Off. Anal. Chem*. 1988 Jul- Aug; 71(4): 798-802.
18. Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity Songklanakarinn J. *Sci. Technol*. 2004; 26(2): 211-219.
19. Kekuda PTR, Shobha KS and Onkarappa R. Studies on antioxidant and anthelmintic activity of two *Streptomyces* species isolated from Western Ghat soil of Agumbe, Karnataka. *J Pharm Res*. 2010; 3(1): 26-29.
20. Yen GC and Chen HY. Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J. Agric. Food Chem*. 1995 Jan; 43(1): 27–32.
21. Hadi Al- Anbari AK and Hasan MA. Antioxidant Activity in Some Citrus Leaves and Seeds Ethanolic Extracts. *International Conference on Advances in Agricultural, Biological and Environmental Sciences*. 2015 Jul; 93-97.
22. Ghasemi K, Ghasemi K and Mohammad Ali E. Antioxidant activity, Phenol and Flavonoid contents of 13 citrus species peels and tissues. *Pak. J. Pharm. Sci*. 2009 Jul; 22(3): 277-281.
23. Toh JJ, Khoo HE and Azrina A. Comparison of antioxidant properties of pomelo [*Citrus grandis* (L) Osbeck] varieties. *Int. Food Res. J*. 2013 Oct; 20(4): 1661-1668.
24. Lee SE, Hwang HJ, Ha JS, Jeong HS and Kim JH. Screening of medicinal plant extracts for antioxidant activity. *Life Sci*, 2003 May; 73(2): 167-179.
25. Van Acker SA, Van Den Berg DJ, Tromp MN, Griffioen DH, Van Bennekom WP, Van der Vijgh WJ and Bast A. Structural aspects of antioxidant activity of flavanoids. *Free Radic. Biol. Med*. 1996; 20(3): 331-342.
26. Anagnostopoulou MA, Kefalas P, Papageorgiou VP, Assimopoulou AN and Boskou D. Radical scavenging activity of various extracts and fractions of sweet orange peel (*Citrus sinensis*). *Food Chem*. 2006; 94: 19-25.
27. Nickavar B, Kamalinejad M and Izadpanah H. *In vitro* free radical scavenging activity of five *Salvia* species. *Pak. J. Pharm. Sci*. 2007 Nov; 20(4):291-294.
28. Mensor LL, Menezes FS, Leitao GG, Reis AS, dos Santos TC, Coube CS and Leitao SG. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. *Phytother. Res*. 2001 Mar; 15: 127-130.
29. Hou WC, Lin RD, Cheng KT, Hung YT, Cho CH, Chen CH, Hwang SY and Lee MH. Free radical scavenging activity of Taiwanese native plants. *Phytomedicine*. 2003 Mar; 10(2-3): 170-175.

Reviewers of this article

Dr. Ganesh Iyer, M.Sc., Ph.D

Head, Department of Life Science,
Ramnarain Ruia College, L. Nappo Road,
Matunga, Dadar East, Mumbai,
Maharashtra 400019, India



Prof. Swaran Kumar G.Y

Associate Professor, Nalanda Institute of
Pharmaceutical Sciences, Sattenapalli,
Guntur, Andrapradesh, India



Prof. Dr. K. Suriaprabha

Asst. Editor, International Journal
of Pharma and Bio sciences.



Prof. P. Muthuprasanna

Managing Editor, International
Journal of Pharma and Bio sciences.

We sincerely thank the above reviewers for peer reviewing the manuscript