



NEW VALIDATED STABILITY INDICATING GRADIENT RP-HPLC METHOD FOR THE ASSAY & RELATED SUBSTANCES OF MICONAZOLE NITRATE AND ASSAY OF BENZOIC ACID IN MICONAZOLE NITRATE 2.0% W/W GEL

S.KUMARAVEL^{1*} AND P.SHANMUGAPANDIYAN²

¹Research Scholar, Prist University, Tanjore, Tamilnadu, India

²Department of Pharmaceutical analysis, Rao College of Pharmacy, Nellore, Andhra Pradesh, India

ABSTRACT

An innovative, simple, economic, selective and sensitive of the stability indicating gradient reverse phase high performance liquid chromatography (RP-HPLC) method has been developed and validated for the analysis of assay of Miconazole nitrate and Benzoic acid and related substances of Miconazole nitrate in gel dosage form with better precision and accuracy. Miconazole nitrate & Benzoic acid are found to be degraded together under different set of conditions as followed according to ICH guidelines and the degradants so formed along with Miconazole nitrate organic impurities were separated by using INERTSIL ODS C18(150mm x 4.6, 3.5 μ) column in gradient mode with mobile phase A consisting of 0.1% v/v solution of phosphoric acid in water and mobile phase B consisting of water : acetonitrile (22:78) with a flow rate of 1.0 mL/min and throughout the gradient program with an UV detection wavelength of 227nm for both the compounds with an injection volume of 10 μ L. Its chromatographic run time was 15.0 min. The method was validated according to ICH guidelines with respect to all validation parameters such as selectivity, specificity, linearity, precision, accuracy, robustness and limits of detection and quantification. All parameters examined were found to be well within the stated guidelines. The results were indicating the method is selective and specific in analysis of both Miconazole nitrate & Benzoic acid in the presence of degradation products formed under various stress conditions. The Proposed Gradient RP-HPLC method is utilized for the simultaneous estimation of Assay of Miconazole nitrate and Benzoic acid and related substance determination in Miconazole Nitrate Gel.

KEYWORDS: Miconazole nitrate, Benzoic acid, Gradient RP-HPLC, INERTSIL ODS C18, Orthophosphoric acid, Acetonitrile, ICH Guidelines.



S. KUMARAVEL*

Research Scholar, Prist University, Tanjore, Tamilnadu, India

Received on: 09-03-2017

Revised and Accepted on: 15-04-2017

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.2.p308-312>

INTRODUCTION

Miconazole nitrate is 2-(2,4-Dichlorophenyl)-2-(2,4-dichlorobenzoyloxy)-1-(1*H*-imidazol-1-yl)-ethane and usually available in different forms such as a 2% cream, lotion, or powder in the treatment of fungal infections of skin including candidiasis, dermatophytosis, and pityriasis versicolor¹⁻⁵. Literature review reveals that the methods for Miconazole nitrate alone or in combined dosage forms are Massaccesi reported the development of a two-phase titration method for the analysis of miconazole and other imidazole derivatives in pure form and in pharmaceutical formulation⁶. Szabolcs determined the active principle in preparations based on miconazole and clotrimazole⁷. The two drugs were determined in ointments by extraction with chloroform, evaporation of the solvent, dissolution of the residue in acetic acid, and titration with 0.1 N perchloric acid in the presence of Gentian Violet. Mapi et al developed a potentiometric method for the determination of the stability constants of miconazole complexes with iron(II), iron(III), cobalt(II), nickel(II), copper(II) and zinc(II) ions⁸. Shamsipur and Jalali described a simple and accurate pH metric method for the determination of two sparingly soluble in water antifungal agents; miconazole and ketoconazole in Micellar medium⁹. Bonazzi et al reported the determination of miconazole and other imidazole antimycotics in creams by supercritical-fluid extraction and derivative ultraviolet spectroscopic method¹⁰. Goger developed a quantitative method for the determination of miconazole in cream formulations that benzoic acid as preservative by second order derivative spectrophotometry¹¹. Erk described a spectrophotometric method for the simultaneous determination of metronidazole and miconazole nitrate in ovules¹². El-Shabouri et al used a charge-transfer complexation method for the spectrophotometric assay of miconazole nitrate and other imidazole antifungal drugs¹³. Chen used a second-derivative spectrophotometric method for the determination of miconazole nitrate in Pikangshuang¹⁴. Erk N, and Altum ML used a ratio spectra derivative spectrophotometric method and a high performance liquid chromatographic method for the analysis of miconazole nitrate and metronidazole in ovules¹⁵. Neill et al described an automated screening procedure using gas chromatography-mass spectrometry for identification miconazole and other drugs after their extraction from

biological sample¹⁶. Aboul-Enein and Ali used a high performance liquid chromatography method by using chiral columns for the analysis of miconazole and other imidazole antifungal agents in commercial dosage forms¹⁷⁻¹⁹. Zhang and Nunes used a high performance liquid chromatographic-coupled with electrospray ionization mass spectrometry (HPLC-ESI-MS) method to study the structure and generation mechanism of a novel degradation product formed by oxidatively induced coupling of miconazole nitrate with butylated hydroxytoluene in a topical ointment²⁰.

MATERIALS AND METHODS

The reference standard was supplied by M/s Microlabs limited, Bangalore, India. Acetonitrile (HPLC grade), methanol (HPLC grade), Orthophosphoric acid (HPLC grade), calcium chloride (AR grade) were purchased from Merck (Mumbai, India). Milli-Q water was used as HPLC grade water. Chromatographic Conditions: The determination was carried out on Waters HPLC equipped with Photo diode array as detector using data handling system – waters empower software. The column used in the development for the determination is INERTSIL ODS C18 (150 x 4.6, 3.5 μ). The mobile phase composition was used as per gradient Program indicated in Table 1. The detector wavelength was set at 227nm. A flow rate of 1.0mL/min was used for the determination. The samples and standards were dissolved in diluent (water: ACN, 30: 70) and 10 μ L sample were injected into HPLC system at the column at a temperature of 35°C.

Mobile phase


1.0 mL of orthophosphoric acid was mixed with 1000mL of milli Q water used as a mobile phase A. Water: Acetonitrile mixture (22:78) used as a mobile phase B. The mobile phases were filtered through 0.45 μ m Membrane filter and degassed by sonication.

RESULTS AND DISCUSSION

All the system suitability parameters according to ICH guidelines was achieved by using INERTSIL ODS C18 (150 x 4.6, 3.5 μ) column with a flow rate of 1.0 mL/min and an UV detection wavelength of 227nm with a injection volume of 10 μ L.

Table 1
Gradient Program

Time(minutes)	Flow Rate (ml/min)	%Mobile Phase A	%Mobile Phase B
0	1.0	80	20
2	1.0	80	20
8	1.0	53	47
21	1.0	50	50
25	1.0	20	80
38	1.0	20	80
40	1.0	80	20
46	1.0	80	20



Peak Results

	Name	RT	Area	% Area	Purity1 Angle	Purity1 Threshold	Purity1 Flag	Int Type
1	RO14821	4.964	35951	0.48	0.517	3.635	No	BB
2	Benzoic Acid	9.449	1180084	15.61	0.199	0.490	No	BB
3	RO66921	11.734	39030	0.52	0.608	2.250	No	BV
4	RO78890	12.928	43079	0.57	0.338	3.764	No	BB
5	Miconazole Nitrate	13.549	6221957	82.32	0.509	1.309	No	BB
6	RO60524	34.326	38299	0.51	0.997	5.069	No	BB

Figure 1
Spiked sample chromatogram

To validate the RP-HPLC method (Figure 1). The detection limit determined for the API and each validation impurity. Detection limit value will be set at 0.05% w/w. The quantitation limit (LOQ) determined for the API and each validation impurity. Quantitation limit value is set equal to reporting threshold value (0.10% w/w) and shall be supported by the accuracy at reporting threshold level. Limit of Detection (LOD) is

confirmed by preparation of solution containing Miconazole Nitrate, IMP-A, IMP-B and IMP-C injected one injection of this solution in the proposed methodology. Linearity solutions prepared from LOQ to 200% of IMP-A, IMP-B and IMP-C of the specification limit concentration and for Miconazole Nitrate from LOQ to 120% of test concentration and the correlation coefficient was found to be 0.9999.

Table 2
Limit of Quantitation

S.No	Component	LOQ	
		Conc.(µg/mL)	Conc. (% w/w)
1	Miconazole	0.202	0.081
2	Miconazole IMP-A	0.234	0.094
3	Miconazole IMP-B	0.214	0.086
4	Miconazole IMP-C	0.238	0.095

Table 3
Linearity Studies

Sample	Miconazole Nitrate		Benzoic acid	
	Concentration (µg/mL)	Area	Concentration (µg/mL)	Area
80.0 %	199.4320	5305644	17.6323	863637
90.0 %	224.3610	5980038	21.4106	1051773
100.0 %	249.2900	6624970	25.1889	1234269
110.0 %	274.2190	7252894	28.9673	1415689
120.0 %	299.1480	7943777	32.7456	1601912
Correlation coefficient (r)	0.999992		0.999981	
Slope	26546.0858		48711.0962	
Intercept	2875.8539		6475.1201	
Residual sum of squares	1570093447.6066		12660869.5301	

Table 4
Precision studies

Parameters Sample No.	Method Precision					Intermediate Precision				
	Assay		Purity (% w/w)			Assay		Purity (% w/w)		
	Benzoic acid	Miconazole Nitrate	IMP-A	IMP-B	IMP-C	Benzoic acid	Miconazole Nitrate	IMP-A	IMP-B	IMP-C
1	99.3	99.1	0.517	0.524	0.491	101.5	99.5	0.504	0.527	0.510
2	98.9	98.6	0.522	0.530	0.493	101.7	99.4	0.513	0.527	0.515
3	99.1	98.7	0.516	0.521	0.491	102.0	99.9	0.504	0.527	0.491
4	99.2	98.9	0.528	0.538	0.503	101.1	98.8	0.506	0.523	0.503
5	98.1	97.8	0.522	0.522	0.491	101.6	99.4	0.504	0.523	0.491
6	98.6	98.3	0.535	0.540	0.501	101.4	99.5	0.505	0.512	0.501
Mean	98.9	98.6	0.523	0.529	0.495	101.6	99.4	0.506	0.523	0.495
%RSD	0.5	0.5	1.3	1.5	1.2	0.3	0.4	0.8	1.1	0.4

The precision studies (expressed as the relative standard deviation (RSD)) were determined for miconazole nitrate and benzoic acid and the values are presented in Table 4.

Table 5
Recovery studies

Drug	Levels	% recovery	% RSD
Miconazole Nitrate	80%	99.0	0.2
	100%	99.1	0.2
	120%	98.9	0.3
Benzoic acid	70%	98.5	0.3
	100%	98.6	0.0
	130%	98.5	0.1
Miconazole Impurity-A	LOQ	96.0	1.3
	0.5%	97.4	0.5
	1.0%	96.9	1.0
Miconazole Impurity-B	LOQ	101.5	1.3
	0.5%	100.4	0.7
	1.0%	99.7	0.7
Miconazole Impurity-C	LOQ	94.0	0.6
	0.5%	97.3	0.6
	1.0%	97.4	0.7

The recovery experiment values in Table 5 obtained were performed by adding known amount of IMP-A, IMP-B and IMP-C impurities spiked into nine different flasks containing placebo with Miconazole nitrate drug substance and for Miconazole nitrate, drug substance stock solution spiked into nine different flasks containing placebo in three different levels, each in triplicate viz., LOQ level, 0.5% & 1.0% impurity level (Figure 2). Miconazole Nitrate at high concentration levels accuracy performed at 80.0%, 100.0% and 120.0%, w/w of sample concentration. Each concentration will be prepared 3 times by separate weighing and contains

placebo (without Active) at 100%, w/w level. Accuracy for Benzoic Acid performed at 70.0%, 100.0% and 130.0%, w/w levels. Each concentration prepared 3 times by separate weighing and contains placebo (without Benzoic Acid) at 100%, w/w level. The stability of sample was checked by forced degradation in different conditions and % of degradation was calculated. The peak purity of the analytes was passed in all conditions (purity angle should be less than the threshold value). The following values in Table 6 indicate that any other impurity is not merging with the main peak.

Table 6
Robustness studies

S.NO	Condition	% RSD	
		Miconazole Nitrate	Benzoic acid
1	Flow (+20%)	1.0	1.0
2	Flow (-20%)	1.0	1.0
3	Temperature (35°C)	0.2	0.2
4	Temperature (40°C)	0.0	0.1
5	Wavelength (+2nm)	0.2	0.2
6	Wavelength (-2nm)	0.1	0.2
7	Sonication time (+2%)	0.5	0.5
8	Sonication time (-2%)	0.5	0.5
9	Column lot-1	0.2	0.3
10	Column lot-2	0.3	0.3

The reliability of the method was determined by made small deliberate variations in method parameters and the RSD values (Table 7) obtained, an indication of its reliability on normal usage. Selectivity solution and reference solution was injected on different days during the validation study. Using the system suitability software, USP resolution between Imp- B and Miconazole Nitrate was calculated from selectivity

solution and USP tailing, USP plate count for Benzoic peak and Miconazole Nitrate were calculated from Reference solution-1. % RSD for five replicate injections for Miconazole nitrate peak and Benzoic acid peak were calculated from reference solution-1. % Recovery for Miconazole nitrate peak and Benzoic acid peak was calculated from reference solution-2 against reference solution-1

Table 7
System suitability studies

Experiment Name	Resolution	USP Tailing		USP Plate count		% RSD		% Recovery	
		Peak-1	Peak-2	Peak-1	Peak-2	Peak-1	Peak-2	Peak-1	Peak-2
System Suitability	2.3	1.1	1.7	50172	32938	0.2	0.1	101.0	100.6
Method Precision, Filter study & Robustness	2.2	1.1	1.7	45529	27239	0.3	0.3	101.0	100.4
Solution stability	1.7	1.1	1.5	52168	19448	0.2	0.1	101.0	99.9
Specificity	2.4	1.1	1.4	37500	31568	1.0	1.0	100.0	100.4
Linearity	2.3	1.0	1.5	45654	28981	0.1	0.1	99.4	99.3
Accuracy	2.4	1.0	1.4	44603	32699	0.1	0.2	98.7	99.5
Stress study & Robustness	2.3	1.0	1.4	36589	29110	0.2	0.3	100.1	100.1

CONCLUSION

The validated stability indicating HPLC method for Assay, Chromatographic Purity of Miconazole Nitrate and Assay of Benzoic Acid in Daktarin Gel (Miconazole Nitrate 2.0% w/w) is specific, linear, accurate, precise and robust. Analytical data of method precision study (performed by reference/sending laboratory, Lab-1) and intermediate precision study (performed by receiving

laboratory, Lab-2) were considered for the analytical method transfer activity. Hence, the analytical methodology for the assay of Miconazole nitrate, assay of Benzoic acid and degradation products of Miconazole nitrate in Daktarin gel is suitable for its intended use.

CONFLICT OF INTEREST

Conflict of interest declared none.

REFERENCES

- O'Neil MJ. The Merck Index: An Encyclopedia of Chemicals, Drugs, and Biologicals. 12th ed. New Jersey: Merck, 1996.p.894.
- Moffat AC. Clarke's Isolation and Identification of Drugs. 2nd ed. London. The Pharmaceutical Press. 1989.p.784.
- Sweetman SC. Martindale: The Complete Drug Reference, 32nd ed. London. The Pharmaceutical Press. 2002.p.384.
- British Pharmacopoeia Commission. *British Pharmacopoeia 1998*. London: TSO; 1998.
- United States Pharmaceutical Convention. *United States Pharmacopoeia 2003*. Rockville; 2003.p.1112.
- Massaccesi M. Two-phase Titration of some Imidazole derivatives in pharmaceutical preparations. *Analyst* 111: 1986; 987-989.
- Szabolcs L. Determination of the active principle in preparations based on Miconazole and clotrimazole. *Acta Pharm Hung*. 1976 Jan; 46(1):43-8.
- Mapsi N, De CJ, Ranter and Jottier W. Potentiometric determination of the stability constants of miconazole complexes with Fe(II), Fe(III), Co(II), Ni(II), Cu(II) and Zn(II) ions]. *J Pharm Belg*. 1987 Jan-Feb; 42(1):29-37.
- Shamsipur M and Jalali F. Acid-base titrimetric determination of ketoconazole and miconazole in micellar media. *Chem Anal*. 2002; 47(6):905-911.
- Bonazzi D, Cavrini V, Gatti R, Boselli E and Caboni M. Determination of imidazole antimycotics in creams by supercritical fluid extraction and derivative UV spectroscopy. *J Pharm Biomed Anal*. 1998 Oct; 18(1-2):235-40.
- Goger NG, Gokcen L. Quantitative Determination of Miconazole in Creams by Second Order Derivative Spectrophotometry. *Anal. Lett*. 1999; 32:2595-2602.
- Erk N. Simultaneous Determination of Metronidazole and Miconazole Nitrate in Ovules by Spectrophotometric Methods. *S.T.P. Pharma Sci*. 1996; 6:312-315.
- El-Shabouri SR, Emara KM, Khashaba PY, Mohamed AM. Charge-Transfer Complexation for Spectrophotometric Assay of Certain Imidazole Antifungal Drugs. *Anal. Lett*. 1998; 31:1367-1385.
- Chen B. Determination of miconazole nitrate in pikangshuang by second order derivative spectrophotometry. *Chinese Journal of Pharmaceuticals*. 1993; 7: Page 66-67.
- Erk N, and Altum ML. Spectrophotometric resolution of metronidazole and miconazole nitrate in ovules using ratio spectra derivative spectrophotometry and RP-LC. *J Pharm Biomed Anal*. 2001 Apr; 25(1):115-22.
- Neill GP, Davies NW, McLean S. Automated screening procedure using gas chromatography-mass spectrometry for identification of drugs after their extraction from biological samples. *J Chromatogr*. 1991 Apr 19; 565(1-2):207-24.
- Aboul-Enein HY and Ali I. Comparative study of the enantiomeric resolution of chiral antifungal drugs econazole, miconazole and sulconazole by HPLC on various cellulose chiral columns in normal phase mode. *J Pharm Biomed Anal*. 2002 Jan 15; 27(3-4):441-6.
- Ali I and Aboul-Enein HY. Enantioseparation of some clinically used drugs by HPLC using cellulose Tris (3,5-dichlorophenylcarbamate) chiral stationary phase. *Biomed Chromatogr*. 2003 Mar-Apr; 17(2-3):113-7
- Aboul-Enein HY and Ali I. Enantiomeric resolution of some imidazole antifungal agents on chiralpak WH chiral stationary phase using HPLC. *Chromatographia*. August 2001, Volume 54, Issue 3, pp 200-202.
- Zhang F and Nunes M. Structure and generation mechanism of a novel degradation product formed by oxidatively induced coupling of miconazole nitrate with butylated hydroxytoluene in a topical ointment studied by HPLC-ESI-MS and organic synthesis. *J Pharm Sci*. 2004 Feb; 93(2):300-9.

Reviewers of this article

Dr S Anbazhagan

Principal cum Prof, Dept of Pharmaceutical chemistry, Surya School of Pharmacy, NH-45, GST Road, Surya Nagar, Vikravandi .



G. Bakhya Shree M.S. (Research)

Affiliation

Coordinator and Trainer, Department of Biotechnology and Life Sciences, Dexter Academy, Madurai, Tamilnadu



Prof. Dr. K. Suriaprabha

Asst. Editor , International Journal of Pharma and Bio sciences.



Prof. P. Muthuprasanna

Managing Editor , International Journal of Pharma and Bio sciences.

We sincerely thank the above reviewers for peer reviewing the manuscript