



## ANTIOXIDANT AND PRELIMINARY PHYTOCHEMICAL STUDIES OF *ANETHUM GRAVEOLENS* LEAVES

**AMRITA KUMARI SRIVASTAVA, DEVESH KUMAR KUSHAWAHA AND GEETA WATAL\***

*Alternative Therapeutics Unit, Drug Development Division, Medicinal Research Laboratory,  
Department of Chemistry, University of Allahabad, Allahabad 211002, U. P., India.*

### ABSTRACT

Plants being rich in polyphenols are an important source of natural antioxidants which can neutralize free radicals generated in the system responsible for causing oxidative stress during diabetes. The present study deals with phytochemical screening and in vitro assessment of antioxidant potential of aqueous leaves extract of *Anethum graveolens*. The phytochemical analysis of the extract revealed the presence of tannins, saponins, carbohydrates, glycosides, alkaloids, flavonoids and proteins. The total phenolics and flavonoids contents present in the *A. graveolens* leaves extract were 66.5 mg of Gallic Acid Equivalent (GAE)/g and 29.8 mg of Quercetin Equivalent (QE) /g respectively. Significant antioxidant efficacy of *A. graveolens* leaves was further confirmed by percent inhibition of DPPH<sup>•</sup>, NO<sup>•</sup>, O<sup>2-</sup> radicals, OH radicals, ABTS<sup>+</sup> radicals and Metal Chelating efficacies as it was found to be 87.57, 68.29, 82.32, 74.02, 97.05 and 61.51% inhibition respectively. Therefore, the data suggest that the aqueous extract of *A. graveolens* leaves could be developed not only as an anti-aging agent but also as an agent for managing oxidative stress due to diabetic complications.

**KEYWORDS:** *Anethum graveolens* leaves, Phytochemicals, Antioxidant, Diabetic complications, Antiaging Agent, Polyphenols.



**GEETA WATAL \***

Alternative Therapeutics Unit, Drug Development Division, Medicinal Research Laboratory,  
Department of Chemistry, University of Allahabad, Allahabad 211002, U. P., India.

\*Corresponding author

Received on: 28.11.2016

Revised and Accepted on: 08.03.2017

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.2.p364-373>

## INTRODUCTION

Medicinal plants act as an indigenous source of phytochemicals possessing a therapeutic value which can be used in drug development. Phytochemicals as secondary metabolites are directly responsible for bioactivities of medicinal plants viz. antidiabetic, antioxidant, antimicrobial, anticancer and anti-inflammatory etc. Hence, bioactive secondary metabolites serve as important leads for the development of much cost-effective and less toxic pharmaceutical drugs and form the backbone of the modern system of medicine<sup>1,2</sup>. Thus, correlation between the phytochemicals and the bioactivity of the plant is desirable to know for developing the safe drugs with specific activities to treat various diseases<sup>3</sup>. Antioxidants are gaining much importance as anti-aging agents since they protect the human system from oxidative stress causing cell damage. Though, the production of free radicals in the human system helps in many cellular processes but sometimes due to over-production of these reactive oxygen species (ROS), resulting from an imbalance between their formation and neutralization causes oxidative stress. This failure of antioxidant defense system damages the cells and hence increases the risk of assaults of different diseases<sup>4</sup>. Both experimental and clinical studies suggest that oxidative stress induced by ROS also plays a major role in the pathogenesis of diabetes mellitus. Abnormally high levels of free radicals and the simultaneous decline of the antioxidant defense mechanisms can lead to damage of cellular organelles and enzymes, an increase of lipid peroxidation, and development of insulin resistance as well. These consequences of oxidative stress can promote the development of diabetic complications. Thus, the search for antioxidants from natural resources has paved the way for exploration of new compounds with better efficacy and lower toxicity. These are safe and easily accessible at lower cost in comparison to synthetic antioxidants. Natural antioxidants present in medicinal plants work mostly as free radical scavengers and therefore, are known to protect the system against ROS-mediated oxidative stress<sup>5</sup>. The use of various medicinal plants rich in antioxidant constituents has already been proposed by our research group as an effective therapeutic approach for controlling diabetes as well as diabetes-induced oxidative stress<sup>6,7</sup>. *Anethum graveolens* L. (Apiaceae), commonly known in English as 'dill' and in Hindi as 'Soya', is an annual herb found in Mediterranean region, Europe, and Central and South Asia. *A. graveolens* is used as an aromatic herb as well as medicine due to its high therapeutic value<sup>8</sup>. It is used in Iranian folk medicine as an anti-

hypercholesterolaemic plant<sup>9</sup>. It has traditionally been used for gastrointestinal ailments<sup>10</sup>. It is well known for its pharmacological effects, such as anti-microbial<sup>11-13</sup>, antioxidant<sup>14</sup>, anti-spasmodic, antisecretory, and mucosal protective effects<sup>10</sup>. It also has significant lipid lowering effect being a promising cardioprotective agent<sup>15</sup>. Various different compounds have also been isolated from its seeds, leaves, inflorescence and fruits<sup>16</sup>. The presence of flavonoids, phenolic compounds and essential oil in the seeds of *A. graveolens* make it an important component for the preparation of gripe water<sup>17</sup>. Therefore, the present study deals with the evaluation of the bioactive potential of *A. graveolens* with special reference to antioxidant potential in addition to their phytochemical analysis.

## MATERIAL AND METHODS

### Plant Material

Fresh leaf of *Anethum graveolens* (500g) were collected from the local area of Allahabad-U.P. (India) and authenticated by Professor Satya Narayan, Taxonomist Department of Botany, University of Allahabad, India (MRL/AG/03). The authenticated fresh leaves were washed and dried under shade. The shade dried leaves (180g) were mechanically crushed, powdered and preserved until further use.

### Chemicals

Folin-Ciocalteu reagent, Griess reagent, gallic acid, potassium ferricyanide, TCA (Tricarboxylic Acid), PMS (Phenazine Methosulphate), NADH (Nicotinamide Adenine Dinucleotide), quercetin, BHT (Butylated Hydroxy Toluene), ascorbic acid, sodium carbonate, FeCl<sub>3</sub>, DPPH (1, 1-Diphenyl-2-picrylhydrazyl, H<sub>2</sub>O<sub>2</sub>, KNO<sub>2</sub>, NaOH, MnO<sub>2</sub>, AlCl<sub>3</sub>, TPTZ (2, 4, 6-tripyridyl-s-triazine), FeSO<sub>4</sub>, HCl, SNP (Sodium Nitroprusside), NBT (Nitroblue Tetrazolium), Ferrozine, sodium salicylate, acetic acid, sodium acetate were purchased from Merck Chemicals, New Delhi. All other chemicals and reagents were of analytical grade.

### Preparation of Extract

The dried leaf powder (180g) was refluxed repeatedly thrice with double distilled water, for 2 hour each time and filtrates were collected, combined and concentrated in rotary evaporator at 45°C±5°C under reduced pressure, to obtain semisolid material, which was then lyophilized to get dark brown powder (yield ~15.5% w/w).

### Phytochemical Analysis

Screening for Phytochemicals was carried out based on the following analytical tests<sup>18-22</sup> as given in Table 1.

**Table-1**  
**Analytical Tests for Phytoconstituents**

Phytoconstituents	Colour Tests	Observation
Alkaloids ( <i>Hager's Test</i> ) ( <i>Wagner's Test</i> )	2ml extract + few drops of Hager's reagent 2 ml extract+ 2 drops HCl (1.5%) + 3c reagent	Yellow precipitate Brown precipitate
Anthraquinones ( <i>Borntrager's Test</i> )	3ml extract + 3ml Benzene + 5ml NH <sub>3</sub> (10%)	Pink, Violet or Red colouration
Coumarins	2ml extract + 3ml NaOH (10%)	Yellow colouration

Carbohydrate (Molisch's Test)	2ml extract (EtOH) + 10ml H <sub>2</sub> O + 2 drops ethanolic α-naphthol (20%) + 2 ml conc.H <sub>2</sub> SO <sub>4</sub>	Reddish violet ring at the junction
Emodins	2ml extract + 2ml NH <sub>4</sub> OH + 3ml Benzene	Red colouration
Flavonoids	1ml extract + 1ml Pb(OAc) <sub>4</sub> (10%)	Yellow precipitate
Glycosides (Liebermann's Test)	2ml extract + 2ml CHCl <sub>3</sub> + 2ml CH <sub>3</sub> COOH	Violet to Blue to Green colouration
Phlobatannins (Precipitate Test)	2ml extract + 2ml HCl (1%) + boil	Red precipitate
Proteins (Xanthoproteic test)	1ml extract + 1ml conc. H <sub>2</sub> SO <sub>4</sub>	White precipitate turned Yellow on boiling
Saponins (Foam Test)	5ml extract + 5ml H <sub>2</sub> O + heat	Froth appearance
Steroids (Salkowski's Test)	2ml extract + 2ml CHCl <sub>3</sub> + 2ml conc. H <sub>2</sub> SO <sub>4</sub>	Reddish Brown colouration at interface
Tannin (Braymer's Test)	2ml extract + 2ml H <sub>2</sub> O + few drops of FeCl <sub>3</sub> (5%)	Green colouration
Terpenoids	2ml extract + EtOH+2ml CHCl <sub>3</sub> + Δ(2 mint.) 3 drops conc. H <sub>2</sub> SO <sub>4</sub>	Deep red colouration

### Antioxidant Assays- in vitro

Antioxidant potential of *A. graveolens leaf extracts* (AGLE) was evaluated using different *in vitro* assays such as estimation of total phenolics, total flavonoids, and evaluation of free radical scavenging activity for DPPH<sup>•</sup>, NO<sup>•</sup>, superoxide and ABTS radicals. All the assays were carried out in triplicates and their average values were taken into consideration.

### Estimation of Total Phenolics

Total phenolic contents were estimated by the modified Folin-Ciocalteu method<sup>23</sup>. An aliquot (100 μl) of a range of different concentration, from 100 to 500 μg/ml, of AGLE was mixed with 2.5 ml Folin-Ciocalteu reagent (diluted with water 1:10 v/v) and 2.0 ml (75 g/l) of sodium carbonate. The tube was vortexed for 15 s and allowed to stand in dark at room temperature. The absorbance was measured at 750 nm using Shimadzu UV-VIS spectrophotometer after half an hour. The content of phenolics in extracts was expressed in terms of mg of Gallic Acid Equivalents (GAE) per g of dried extract.

### Estimation of Total Flavonoids

Total flavonoid contents were estimated using the method of Ordon et al.<sup>24</sup> Solutions of different concentrations from 100 to 500 μg/ml of AGLE were prepared separately in methanol. A volume of 0.5 ml of 2% of AlCl<sub>3</sub> solution was added to 0.5 ml of different concentrations of each sample. The absorbance was measured at 420 nm after one hour. The content of flavonoids in extracts was expressed in terms of mg of Quercetin Equivalent (QE) per g of dried extract.

### DPPH radical Scavenging assay

The effect of AGLE on DPPH radical scavenging was determined using the method of Liyana-Pathiranan and Shahidi.<sup>25</sup> About 1ml of sample in varied concentrations ranging from 100 to 500 μg/ml dissolved in deionised water was mixed with 1.0 ml of 0.135 mM DPPH. The mixture was vortexed thoroughly and left to stand for 30 min in the dark, and the absorbance was then measured at 517 nm against a control. Ascorbic acid (AA) was used as standard. The ability to scavenge DPPH radical was calculated by the following equation

$$\text{DPPH radical scavenging activity (\%)} = \left\{ \frac{A_c - A_s}{A_c} \right\} \times 100$$

where,  $A_c$  is the absorbance of control and  $A_s$  is the absorbance of sample.

### ABTS radical Scavenging assay

The ability of AGLE, to scavenge ABTS Cation radical was determined by Re et al.<sup>26</sup> The ABTS<sup>+</sup> radicals were pregenerated by mixing 7mM ABTS stock solution and 2.45 mM potassium persulfate. The solution was incubated for 12 hours in the dark at room temperature until the reaction was complete and the absorbance was

stable. 1 ml of the above solution was mixed with 1 ml of varied concentration of sample ranging from 25 to 125 μg/ml. After 6 minutes of incubation, absorbance was measured at 734 nm. Ascorbic acid (AA) was used as standard. The ability to scavenge ABTS<sup>+</sup> radical was calculated by the following formula:

$$\text{ABTS}^+ \text{ radical scavenging activity (\%)} = \left\{ \frac{A_c - A_s}{A_c} \right\} \times 100$$

where,  $A_c$  is the absorbance of Control and  $A_s$  is the absorbance of Sample.

### Nitrosyl radical Scavenging assay

Scavenging of nitrosyl radical by AGLE was determined by the method of Green et al.<sup>27</sup> Reaction started by incubating 0.5 ml SNP solution (10mM in Phosphate Buffered Saline) with varied concentration ranging from 100 to 500 μg/ml of sample at 25°C. After 150 minutes,

0.25 ml of sulphanilamide and 0.25 ml of NED were added. The mixture was allowed to stand for 30 minutes. Ascorbic acid (AA) was used as standard. The absorbance of sample was measured at 540 nm. The ability of sample to scavenge NO radical was calculated by the following formula:

$$\text{NO radical scavenging activity (\%)} = \{A_C - A_S / A_C\} \times 100$$

where,  $A_C$  is the absorbance of control and  $A_S$  is the absorbance of Sample.

#### Hydroxyl radical Scavenging assay

Hydroxyl radical scavenging activity of AGLE was measured according to a modified method of Smirnoff and Cumbes.<sup>28</sup> The OH radicals were generated from the mixture of  $\text{FeSO}_4$  and  $\text{H}_2\text{O}_2$ . The reaction mixture contained 1 ml  $\text{FeSO}_4$  (1.5mM), 0.7ml  $\text{H}_2\text{O}_2$  (6mM), 0.3 ml sodium salicylate (20mM) and 1ml of varied

concentrations of sample ranging from 100 to 500 $\mu\text{g/ml}$ . After incubation, for 1 h at 37°C, the absorbance of the complex was measured at 562 nm against control. Ascorbic acid (AA) was used as standard. The ability to scavenge OH radical was calculated by the following equation:

$$\text{OH radical scavenging activity (\%)} = [1 - \{A_S - A_B / A_C\}] \times 100$$

where,  $A_C$  is the absorbance of control and  $A_S$  is the sample,  $A_B$  is the absorbance of blank (without sodium salicylate).

#### Superoxide Anion radical Scavenging assay

Superoxide anion radicals were generated by the PMS/NADH system according to the method of Kakkar et al.<sup>29</sup>. The reaction mixture was composed of 1 ml of NBT (156  $\mu\text{M}$ ), 1 ml NADH (468  $\mu\text{M}$ ) and 100  $\mu\text{l}$  of AGLE ranging from 20- 100  $\mu\text{g/ml}$ . The reaction was

started by addition of 100  $\mu\text{l}$  of PMS (60  $\mu\text{M}$ ) to the mixture. After 5 min incubation at 25°C, absorbance was measured at 560 nm. Ascorbic acid (AA) was used as standard. The ability to scavenge  $\text{O}_2^-$  radical was calculated by the following equation:

$$\text{O}_2^- \text{ radical scavenging activity (\%)} = \{A_C - A_S / A_C\} \times 100$$

where,  $A_C$  is the absorbance of control and  $A_S$  is the absorbance of sample.

#### Metal Chelating Assay

The chelation of ferrous ions by AGLE was determined by the method of Dinis et al.<sup>30</sup> A solution of 2mM  $\text{FeCl}_2$  (0.05 ml) was added to varied concentration from 100 to 500 $\mu\text{g/ml}$  of sample. The reaction was initiated by the addition of 5mM ferrozine (0.2 ml) and the mixture was

shaken vigorously and kept at room temperature for 10 min. The absorbance of the reaction mixture was measured at 562 nm against control. Ascorbic acid (AA) was used as standard. The ability to chelate ferrous ions was calculated by the following equation

$$\text{Metal chelating activity (\%)} = \{A_C - A_S / A_C\} \times 100$$

where,  $A_C$  is the absorbance of control and  $A_S$  is the absorbance of Sample.

## STATISTICAL ANALYSIS

Data was statistically evaluated using two-way ANOVA, followed by a post hoc Scheffe's test using the statistical package PRISM 3.0 version. The values were considered significant when  $P < 0.05$ .

## RESULTS AND DISCUSSIONS

#### Phytochemical Screening

Table 2, reveals the results of phytochemical screening of *Anethum graveolens* leaf which indicates the presence of high amount of Carbohydrates, glycosides flavonoids and tannins. Though, alkaloids, proteins and saponins were also present but in lesser extent. On the contrary phytochemical screening also revealed the complete absence of terpenoids, coumarins, anthocyanins, anthraquinones, emodins, leucoanthocyanins, phlobatanins, and steroids. Thus the medicinal values of the plant leaves may be due to these specific classes of phytochemicals present in it. Plants containing carbohydrates and glycosides are

known to exert a beneficial action on immune system by increasing body strength and hence are valuable as dietary supplements<sup>42</sup>. Glycosides also have vast therapeutic efficacy as they are found in almost every medicinal plant. Flavonoids and other phenolics compounds also acts as a potent water-soluble antioxidant and free radical scavenger, which prevent oxidative cell damage<sup>31-33</sup>. It also helps in managing diabetes induced oxidative stress<sup>43</sup>. Tannins are known to be useful in the treatment of inflamed or ulcerated tissues and they have remarkable anticancer property<sup>44</sup>. Moreover, alkaloids represent a class which affects the central nervous system, reduces appetite and behaves as diuretic<sup>45</sup>. Proteins are the building blocks of life. The body needs protein to repair and maintain itself. Numerous studies have confirmed that saponins possess the unique property of precipitating and coagulating red blood cells<sup>34,35</sup>. Thus, the AGLE has a range of phytochemical compounds having therapeutic efficacy with special reference of antioxidant which can effectively protect from oxidative damage caused by free radicals.

**Table 2**  
**Phytochemical Screening of Aqueous Extract Of *A. Graveolens* Leaf**

Phytoconstituents Tested	Results Observed
Tannins	++
Carbohydrates	++
Glycosides	++
Flavonoids	++
Alkaloids (Wagner's Test); (Hager's Test)	+
Proteins	+
Saponins	+
Terpenoids	-
Steroids	-
Phlobatannins	-
Coumarins	-
Antraquinones	-
Anthocyanins	-
Leucoanthocyanins	-
Emodins	-

(++) = Significant Presence, (+) = Presence, (-) = Absence

### Antioxidant assay - in vitro

#### Total Phenolics

Table 3, shows the results of total phenolic content of *A. graveolens* leaf extracts expressed as Gallic acid Equivalents (GAE). Standard curve equation of Gallic acid was calculated with  $y = 0.0002x + 0.029$ ,  $R^2 = 0.995$ . Total phenolic content was estimated 65.5 mg/g gallic acid equivalent. Thus, *A. graveolens* leaves were found to be a potential source of phenolics and hence could be explored as natural antioxidant. Phenolics are the well-known group of secondary metabolites and comprise a large group of biologically active compounds<sup>36</sup>. The antioxidant activities of phenols were credited to their redox properties, which permit them to act as reducing agents, hydrogen donors and singlet

oxygen quenchers, as well as their metal chelating abilities.

#### Total Flavonoids

Table 3, also shows the results of total flavonoid content of *A. graveolens* leaf extracts, expressed as Quercetin Equivalent (QE). Standard curve equation of Quercetin was calculated with with  $y = 0.0536x - 0.0038$ ,  $R^2 = 0.968$ . Total flavonoid content was estimated 29.8mg/g quercetin equivalent. Flavonoids are important secondary metabolites present in plants which modulate lipid peroxidation involved in atherogenesis, thrombosis and carcinogenesis. It is well established that pharmacological effect of flavonoids is associated with antioxidant activities<sup>37</sup>.

**Table 3**  
**Quantitative Estimation Of Total Phenolics And Flavonoids Of *A. Graveolens* Leaf**


Total Phenolic contents (mg/g of GAE)	Total Flavonoid contents (mg/g of QE)
65.5±1.23	29.8± 0.32

### Free Radical Scavenging Activity

#### DPPH Radical

Fig. 1 clearly reveals the results of scavenging activity of AGLE and AA was found to exhibit 87.57 and 96.47% inhibition, respectively at the highest evaluated concentration of 400µg/ml. Moreover, Fig. 7 gives an

account of IC<sub>50</sub> values of AGLE and AA which were found to be 122.4 and 100.24µg/ml, respectively. The data validates the primary antioxidant nature of *A. graveolens* leaves which could be due to hydrogen donating ability of its phenolic constituents.




**Figure 1**  
**DPPH radical scavenging activity of *A. graveolens* leaf extract and ascorbic acid**

**Nitric oxide Radical Scavenging**

Fig. 2 show the results of NO radicals scavenging activity of AGLE and AA was found to exhibit 68.29 and 83.43% inhibition, respectively at the highest evaluated concentration of 400µg/ml. Moreover, Fig. 7 gives an account of IC<sub>50</sub> values of AGLE and AA which were found to be 244.3 and 114.61 µg/ml, respectively. The data is confirming *A. graveolens* leaf could be considered as better antioxidant due to their NO<sup>•</sup> scavenging potential, which is widely used as a

measure of antioxidant activity of plant extract. Nitric oxide radical (NO<sup>•</sup>) is known to be a ubiquitous free radical distributed in tissues or organ systems and have a vital role in neuromodulation or neurotransmitter in the central nervous system. In this assay, SNP gets decomposed in the aqueous solution at physiological pH 7.4 and produced NO radical. This NO radical reacts with oxygen to produce nitrate and nitrite, which was determined by using Griess reagent<sup>38</sup>.




**Figure 2**  
NO radical Scavenging activity of *A. graveolens* leaf extract and ascorbic acid

**Superoxide Radical**

Fig.3, depicts the results of Superoxide anion radical scavenging activity of both, the Extract and the Standard at varied range of concentration. Results depicts that the scavenging ability in both the cases was concentration dependent and was found to be maximum at a concentration of 400µg/ml with inhibition of 82.32% in

case of AGLE and 92.46% in case of Standard, validating thereby the significant antioxidant potential of *A. graveolens* leaf extract. The IC<sub>50</sub> values of the Extract and the Standard were found to be 173µg/ml and 117.22µg/ml respectively, which further confirmed that *A. graveolens* leaves could be developed as an effective antioxidant agent.




**Figure 3**  
Superoxide radical scavenging activity of *A. graveolens* leaf extracts and ascorbic acid

**ABTS cation Radical**

Fig.4, exhibits the results of ABTS cation radical scavenging activity of both, the Extract and the Standard at varied range of concentration. Results depicts that the scavenging ability in both the cases was concentration dependent and was found to be maximum at a concentration of 150µg/ml with inhibition of 97.05% in case of AGLE and 99.12% in case of Standard, validating thereby the significant antioxidant potential of *A. graveolens* leaf extract. The IC<sub>50</sub> values of the Extract and the Standard were found to be 33.45µg/ml and 66.61µg/ml respectively, which further confirmed that *A.*

*graveolens* leaves could be developed as an effective antioxidant agent. The data shows appreciably higher scavenging activity of *A. graveolens* leaf extract against ABTS<sup>•+</sup> radical. This implies that the extract may be useful for treating radical related pathological damage. Proton radical scavenging is an important attribute of antioxidants. It is often used for screening complex antioxidant mixtures such as plant extracts, beverages and biological fluids. The stability of ABTS<sup>•+</sup> for more than 2 days over a wide pH range raised the interest in the use of ABTS<sup>•+</sup> for the estimation of antioxidant activity.




**Figure 4**  
**ABTS radical scavenging activity of *A. graveolens* leaf extracts and ascorbic acid**

#### Hydroxyl Radical

Fig.5, exhibits the results of hydroxyl radical scavenging activity of both, the Extract and the Standard at varied range of concentration. Results depicts that the scavenging ability in both the cases was concentration dependent and was found to be maximum at a concentration of 400µg/ml with inhibition of 74.02% in case of AGLE and 92.22% in case of Standard, validating thereby the significant antioxidant potential of *A. graveolens* leaf extract. The IC<sub>50</sub> values of the Extract

and the Standard were found to be 116.42µg/ml and 20.54 µg/ml respectively, which further confirmed that *A. graveolens* leaf could be developed as an effective antioxidant agent. Hydroxyl radicals are known to be the most reactive of all the reduced forms of dioxygen and well known for initiating cell damage in vivo<sup>39</sup> and causing lipid peroxidation<sup>40</sup>. The evaluation of radical scavenging activity was based on the generation of OH radical by Fenton reaction<sup>46</sup>.




**Figure 5**  
**Hydroxyl radical scavenging activity of *A. graveolens* leaf extract and ascorbic acid**

#### Metal Chelating

Fig.6, depicts the results of metal chelating activity of both, the Extract and the Standard at varied range of concentration. Results depicts that the scavenging ability in both the cases was concentration dependent and was found to be maximum at a concentration of 400µg/ml with inhibition of 61.51% in case of AGLE and 85.18% in case of Standard, validating thereby the significant antioxidant potential of *A. graveolens* leaf extract. The IC<sub>50</sub> values of the Extract and the Standard were found to be 267.89µg/ml and 51.30µg/ml respectively, which further confirmed that *A. graveolens*

leaves could be developed as an effective antioxidant agent. Metal chelating assay is also a simple yet an important assay as ferrozine can quantitatively form complexes with Fe<sup>2+</sup> producing a intense violet Fe<sup>2+</sup>/ferrozine complex. The chelating activity of antioxidant was measured by monitoring the decrease in the intensity of violet colour of the complex<sup>41</sup>. Chelating agents may inhibit lipid oxidation by stabilising transition metals. In this assay, the ability of antioxidants to disrupt the formation of the complexes or to prevent interaction between metal and lipid was measured in terms of their iron-chelating capacity.



**Figure 6**  
**Metal Chelating activity of *A. graveolens* leaf extract and ascorbic acid**




Figure 7

Free radicals Scavenging potential and Metal chelating activity of AGLE and AA in terms of IC<sub>50</sub> (each value represents a mean ± SD (n=3))

## CONCLUSION

Conclusively it could be stated that *A. graveolens* leaf can be explored further in order to develop a novel antioxidant agent with high therapeutic efficacy due to high content of their polyphenolics including tannins, carbohydrates and glycosides in addition to flavanoids. The highly significant antioxidant efficacy of *A. graveolens* leaf was also evident from its high free radical scavenging potential and low IC<sub>50</sub> values in comparison to the standard drug, ascorbic acid, taken as reference.

## REFERENCE

- Bobbarala V, Katikala PK, Naidu KC, Penumajji S. Antifungal activity of selected plant extracts against phytopathogenic fungi *Aspergillus niger*. Indian J. Sci Technol. 2009 April; 2 (4): 87-90.
- Goh SH, Chuah CH, Mok JSL, Soepadmo E. Malaysian Medicinal Plants for the Treatment of Cardiovascular Diseases. Selangor Darul Ehsan: Pelanduk Publication. Kaula Lumpur, Malaysia; 1995. p. 162.
- Pandey P, Mehta R, Upadhyay R. Physicochemical and preliminary phytochemical screening of *Psoralea corylifolia*. Arch Appl Sci Res. 2013; 5:261-5.
- Halliwell B. Reactive oxygen species in living systems: Source, biochemistry, and role in human disease. Am J Med Sci. 1991 Sep 30; 91(3C): 14-22.
- Power M, Pratley R. Alternative and complementary treatments for metabolic syndrome. Curr Diab Rep. 2011 Jun; 11 (3): 173-178.
- Sharma RKS, Chatterji S, Rai DK, Mehta S, Singh RK, Rai PK, et al. Antioxidant potential and phenolic contents of the aqueous extract of some Indian medicinal plants. J Med Plants Res. 2009 Nov; 3 (11): 944-948.
- Rai PK, Jaiswal D, Rai DK, Sharma B, Watal G. Antioxidant potential of oral feeding of *Cynodon dactylon* extract on diabetes-induced oxidative stress. J Food Biochem. 2010; 34: 78-92.

## ACKNOWLEDGEMENT

The first author, Amrita Kumari Srivastava is thankful to University Grants Commission (UGC), New Delhi, India for providing financial assistance in the form of Senior Research Fellowship (NET-SRF).

## CONFLICT OF INTEREST

Conflict of interest declared none.



8. Bahramikia S, Yazdanparast R. Efficacy of different fractions of *Anethum graveolens* leaves on serum lipoproteins and serum and liver oxidative status in experimentally induced hypercholesterolaemic rat models. *Am J Chin Med* 2009; 37 (4): 685-699.
9. Bahramikia S, Yazdanparast R. Antioxidant and Free radical scavenging activities of different fractions of *Anethum graveolens* leaves using in vitro model. *Pharmacology online*. 2008; 2: 219-33.
10. Radulescu V, Lidiapopescu M, Lies D C. Chemical composition of the volatile oil from different plant parts of *Anethum graveolens* L. (umbelliferae) cultivated in Romania. *Farmacia*. 2010; 58(5): 594-600.
11. Kaur GJ, Arora S. Bioactive potential of *A. graveolens*, *Foeniculum vulgare*, *Trachyspermum ammi* belonging to the family Umbelliferae, Current status. *J Med Plant Res*. 2010 Jan 18; 4(2): 87-94.
12. Kazemi M, Rostami H, Shafiei S. Antibacterial and Antifungal Activity of some Medicinal Plants from Iran. *J Plant Sci*. 2012; 7(2): 55-66.
13. Albayrak S, Aksoy A, Sagdic O, Albayrak S. Antioxidant and antimicrobial activities of different of some medicinal herbs consumed as tea and spices in Turkey. *J Food Biochem*. 2012; 36: 547-554.
14. Shyu YS, Lin JT, Chang YT, Chiang CJ, Yang DJ. Evaluation of antioxidant ability of ethanolic extract from dill (*Anethum graveolens* L.) Flower. *Food Chem*. 2009; 115(2): 515-521.
15. Hajhashemi V, Abbasi N. Hypolipidemic activity of *Anethum graveolens* in rats. *Phytother Res*. 2008; 22: 372- 34.
16. Santos AG, Figueiredo AC, Lourenco PML, José G Barroso, Luis G Pedro, Margarida Oliveira M, et al. Hairy root cultures of *Anethum graveolens* (dill): establishment, growth, time-course study of their essential oil and its comparison with parent plant oils. *Biotech Lett*. 2002; 24: 1031-1036.
17. Heamalatha S, Swarnalatha S, Divya M, Gandhi LR, Ganga DA, Gomathi E. Pharmacogonostical, Pharmacological, Investigation on *Anethum graveolens* Linn: A Review. *Res J Pharm Biol Chem Sci*. 2011Dec; 2(4):564-574.
18. Santosh CH, Attialla IH, Mohan MM. Phytochemical analysis, antimicrobial and antioxidant activity of ethanolic extract of *Vernonia anthelmintica*. *Int J Pharm Bio Sci*. 2013; 4(1): 960-966.
19. Yadav M, Chatterji S, Gupta SK, Watal G. Preliminary phytochemical screening of six medicinal plants used in traditional medicine. *Int J Pharm Pharm Sci*. 2014; 6(5): 539-542.
20. Rana Md M, Sayeed Md A, Nasrin Mst S, Islam M, Rahman Md M, Alam MF. Free radical scavenging potential and phytochemical analysis of leaf extract from *Oscimum sanctum* Linn. *J Agric Tech* 2015; 11(7): 1615-1623.
21. Angappan MS, Karuppaiah J. Screening of phytochemical and in vitro antioxidant property of n-miracle (polyherbal formulation). *World J Pharm Sci*. 2015 April 02; 4(6): 1702-1717.
22. Devi MR, Krishnakumari S. Quantitative estimation of primary and secondary metabolites in hot aqueous extract of *Pleurotus sajor caju*. *J Pharmacogn Phytochem*. 2015 Aug 16; 4(3): 198-202.
23. Wolfe KWX, Liu RH. Antioxidant activity of apple peels. *J Agric Food Chem*. 2003 Jan 29; 51(3): 609-614.
24. Ordon EEAL, Gomez JD, Vattuon MA, Isla MI. Antioxidant activities of *Sechium edule* (Jacq.) Swart extracts. *Food Chem*. 2006 Aug; 97(3): 452-458.
25. Liyana-Pathiranan CM, Shahidi F. Antioxidant activity of commercial soft and hard wheat (*Triticum aestivum* L) as affected by gastric pH conditions. *J Agric Food Chem*. 2005 Apr 6; 53 (7): 2433-2440.
26. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med*. 1999 May; 26 (9-10): 1231-1237.
27. Green LC, Wanger DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. Analysis of nitrate, nitrite, [15 N] nitrate in biological fluids. *Anal Biochem*. 1982 Oct; 126 (1):131-138.
28. Smirnoff N, Cumbes QJ. Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry*. 1989; 28(4): 1057-1060.
29. Kakkar P, Das B, Viswanathan PN. A modified spectrophotometric assay of superoxide dismutase. *Indian J Biochem Biophys*. 1984 Apr; 21: 130-132.
30. Dinis TC, Madeira VM, Almeida LM. Action of phenolic derivatives (acetaminophen, salicylate and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation as peroxyl radical scavengers. *Arch Biochem Biophys*. 1994 Nov 15; 315(1): 161-9.
31. Shi J, Yu J, Pohorly J, Young C, Bryan M, Wu Y. Optimization of the extraction of polyphenols from grapes seed meal by aqueous ethanol solution. *Food Agric Environ*. 2003; 1(2): 42-47.
32. Rio DA, Obdulio BG, Casillo J. Marin FG, Ortuno A. Uses and properties of citrus flavonoids. *J Agric Food Chem*. 1997 Dec; 45(12):4505-4515.
33. Salah N, Miler NJ, Pagange G, Tijburg L, Bolwell GP, Rice E. Polyphenolic flavonoids as scavenger of aqueous phase radicals as chain breaking antioxidant. *Arch Biochem Broph*. 1995 Oct 1; 322(2):339-46.
34. Okwu DE. Phytochemicals and vitamin content of indigenous spices of southeastern Nigeria. *J Sustain Agric Environ*. 2004; 6(1):30-37.
35. Sodipo OA, Akiniyi JA, Ogunbamusu JU. Studies on certain characteristics of extracts of bark of *Pansinystalia macruceras* (Kschemp) Pierre Exbeille. *Global J Pure Appl Sci*. 2000; 6(1):83-87.
36. Dreosti IE. Antioxidant polyphenols in tea, cocoa, and wine. *Nutrition*. 2000; 16(7-8): 692-694.
37. Shi J Yu J, Pohorly J, Young CJ, Bryan M, Wu Y. Optimization of the extraction of polyphenols from grapes seed meal by aqueous ethanol solution. *Food Agric Environ*. 2003; 1(2): 42-47.

38. Adedapo AA, Jimoh FO, Afolayan AJ, Masika PJ. Antioxidant Properties of the Methanol Extracts of the Leaves and Stems of *Celtis Africana*. Rec Nat prod. 2009; 3(1): 23-31.
39. Rollet-Labelle E, Grange MJ, Elbim C, Marquetty C, Goggerot-Pocidallo MA, Pasquier C. Hydroxyl radical as a potential intracellular mediator of polymorphonuclear neutrophil apoptosis. Free Rad Biol Med. 1998 Mar 1; 24(4): 563-572.
40. Aurand LW, Boone NH, Giddings GG. Superoxide and singlet oxygen in milk lipid peroxidation. J Dairy Sci. 1977; 60: 363-369.
41. Satoh K, Sakagami H. Effect of metal ions on radical intensity and cytotoxic activity of ascorbate. Anticancer Res. 1997 Mar- Apr; 17(2A): 1125-1129.
42. Patil US, Deshmukh O S. Preliminary Phytochemical Screening of six medicinal plants used as Traditional Medicine. Int J Pharm Bio Sci. 2016 Jan; 7(1): 77 – 81.
43. Yadav M, Chatterji S, Gupta S K, Watal G. Preliminary Phytochemical Screening of Six medicinal Plants used in traditional medicine. Int J Pharm Sci. 2014; 6(5): 539-542.
44. Okwu DE. Phytochemicals and Vitamin content of indigenous spices of South Eastern Nigeria. Journal of sustenance of Africa Environment, 2004; 6:30-3.
45. United States Department of Agriculture. Center for Nutrition Policy and Promotion. Dietary Guidelines for Americans. National Academy Press, Washington DC: USA, (2010).
46. Klein SM, Cohen G, Cederbaum AI. Production of formaldehyde during metabolism of dimethyl sulphoxide by hydroxyl radical generating system. Biochemistry. 1991;20:6006–6012.

## Reviewers of this article

### **Dr. Sanjukta Chatterji**

Research Associate, Department of  
Chemistry,  
University of Allahabad,  
Allahabad - 211002  
U.P. India



### **Asst. Prof. Dr. Sujata Bhattacharya**

Assistant Professor, School of Biological  
and Environmental Sciences, Shoolini  
University, Solan (HP)-173212, India



### **Prof. Dr. K. Suriaprabha**

Asst. Editor, International Journal  
of Pharma and Bio sciences.



### **Prof. P. Muthuprasanna**

Managing Editor, International  
Journal of Pharma and Bio sciences.

We sincerely thank the above reviewers for peer reviewing the manuscript