



FATTY ACID COMPOSITION OF BHALLATAKA OIL AND THEIR BIOLOGICAL PROPERTIES

VUSTELAMURI PADMAVATHI AND BHATTIPROLU KESAVA RAO*

*Prof.B.Kesava Rao, Chairman-BOS, Department of Chemistry,
University College of Sciences, Acharya Nagarjuna University,
Nagarjunanagar- 522 510, Guntur District, Andhra Pradesh, India.

ABSTRACT

Semecarpus anacardium L.f is a plant well known for its therapeutic value in ayurvedic and siddha system of medicine, in our survey we found that, the Anacardiaceae family has 77 genera and 850 species surviving as trees, shrubs and vines. Out of these, the commonly known Cashew family (or) Sumac family has great viable importance. Due to the presence of several anticancer drugs isolated from Anacardiaceae family, we have selected *Semecarpus anacardium* L.f, for its high medicinal value in Ayurveda and Siddha systems and isolated numerous active constituents. In our research, *Bhallataka oil* (cape 32) found to have a free fatty acid (0.50) hence; alkaline *Tran's*-esterification was done by using anhydrous methanol at a molar ratio of 6:1 and 3g/ Litre of sodium hydroxide as catalyst, and the results showed that Fatty acids with highest weight percentage 18:1 - 51.25403 %, 16: 0 -13.41810 %, 18:3 - 6.042404 % and also it consists of the antimicrobial activity against both gram positive and gram negative bacteria.

KEY WORDS: *Anacardiaceae family, Semecarpus anacardium L.f, Fatty acid methyl esters, Antimicrobial activity.*



BHATTIPROLU KESAVA RAO*

Prof.B.Kesava Rao, Chairman-BOS, Department of Chemistry, University College of Sciences, Acharya Nagarjuna University, Nagarjunanagar- 522 510, Guntur District, Andhra Pradesh, India.

Received on: 17-05-2017

Revised and Accepted on : 30-05-2017

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.3.p81-92>

INTRODUCTION

Herbal drugs are contributing a lot to human health in our tradition especially in 21st century. Natural medicine progresses the inner immune system of the human body and there will be no side effects. Hence, the herbal drug acts more effectively than the modern medicine. The word, *Semecarpus*¹ is derived from Simeion in Greek means marking/tracing and carpus in Greek means nut. *Anacardium* means like cardium, i.e. heart shaped marking nut. *Bhallatak* literary means sharp like spear. *Bhallatak* or marking nut is known to the world since ancient times. The stem, sap, fruit and seeds are used by mankind for diverse purposes such as timber, paint, waterproofing, food and medicine. In view of its several potent medicinal properties, it is acclaimed as *Ardhavaidya* in Ayurveda and as a Golden acorn at the time of Galen in the western world. Severe skin manifestations, precludes its use for medical purposes. In India, the plant is used by Ayurvedic practitioners, traditional healers across the country albeit with caution. However, manufacturing units are overtly scared in view of its apparent toxic nature, and hence there are a few takers. The plant belonging to *Anacardiaceae* family has potential to produce allergic manifestations through contact dermatitis. Phytoconstituents, viz. alkyl catechols, phenols, quinols and resorcinols are believed to be responsible for skin reactions. The plant growing naturally in tropical and dry climate is a deciduous tree having a height of about 10 m. Flowering time is

May to October, whereas fruiting time is December to March. The fruit is eaten when ripe; whereas the kernel of the seed is eaten after removing the pericarp. Marking nut seed, which sinks in water, is used for medicinal purposes, Ayurvedic properties of *Bhallatak* are madhur, kashay ras, ushna virya, madhur vipak and laghu, snigdha, tikshna, and ushna gunas². It has several karmas like *Kaphavatashamak* (alleviates kapha & Vata dosha), *Bhootanashan* (anti-devil) *Pittasanshodhak* (expels out pitta dosha), *Medhya* (beneficial to brain), *Vanhikar* (improves digestive fire), *Vrishya* (aphrodisiac), *Chedana* (excisional functions), *Bhedan* (incisional function), *Bruhan* (anabolic in effect), and hence indicated for many diseases like *Arsha* (haemorrhoids), *Udar* (ascites), *Grahani* (inflammatory bowel diseases), *Shotha* (inflammation), *Krumi* (helminthiasis), *Kushtha* (skin disorders, like psoriasis), *Vran* (wounds), *Shwittra* (vitiligo), *Gulma* (abdominal mass), *Jwar* (fever), *Adhman* (flatulence), etc.

PHARMACOLOGICAL ACTIVITIES

Recent studies suggested that single and compound formulations of *Semecarpus anacardium* L.f. has multiple pharmacological activities such as anti-inflammatory activity³, anti-arthritic activity⁴, hypoglycaemic activity⁵, Anti-cancer activity⁶, contraceptive agent⁷, lipoxygenase inhibitory activity⁸, hypolipidemic activity⁹ & hypocholesterolemic activity⁹, antimicrobial activity¹⁰, antistress activity¹¹ and immune modulatory activity¹² in (Fig:1)




Figure1
Pharmacological activity properties shown by *Semecarpus anacardium* L.f.

MATERIALS AND METHODS

Semecarpus anacardium L.f. nuts were collected from field area very nearer to the village Nandgaon, Kolhapur City, Maharashtra, India. All plant material specimens

were identified by Dr Vatsavaya S. Raju, Plant Systematic Laboratory, Department of Botany, Kakatiya University, Warangal (A. P. State) and conformed as *Semecarpus anacardium* L.f. (syn: *Anacardium latifolium* Lam., *A. orientale* Steud.) of *Anacardiaceae*

and plant specimen deposited at Kakatiya University Herbarium, Warangal (KUW) with accession number 1874. It is locally known as 'nalla jeedi' and popularly known as 'marking nut/dhobi nut'.

Extraction and purification

The crushed nuts (3.5 kg) were extracted with cold petroleum ether. Solvent from this extract was removed under reduced pressure. 129gms of blackish gummy substance was obtained, this was designated as *Semecarpus anacardium* cold hexane extract (SaCHE). The residue left after cold extraction was Soxhleted with hot petroleum ether (60-80^o) and this was concentrated under reduced pressure which resulted blackish gummy mass of 680gms and was designated as *Semecarpus anacardium* hot hexane Extract (SaHHE). The left residue was extracted with

cold acetone; the solvent from this extract was removed under reduced pressure which resulted dark blackish substance of 420gms which designated as *Semecarpus anacardium* cold acetone extracts (SaCAE). Finally the residue was Soxhleted with hot acetone, the solvents from this extract was removed under reduced pressure, resulted the yield of dark reddish brown substance of 959gms which is designated as *Semecarpus anacardium* hot acetone extract (SaHAE). All the four filtrates (SaCHE, SaHHE, SaCAE, and SaHAE) were subjected for column chromatography on silica gel and column was eluted with increasing polarity of solvents individually. And the oil obtained from the fraction of CAPE (cold acetone petroleum ether extract) 32 was under reduced pressure the oil (named CAPE 32 (Bhallataka Oil-3)) was concentrated by vacuum evaporation and given for spectral analysis.

Process of Flow Diagram of *Semecarpus anacardium* L.f Nut oil (Bhallataka Oil-3).




Figure 2
Process of Flow Diagram of *Semecarpus anacardium* L.f Nut oil (Bhallataka Oil-3).

RESULTS AND DISCUSSION

In our investigation, the seed oil of *Semecarpus anacardium* L.f (Bhallataka oil) has number of bioactive compounds and the Oil was characterized by FT-IR, ^1H NMR, ^{13}C NMR, GC methods, it was found that it contains free fatty acids, triglycerides. Bhallataka Oil-3 was obtained from the cold acetone extract of 31-32 fractions which was eluted with Hexane, the fractions were examined by TLC, individually, Basing on TLC, fractions from 31-32 were showing similar spot pattern and designated as CAPE-32, it was obtained as Brown oil with B.P 74-78.5°C. The FT-IR Spectrum, displayed bands at 3474.70 cm^{-1} (O-H, hydrogen bonded alcohol, phenols), 3006.14 cm^{-1} (hydrogen bonded alcohol, phenols), 2925.96 cm^{-1} , 2855.77 cm^{-1} (CH, alkanes), 1746.76 cm^{-1} (C=O, aldehydes, ketones, carboxylic acids, esters), 1461.47 cm^{-1} , 1376.63 cm^{-1} (C-H, alkanes), 1237.81 cm^{-1} , 1118.90 cm^{-1} , 1163.85 cm^{-1} (C-O, alcohol, acids, ethers, carboxylic acids, esters.), 722.78 cm^{-1} (C-H, alkanes). The proton NMR Spectrum of Bhallataka Oil-3 (CAPE 32) : 5.360-5.323 δ ppm is due to the unsaturated acids -CH=CH-Olefinic protons, The Sn-2 proton (attached to the carbon CHO) of the Glycerol backbone of the triacyl glycerols causes the small clusters of the peaks at 5.220

ppm, 4.322 – 4.113 ppm due to Sn-1 and Sn-3 protons (attached to the two terminal carbons; CH_2O) of the Glycerol backbone of the triacylglycerides, 2.789-2.751 ppm [bisallylic protons, i.e., protons attached to carbons situated directly next to two C=C double bonds -CH=CH- CH_2 -CH=CH-], 2.334-2.284 ppm [Protons on the second carbon in the fatty acid chains -O-C(=O)- CH_2], 2.019-2.000 ppm due to allylic protons i.e., protons attached to carbons next to C=C double bonds, 1.606 ppm due to Methylene protons due to all acyl chain - CH_2 - CH_2 -COOH, 1.301-1.255 ppm. -(CH_2) $_n$ - due to all acyl chains, 0.109-0.857 due to Terminal methyl protons -(CH_2) $_n$ - CH_3

^{13}C NMR (CDCl_3)

173.228, 172.804 (C-1, Sn-2 due to triacylglyceride moiety), 130.175 (C-10 due to oleyl), 129.965, 129.657 (C-9 oleyl), 128.031, 127.851 (Linoleyl), 68.856 (Triacylglycerides), 62.070 (CH_2 -Sn-1,3 Triacylglycerides), 35.810, 34.164, 34.003 (C-2, Sn-1,3 acyl chain), 31.884, 31.750 (W_3 , sat., n-9 and n-6 acids), 31.497, 31.262, 30.878, 29.679, 29.500, 29.295, 29.092, ((CH_2) $_n$ acyl chains), 27.174 (due to oleyl and linoleyl), 25.599 (Diallylic, linoleyl, linolenyl), 22.655, 22.544, 14.0073 and 13.768 (due to acyl chain).


Fatty acid composition of Bhallataka Oil-3

Table 1
Fatty acid composition of Bhallataka Oil-3

S.No	LIPID VALUE	FATTY ACID	WT%
1	12:0	Lauric acid	0.18
2	14:0	Myristic acid	0.25
3	16:0	Palmitic acid	13.41
4	16:1	Palmitoleic acid	0.1474
5	18:1	Oleic acid	51.254
6	18:2	Linoleic acid	5.639
7	18:3	α -Linoleic acid	5.320
8	20:0	Arachidic acid	3.819
9	20:1	Paullinic acid	1.09
10	20:2	Eicosadienoic acid	1.189
11	22:0	Behenic acid	5.076
12	23:0	Dihomo gammaLinolenic acid	1.02
13	24:0	Tetracosanoic acid	1.18

The structural elucidation of the Bhallataka Oil-3 is finalized by the linkage of connecting the major composition of the fatty acids by the increasing order of Molecular weight of the fatty acids. According to this fatty acids present in Bhallataka Oil-3 are

(1) Palmitic acid: lipid value (16:0) with weight percentage 13.41



Palmitic acid

Figure 3
Palmitic acid

(2) Oleic acid: Lipid value (18:1) with weight percentage 51.254




Figure 4
Oleic acid

(3) α - Linoleic acid : Lipid value (18:3) with weight percentage 5.320




Figure 5
 α - Linoleic acid

The tentative structure of Bhallataka Oil-3 is




Figure 6
Structure of [(7Z, 10Z)-2-((Z)-octadec-7-enoyloxy)-3-(stearoyloxy) propyl]icoso-7,10-Dienoate

The proposed structure of Bhallataka Oil -3 (CAPE 32) is confirmed by the Spectral and chemical Trans-esterification.

Tran's -esterification process^{13, 14}

In our investigation, *Semecarpus anacardium* nut oil cape 32 had found to have a free fatty acid (0.50) hence; alkaline *Tran's*-esterification was done by using anhydrous methanol at a molar ratio of 6:1 and 3g/Liter of sodium hydroxide as catalyst. The processor was stirred at 600 rpm and at a temperature of 60⁰ C for 2 hours, after which the mixture was poured into a

decanter and allowed to settle for 3 hours so that the reaction can be driven to completion. By following mechanism as in Figure 6 and found that the mixture has been separated in to its corresponding methyl ester. The glycerol at the bottom in the ester was drained off by gravity. The excess methanol in the ester was removed by using a flash rotary evaporator. The impurities were removed from the methyl ester by washing with distilled water of volume ratio 3 to 1 three times. Finally, the washed methyl ester was dried by passing it through anhydrous sodium sulphate (Na₂SO₄).

Figure 7
General Mechanism of Trans-esterification process of Oil.




Figure 8
Structure of [(7Z, 10Z)-2-((Z)-octadec-7-enoyloxy)-3-(stearoyloxy) propyl icoso-7, 10-dienoate, Bhallataka oil-3(Kabala triacylglyceride) of major fatty acids.

The Identification of Cold acetone pet.ether 32 fraction of Nuts was Identified according to IUPAC Name was (7Z, 10Z)-2-((Z)-octadec-7-enoyloxy)-3-(stearoyloxy) propyl icoso-7, 10-dienoate] and found to be as new triacylglyceride "isolated from Semecarpus anacardium L.f was characterized and named as Kabala

triacylglyceride" Basing on UV, FT-IR, ¹H NMR, ¹³C NMR, MASS, HPLC, LCMS, GCMS, GC methods, it was found that it contains free fatty acids, triglycerides and others. Basing on spectral data and their physical properties Semecarpus anacardium L.f. nut oil was selected for biodiesel production.

The spectral data of CAPE 32 was given below as:




Figure 9
FT-IR of CAPE 32

Table 2
FT-IR Spectral data of Bhallataka Oil (CAPE 32)

CAPE -32	Functional group assignment	Group frequency cm-1
3474.70	Hydrogen bonded Alcohol, Phenols.	3200-3600
3006.14	Aromatic H-X group	3000-3100
2925.96	Alkanes	2850-3000
2855.77	Alkanes	2850-3000
1746.76	Saturated aldehyde	1720-1740
1461.47	Alkenes	1610-1680
1376.63	Alkanes	1340-1470
1237.81	Alcohol, Ether, Carboxylic acid, Ester	1050-1300
1163.85	Alcohol, Ether, Carboxylic acid, Ester.	1050-1300
1118.90	Alcohol, Ethers, Carboxylic acids, Esters.	1050-1300
722.78	Alkanes	679-995
	Alkanes	679-995
	Alkanes	679-995








Figure 10
¹H NMR of CAPE 32

Table 3
FT-IR Spectral data for Bhallataka Oil

Expanded form of CAPE 32 ¹H NMR

S.NO	PEAK	δ (ppm) Values	Assignment of compound
1		(5.44-5.22) δ (ppm)	Unsaturated acids -CH=CH- olefinic protons, the SN-2 protons (attached to the centre carbon; CHO) of the glycerol backbone of the triacylglycerols causes the small cluster of peak at 5.25 ppm.
2		(4.22-4.10, 4.40-4.22) δ (ppm)	SN-1 OR SN-3 unsaturated fatty acids -CH-OCOR, (attached to the two terminal carbons; CH ₂ O) of the glycerol back bone of the triacylglycerols.
3		(2.85-2.70) δ (ppm)	bis-allylic protons, i.e. protons attached to carbons situated directly next to two C=C double bonds, -CH=CH-CH ₂ -CH=CH-.
4		(2.40-2.25) δ (ppm)	Protons on the second carbon in the fatty acid chains-O-C(=O)-CH ₂ -.




Table 3
Expanded form of CAPE 32 ¹H NMR

¹³C NMR Spectral data for CAPE 32:

CAPE 32	peaks
173.228	C ₁ ,Sn-1,3
172.804	C ₁ ,Sn-2
130.175	C ₁₀
129.965	C ₉
129.657	
128.031	C ₁₀
127.851	C ₁₂
68.656	-CHO- Sn-2
62.070	CH ₂ O Sn-1,3
	C-2 Sn-2
35.810	c-2 Sn-2
34.164	C-2 Sn-1,3
34.003	
	C-2 Sn-1,3
	W3 saturated
31.884	
31.750	
31.497	saturated
31.262	
30.878	
29.679	(CH ₂)_ allyl chain
29.500	
29.295	
29.092	
27.174	Acyl chain c8- c11;c-8-c-14
25.599	Acyl chain
	Acyl chain
	Diallylic c-11,c-14
24.819	Acyl chain
22.655	Acyl chain
22.544	Acyl chain
14.073	Terminal
13.768	methylene.

Table 4
¹³C NMR Spectral data for CAPE 32 GC analysis


The fatty acid composition was determined by GC and the results are given below, the acid value and it was found that it is varied from 0.52-0.31. For the analysis the method followed for Gas chromatography (GC) is as

follows:GC Apparatus:Agilent Technologies, USA., Model no →6890 N, Column → DB-225 [(50% cyanopropylphenyl)-dimethylpolysiloxane, 30M×0.250mm1D×0.25µm film thickness Inlet temp →

230^oc,Oven prog → 160^oc(2min)-5^oc/min-230^oc(20min) , Det. temp → 270^oc ,HY → 30ml/min ,Air → 300ml/min ,Inlet type → Split/Split less , Det. type → FID ,Soft ware → GC Chem. station , A.10.02(1757) The fatty acid composition was

determined by GC and the results are given below; it was found that the acid value varied from 0.52-0.31.Fatty acid composition of the Bhallataka oil (CAPE 32).

Figure 11
GC analysis of *Semecarpus anacardium* L.f Oil



rted By : Retention Time
 Multiplier : 1.0000
 Dilution : 1.0000
 Sample Amount : 1.00000 [ng/ul] (not used in calc.)
 Use Multiplier & Dilution Factor with ISTDs

Signal 1: FID1 A,

Peak #	RetTime [min]	Sig	Type	Area [pA*s]	Height [pA]	Area %
1	4.512	1	BB	25.64367	6.22549	0.18675
2	6.651	1	BP	35.49586	13.30136	0.25849
3	9.409	1	BB	1842.54785	480.79678	13.41810
4	9.603	1	PB	20.24084	7.05006	0.14740
5	12.381	1	BV	1326.60620	192.38126	9.66082
6	12.701	1	VB	7038.10791	1063.49280	51.25403
7	13.156	1	VV	774.37714	228.03784	5.63929
8	13.320	1	VB	730.65979	212.54608	5.32093
9	13.844	1	BP	99.02870	30.61722	0.72116
10	15.178	1	BP	524.48413	145.53618	3.81948

Sample file: C:\APPLICHEM\DATA\MISC\ANP00008.D

Sample Name:

Peak #	RetTime [min]	Sig	Type	Area [pA*s]	Height [pA]	Area %
11	15.439	1	BB	150.90208	43.02037	1.09892
12	15.811	1	BP	163.37068	40.40532	1.18972
13	18.240	1	BV	697.12018	135.39716	5.07668
14	20.356	1	PB	141.16777	18.89869	1.02803
15	22.573	1	PB	162.06049	21.10181	1.18018

Totals : 1.37318e4 2638.80843

Results obtained with enhanced integrator!

*** End of Report ***

Table 5
Fatty acid composition of CAPE-32

Fatty acid	12:0	14:0	16:0	16:1	18:1	18:2	18:3	20:0	20:1	20:2	22:0	23:0	24:0
Wt%	0.18675	0.25849	13.41810	0.14740	51.25403	5.63929	6.04209	3.81948	1.09892	1.18972	5.07668	1.02803	1.18018

Antimicrobial activity

Determination of Antimicrobial activity

The antimicrobial activity of the isolated compounds and their derivatives was determined by using well diffusion method (Amsterdam, 1996) against different pathogenic reference strains procured from the Microbial Type Culture Collection and Gene Bank (MTCC), CSIR-Institute of Microbial Technology, Chandigarh, India. The pathogenic bacteria and *Candida* reference strains were seeded on the surface of the media Petri plates, containing Muller-Hinton agar with 0.1 ml of previously prepared microbial suspensions individually containing 1.5×10^8 cfu ml⁻¹ (equal to 0.5 McFarland). Wells of 6.0

mm diameter were prepared in the media Petri plates using a cork borer and the isolated compound and their derivatives at a dose range of 300 - 1.4 µg well⁻¹ was added in each well under sterile conditions in a laminar air flow chamber. Standard antibiotic solutions of Neomycin and Miconazole at a dose range of 300 -1.4 µg well⁻¹ and the well containing methanol served as positive and negative controls, respectively. The plates were incubated for 24 h at 30°C and the well containing the least concentration showing the inhibition zone was considered as the minimum inhibitory concentration. All experiments were carried out in duplicates and mean values are represented.

Table 6
Antimicrobial activity of Bhallataka oil

COMPOUNDS	<i>Staphylococcus aureus</i> MTCC 96	<i>Klebsiella planticola</i> MTCC 530	<i>Bacillus subtilis</i> MTCC 121	<i>S.aureus</i> MLS16 MTCC 2940	<i>Micrococcus luteus</i> MTCC 2470	<i>Escherechia coli</i> MTCC 739	<i>Pseudomonas aeruginosa</i> MTCC 2453	<i>Candida albicans</i> MTCC 3017
BO	---	150	75	75	---	---	150	150
FAME BO	9.37	9.37	18.75	9.37	9.37	9.37	9.37	9.37
Neomycin	18.75	18.75	18.75	18.75	18.75	18.75	18.75	--
Miconazole	--	--	--	--	--	--	--	9.37




Figure 12
Graphical representation Antimicrobial activity of Bhallataka oil

The seed oil of *Semecarpus anacardium* (Bhallataka oil) has number of bioactive compounds. In our investigation, Oil was characterized by FT-IR, ¹H NMR, ¹³C NMR, GC methods; it was found that it contains free fatty acids, triglycerides and others. *Semecarpus anacardium* nut oil cape 32 had found to have a free fatty acid (0.50) hence; alkaline Tran's-esterification was done by using anhydrous methanol at a molar ratio of 6:1 and 3g/Liter of sodium hydroxide as catalyst. SA Nut oil to yield Fatty Acid Methyl ester, and glycerine as a by-product and the Derivative (FAME) determined by Trans-esterification process (Fig.7) compared with the Various Oil Spectral data & Various properties, which were not reported earlier¹⁵⁻²⁴, Bhallataka oil derivative is

most active against both gram +ve and gram -ve bacteria when compared to Bhallataka oil (Table.6). It was found that it contains free fatty acids, triglycerides and others. Basing on spectral data and their physical properties *Semecarpus anacardium* L.f. nut oil was selected for biodiesel production.

CONCLUSION

Bhallataka oil has a specific free fatty acid value of 0.50. Hence, alkaline Tran's esterification process was chosen. These results were studied by Gas Chromatography and found the methyl esters of the following acids as : Oleic acid(18:1)-51.25403%;

Palmitic acid(16:0) -13.41810%; α -Linoleic acid(18:3)-6.04209%;Linoleic acid(18:2) -5.63929%;Docosanoic acid(22:0) -5.07668; Eicosanoic acid(20:0) -3.81948;Eicosadienoic acid(20:2)-1.18972;Lignoceric acid(24:0)-1.18018;Eicosenoic acid(20:1)-1.09892; Mead acid(23:0)-1.02803;Myristic acid(14:0)-0.25849;Lauric acid(12:0)-0.18675;Palmitoleic acid(16:1)-0.14740%.These results showed, that all the above fatty acids were totally converted to their corresponding methyl esters. Further we found that, the Bhallataka oil derivative is most active against gram +ve and gram -ve bacteria when compared to Bhallataka oil. The applications of these various compositions of

Bhallataka Oil as Biodiesel and Biolubricants are under process.

ACKNOWLEDGMENT

We would like to thank Sri Dr.S.Chandrasekhar, Director of IICT-CCMB, Sri Dr. R.B.N. Prasad Chief Scientist Department of Lipid Research, CSIR-IICT, Hyderabad, Dr. M.S.L Karuna, A. Tirupathi, for their kind cooperation and help during the tenure of this research.

CONFLICT OF INTEREST

Conflict of interest declared none.

REFERENCES

1. Maheshwari JK, *Addition to the Flora of Delhi*, Department of Botany, University of Delh 1996: 48.
2. MurthySSN,A biflavanoid Semcarpuflavanone , from *Semecarpus anacardium*.Phytochemistry.1983;;22:518-520.
3. Ramprasath V.R., Shanthi P. and Sachdanandam P., Anti-inflammatory effect of *Semecarpus anacardium* Linn.nut extract in acute and chronic inflammatory conditions, Biol Pharm Bull. 2004; 27(12): 2028-2031.
4. Vijayalakshmi T., Muthulakshmi V. and Sachdanandam P., Effect of the milk extract of *Semecarpus anacardium* nut on adjuvant arthritis - A dose dependent study in Wistar albino rats, Genetical Pharmacology.1996; 27(7): 1223-1226 .
5. Kothai R., Arul B., Suresh Kumar K. and Christina A.J.M., Hypoglycemic and antihyperglycemic effects of *Semecarpus anacardium* Linn. in normal and Alloxan included diabetic rats.Journal of Pharmacotherapy. 2005;5(2):4956.
6. Premalatha B. and Sachdanandam P.,Effect of *Semecarpus anacardium* nut extract against aflatoxin B -induced hepatocellular carcinoma, Fitoterapia.,1999;;70(5): 484-492 .
7. Narayan J.P., John M.S., Ghosh P.K.,Singh J.N., Jha O.P. and Jha I.S., Screening of some medicinal plants for spermatostatic and spermicidal properties,Proceedings of symposium on phytochemistry and Botanical Classification, CBS publishers and Distributors Pvt. Ltd.,Delhi, India. 1985;;211-216 .
8. Nagabhushanaa K.S., Umamaheshwari S., Tocolic F.E., Prabhu S.K., Green I.R. and Ramadoss C.S., Inhibition of soybean and potato Lipoxygenases by Bhilawanols from Bhilawan (*Semecarpus anacardium*) nut shell liquid and some synthetic Salicylic acid Analogues ,Journal Enzyme Inhib Medl Chem. 2002; 17(4):255-259.
9. Tripathi Y.B. and Pandey R.S., *Semecarpus anacardium* L, nuts inhibit lipopolysaccharide induced NO production in rat macrophages along with its hypolipidemic property, Indian JExp Biol.2004;;42(4): 432-436.
10. Nair A. and Bhide S.V., Antimicrobial properties of different parts of *Semecarpus anacardium*, Indian Drugs.1996;33: 323-328.
11. Shukla S.D., Jain S., Sharma K. and Bhatnagar M., Stress induced neuron degeneration and Protective effects of *Semecarpus anacardium* Linn. And *Withania somnifera* Dunn. In Hippocampus of albino rats: Anultrastructural study. Indian J Exp Biol. 2000; 38(10):1007-13.
12. Premalatha B. and Sachdandam P., Immunomodulatory Activity of *Semecarpus anacardium* Linn. Nut Milk Extract in Aflatoxin B – induced Hepatocellular: Carcinoma in Rats. Pharm. Phmacol. Commun., 1998;;4: 507-510.
13. Yoko Miyake,Kazuhisa Yokomizo Determination of Unsaturated Fatty Acids Composition By High-Resolution Nuclear Magnetic Resonance Spectroscopy, JAOCS.,1998; 75(9):1091- 1093.
14. Christie WW.Sampling and analysis of commercial fats and oils. AOCS Official Method Cd3d-63, Acid value 20036 W. Christie, in Advances inLipid.Methodology-Two.1993; 69-111.
15. Jagan Mohan R.S¹, Padmavathi V¹, Kesava R. B^{1*} and Noboru Motohashi^{**},Cardenolides and Relates of Mainly *Calotropis gigantea* and *C.Procera* in the Family *Asclepiadaceae*” Text Book on Occurrences, Structure ,Biosynthesis, and Health Benefits Based on their Evidences of Medicinal Phytochemicals in Vegetables and Fruits,2015; 4(4):109-180.
16. Padmavathi V¹, Kesava R. B^{1*}, Noboru Motohashi², Sridhara Janardhan³ Narahari Sastry G³, “Comparative and Computer Assisted Drug Designing of Fatty Acids Isolated From Flowers, Leaves, Stem bark, Root bark and Nuts of *Semecarpus anacardium* L.f (Anacardiaceae).” Journal of Pharmacy and Pharmacology.,2014; 2, 582-591,USA Print ISSN: 2328-2150.
17. Padmavathi V, Kesava Rao B*,”Heterocyclic Compounds and their Biological Applications of *Semecarpus anacardium* L.f, Special Chapter in the Text Book on Hetero cyclic compounds ,2017(In press).
18. Padmavathi.V & KesavaRao.B*,Proceedings on FTIR Studies in Drug Designing and Discovering

- the potent molecules as drugs from *Semecarpus anacardium* L.f, [DDD-2013] ,Kolhapur, ISBN: 978-93-5126-349-4, RA-1 (pp 14-(17).2013, which has got best oral paper presentation award in this conference.
19. Padmavathi .V, Kesava R.B*. Naturally Occurring Biflavonoids from *Semecarpus anacardium* L.f and Their Biological Activity-A Review, Proceedings of Andhra Pradesh Academy of Sciences(PAPAS),2014;16(1)January- June:39-44.
 20. Padmavathi.V&Kesava Rao.B*.*Proceedings on New Dimentions in Chemistry and Chemical Technologies Applications in Pharma Industry.,2014; (NDCT-2014)June 23-25.ISBN 978-93-82829-90-4.(Pp-422- 427).*
 21. Padmavathi.V&Kesava Rao.B* *Proceedings on Chemical constituents From Methanolic extract of Semecarpus anacardium*L.f Flowering Buds, Antimicrobial Activity, AnticancerActivity And their Docking Studies *Indian youth Science congress, 2015; Jan 19-21, ANU Campus* which has got Young Scientist Award in this Conference.
 22. Padmavathi.V& Kesava Rao.B* , *Proceedings on "Trend Setting Innovations of Biflavonoids from Semecarpus anacardium* L.f of Docking studies and Biological activates in Pharma Industry", TSCST-JNTUH, October 16-18,2015; ISBN 978-93-82829-48-5,pp 384-388,
 23. Padmavathi .V and Kesava Rao.B*, Proceedings on "Physicochemical Properties of *Semecarpus Anacardium* L.f",Trend Setting Innovations in Chemical Sciences and Technology-Nature inspired Chemistry and Engineering,TSCST-NICE 2016;4-6 October, JNTUH,Hyderabad2016, OP-3, ISBN978-93-82829-14-0. (Pp-224-231).
 24. Padmavathi .V and Kesava Rao.B*, Correlation Between Metal Ions and Organic Compounds from *Semecarpus anacardium* L.f., Their Biological Aspects and Docking Studies, International Journal of Recent Scientific Research (IJRSR),December2016; 7(12):14952-14959.
 25. Kesava Rao.B* and Usha Rani.G,GC-MS Analysis of volatile components in petroleum ether extracts Of *Coldenia procumbens* Linn, Internal Journal of Pharma and Biosciences.2016;7(2):241-245.

Reviewers of this article



Dr. Shalini Jaiswal

Coordinator & HOD,
Department of Chemistry,
United College of Engineering and Research,
Noida, India

Dr C Suresh Reddy

Professor of Chemistry
S.V.U.College of Sciences
S.V.University
Tirupati-517502



Prof. Srawan Kumar G.Y

Associate Professor, Nalanda Institute of
Pharmaceutical Sciences, Sattenapalli,
Guntur, Andrapradesh, India



Prof. Dr. K. Suriaprabha

Asst. Editor, International Journal
of Pharma and Bio sciences.



Prof. P. Muthuprasanna

Managing Editor, International
Journal of Pharma and Bio sciences.

We sincerely thank the above reviewers for peer reviewing the manuscript