



## ISOLATION AND CHARACTERIZATION OF PHYTOSTEROID AND TRITERPENES FROM LEAVES OF ABUTILON CRISPUM

**DR.M.RAM MOHAN<sup>1\*</sup>, S.GANAPATY<sup>2</sup>, B.VASUDHA<sup>1</sup>**

<sup>1\*</sup> Associate professor, Department of Pharmacognosy and Phytochemistry,  
Anurag School of Pharmacy, Hyderabad, Telangana, India

<sup>2</sup> GITAM Institute of Pharmacy, GITAM University, Gandhi Nagar,  
Visakhapatnam-530 045, Andhra Pradesh, India.

### ABSTRACT

Natural products have served as an important source of drugs since ancient times and about half of the useful drugs today are derived from natural sources. The study of natural products has an advantage over synthetic drugs. Most of the phytochemicals derived from natural sources are extremely useful as lead structures for synthetic modifications and optimization of bioactivity. The aim of the study is to identify and characterize the bioactive compounds from leaves of *Abutilon crispum*. For isolation of the compounds, the dried leaves powder of *Abutilon crispum* was subjected to extraction with chloroform and subjected to column chromatography with different solvent fractions. Three compounds were isolated from the chloroform extract and purified. These compounds were eluted with benzene and petroleum ether. The structures of these compounds were characterized by using IR, NMR and Mass spectrometry. These compounds were isolated from this plant for the first time.

**KEYWORDS:** *Abutilon crispum*, Isolation, IR, NMR.



**DR.M.RAM MOHAN \***

Associate professor, Department of Pharmacognosy and Phytochemistry,  
Anurag School of Pharmacy, Hyderabad, Telangana, India.

Received on : 27-01-2017

Revised and Accepted on : 14-06-2017

DOI: <http://dx.doi.org/10.22376/ijpbs.8.3.p190-195>

## INTRODUCTON

*Abutilon crispum* belonging to family Malvaceae is a trailing perennial, weak, shrub. This plant is commonly distributed in the shady undergrowth forest on hilly slopes found throughout India. It is known as Errabenda in Telugu<sup>1</sup>. The plant finds its application in the traditional system of medicine. In India, this plant is used in the treatment of Asthma in Rayalaseema Region of Andhra Pradesh and it is also used for the treatment of Piles by the Kondadora Tribe of Northern Andhra Pradesh<sup>2</sup>. Its leaves are used for treating asthma, piles, ulcers, cough, jaundice and diabetics by tribal people of Andhra Pradesh and its fruits are used in the treatment of piles in Tamilnadu. The plant is used as a leafy vegetable by the Paliyar tribes of Pachalur in Dindigul district of Tamilnadu<sup>3-6</sup>.

## MATERIALS AND METHODS

### Plant Collection and Preparation

The fresh leaves of *Abutilon crispum* collected from Pakala (Warangal) in the month of August were subjected to be authenticated by Prof. V. S. Raju, Department of Botany, Kakatiya University and Warangal, voucher specimen (MRM/03/2012) was deposited in the College of Pharmaceutical Sciences, Andhra University, Visakhapatnam. After confirmed authenticity and leaves were manually separated. The plant material was washed with water to remove soil, mud, debris and other adhering materials and dried thoroughly in air under shade at room temperature. Coarse prepared powdered drug was passed through sieve no.40 and stored in air tight containers.

### Extraction and Isolation

About (1kg) of powdered leaves of *Abutilon crispum* was extracted with chloroform using Soxhlet apparatus. The extract was concentrated with rotary evaporator. the chloroform extract was stored in screw cap vial at 4°C until further use. The chloroform extract upon concentration under reduced pressure left a dark gummy residue (15g). The extract showed positive Liebermann-Burchard reaction (pink-Blue-Green) and negative ferric chloride, Shinoda tests, indicates the presents of steroidal compounds. The 15g of chloroform extract was admixed with 15g silicagel (60/120 meshes) to get uniform mixing. 150 g of silica gel (70/325 meshes). The column was kept aside for 1 hr and allowed for close packing. Admixture was then added at the top of the stationary phase and separating of the compounds as started by eluting with various solvent such as petroleum ether, benzene and petroleum ether with different ratios with increasing of polarity. All the column fractions were collected separately and concentrated.

### Spectroscopic characterization


Different spectroscopic methods were used to elucidate the structure of compounds. Among the spectroscopic techniques IR, <sup>1</sup>H Nuclear Magnetic Resonance and <sup>13</sup>C NMR as carried out. The infra red spectrum as recorded on Perkin Elmer FTIR88, <sup>1</sup>HNMR spectra were recorded on a 90MHz, Jeol JNM EX-90 FT NMR, 300 MHz Bruker FT DRX-300. The <sup>1</sup>HNMR was recorded using CDCl<sub>3</sub>, as solvent with Tetramethyl silane (TMS) as an internal standard.

## RESULTS AND DISCUSSION


Repeated chromatographic separation and purification of the chloroform extract of the leaves of *Abutilon crispum* provided a total of three compounds (compound 1, 2, and 3). The structures of which were determined by extensive IR, <sup>1</sup>HNMR and <sup>13</sup>CNMR spectral analysis as well as by comparison of their spectral data with previously reported values.

### Characterization of compound -1


It crystallized as needles from petroleum ether, m.p. 136-138°C. It showed colour reactions in L.B. test, characteristic of sterols (play of colours) and gave a single spot on silver nitrate impregnated TLC. The IR absorption spectrum showed absorption peaks at 3373cm<sup>-1</sup>(O-H stretching.); 2940cm<sup>-1</sup>and 2867cm<sup>-1</sup> (aliphatic C-H stretching); 1641cm<sup>-1</sup> (C=C absorption peak); other absorption peaks includes 1457cm<sup>-1</sup> (CH<sub>2</sub>); 1381cm<sup>-1</sup> (OH def), 1038 cm<sup>-1</sup>(cycloalkane) and 881.6 cm<sup>-1</sup>. The <sup>1</sup>HNMR spectrum of the compound (1) (Fig-1) has revealed a one proton multiplet at δ 3.51, the position and multiplicity of which was indicative of H-3 of the steroid nucleus. The typical H-6 of the steroidal skeleton appeared as a multiplet at δ 5.39 for one proton. The resonance signals at δ 0.72 and δ 1.05 (3H each) was due to two tertiary methyl groups at C-13 and C-10 respectively. Two doublets centered at δ 0.87 with coupling constant 6.7 Hz and δ 0.85 with coupling constant 6.7 Hz were assigned for two methyl groups at C-25. The <sup>1</sup>HNMR spectrum showed a doublet at δ 0.96 with coupling constant 6.5 Hz for a methyl group at C-20. The spectrum also showed three protons triplet at δ 0.89 which could be assigned to the primary methyl group at C-28 and <sup>13</sup>CNMR has given signal at 150.98, 145.2 (C-5), 139.8 (C-22), 121.7, 118.89(C-6), 79.03 (C-3), 55.3(C-14), 55.18(C-17), 50.45 (C-9), 48.3 (C-9), 40.8 (C-20), 40.1(C-12), 39.2 (C-13), 38.9 (C-4), 38.6 (C-12), 37.18 (C-1), 37.12 (C-10), 36.3 (C-8), 35.59(C-20), 34.29 (C-22), 34.24 (C-7), 32.66 (C-8), 29.86 (C-25), 29.71 (C-16), 28.41 (C-2), 28.1 (C-15), 27.4 (C-28), 26.1 (C-11,26), 21.6 (C-27), 19.32 (C-19), 17.71 (C-21), 15.6 (C-18, 29). On this basis, compound-1 was characterized as β-Sitosterol, the identity of which was confirmed by comparison of the spectral data with previously reported values<sup>7</sup>




**Figure1**  
*Chemical structure of  $\beta$ -Sitosterol*



**Figure 1.1**  
*<sup>1</sup>H NMR of  $\beta$ -Sitosterol*



**Figure 2**  
*Chemical structure of Lupeol*



**Figure 2.1**  
**<sup>1</sup>H NMR of Lupeol**

### **Characterization of compound -2**

It crystallized from petroleum ether-Benzene as needles, m.p. 212-213°C, It gave pink colour in L.B. reaction and yellow colour with tetranitromethane. In TLC, it corresponded to Lupeol and the identity was confirmed by comparison with an authentic sample (mmp and Co-TLC). The compound showed IR peaks corresponding to hydroxyl (3410, 1042 cm<sup>-1</sup>) and vinylidene group (1640 and 881 cm<sup>-1</sup>), 2940, 2870, 1640 (>=CH<sub>2</sub>), 1453, 1380 (-C-Me<sub>2</sub>), 1042 (OH), 881 (>=CH<sub>2</sub>). The <sup>1</sup>H NMR spectrum of the compound (2) (Fig-2) that revealed signals at  $\delta$  4.56 appeared as one proton doublet with coupling constant 0.4 Hz and signals at  $\delta$  4.67, as one proton doublet with coupling constants 0.4 Hz and 0.5 Hz centered at for an exomethylene group. The signals at  $\delta$  3.18 appeared as double doublets with coupling constant 9.6 Hz and 6.2 Hz centered at could be assigned for a secondary carbinol group. The spectrum also showed a broad doublet at  $\delta$  1.66 with coupling constant 0.5 Hz attributed to a methine group. The resonance signals at  $\delta$  0.75, 0.77, 0.80, 0.92, 0.94 and 1.02 exhibited six tertiary methyl singlets and <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz)  $\delta$ : 38.77 (C - 1), 27.46 (C - 2), 79.02 (C - 3), 38.87 (C - 4), 55.31 (C - 5), 18.33 (C - 6), 34.42 (C - 7), 40.84 (C - 8), 50.45 (C - 9), 37.18 (C - 10), 20.94 (C - 11), 25.15 (C - 12), 38.07 (C - 13), 42.84 (C - 14), 27.46 (C - 15), 35.59 (C - 16), 43.01 (C - 17), 48.72 (C - 18), 48.32 (C - 19), 150.99 (C - 20), 29.86 (C - 21), 40.01 (C - 22), 27.99 (C - 23), 15.37 (C - 24), 16.12 (C - 25), 15.98 (C - 26), 14.55 (C - 27), 18.01 (C - 28), 109.32 (C - 29), 19.31 (C - 30). These data were in close agreement with those reported for a

typical pentacyclic triterpenoid Lupeol and further confirmed the identity of compound- 2 as Lupeol<sup>8</sup>.

### **Characterization of compound -3**

It crystallized from acetone as colourless plates, m.p. 199-200°C and showed colour reactions characteristics for triterpenes. In TLC, it corresponded to  $\beta$ -Amyrin and the identity was confirmed by comparison with an authentic sample (mmp and Co-TLC). The IR absorption spectrum showed absorption peaks at 3373cm<sup>-1</sup> (O-H stretching.); 2940 cm<sup>-1</sup> and 2867cm<sup>-1</sup> (aliphatic C-H stretching); 1641 cm<sup>-1</sup> (C=C absorption peak); other absorption peaks includes 1457cm<sup>-1</sup> (CH<sub>2</sub>); 1381 cm<sup>-1</sup> (OH def), 1038.7cm<sup>-1</sup> (cycloalkane) and 881 cm<sup>-1</sup>. The <sup>1</sup>H NMR spectral data showed the presence of a downfield methine proton at  $\delta$  3.3 (1H, m) characteristic of H-18 of oleanene derivatives, an olefinic proton signal at  $\delta$  5.4 (br, s, H-12). The spectra also <sup>1</sup>H NMR showed the presence of eight tertiary methyls in the region  $\delta$  1.25-0.78 indicative of oleanene skeleton in compound and <sup>13</sup>C NMR  $\delta$  39.59 (C-1), 27.93 (C-2), 80.8 (C-3), 39.59 (C-4), 55.24 (C-5), 18.19 (C-6), 33.73 (C-7), 38.37 (C-8), 47.4 (C-9), 35.55 (C-10), 23.59 (C-11), 121.62 (C-12), 145.20 (C-13), 42.05 (C-14), 28.73 (C-15), 27.93 (C-16), 32.85 (C-17), 59.0 (C-18), 40.01 (C-19), 41.52 (C-20), 31.23 (C-21), 42.01 (C-22), 29.69 (C-23), 15.72 (C-24), 15.72 (C-25), 16.84 (C-26), 23.59 (C-27), 28.73 (C-28), 17.50 (C-29), 21.39 (C-30). These data were in close agreement with those reported for a typical pentacyclic triterpenoid and further confirmed the identity of compound- 3 as  $\beta$ - Amyrin.- 3.




Figure 3  
Chemical structure of  $\beta$ - Amyrin




Figure 3.1  
 $^1\text{H}$  NMR of  $\beta$ - Amyrin

## CONCLUSION

Chemical examination of the leaves of *Abutilon crispum* on conventional extraction and a sequential chromatography afforded three compounds. From the chloroform extract three compounds were separated and identified as  $\beta$ -Sitosterol, Lupeol and  $\beta$ -Amyrin. All the compounds were characterized by physical, chemical and spectral studies using IR,  $^1\text{H}$  NMR, and  $^{13}\text{C}$  NMR. This is the first report of occurrence of this compound in this plant. Therefore future biological investigations are needed for these isolated compounds.

## REFERENCES

- RamMohan M, Srinivas K, Ganapaty, S. Hepatoprotective activity of the leaves of *Abutilon crispum*, World J Pharma Pharmaceutical Sci. 2014 Nov; 3(11): 774-9.
- Sekhar PC, Kumar YV, Grace JR, Murty PP. Some Ethno Medicinal Plants used for the treatment of piles by the kondadora tribe of Northern Andhra Pradesh, AP, India. Int J Ayur and Herbal Medicine. 2012 Oct; 2(5): 803-9.
- Anjaneyulu E, Sudarsanam G. Folk medicinal plants used in the treatment of Asthma in Rayalaseema region of Andhra Pradesh, India, India Res J Pharm, Bio and Chem Sci. 2013 Mar; 4(1): 833-9.
- Padal SB, Ramakrishna H, Devender R. Ethnomedicinal studies for endemic diseases by the tribes of Munchingiputtu Mandal, Visakhapatnam district, Andhra Pradesh, India.

## ACKNOWLEDGMENTS

The authors are thankful to Gitam University, Vishakapatnam and Andrapradesh for providing necessary facilities for carrying out the research work.

## CONFLICT OF INTEREST

Conflict of interest declared none.

- Int J Med and Aromatic Plants.2012 Sep; 2(3): 453-9
5. Shanmugam S, Annadurai M, Rajendran K. Ethnomedicinal plants used to cure diarrhoea and dysentery in Pachalar hills of Dindigul district in Tamil Nadu, Southern India. Journal of Applied Pharmaceutical Science. 2011 Oct 1;1(8):94.
  6. Ram Mohan M, Ganapaty S. Anti-Inflammatory and Toxicity Studies of the Leaves of *Abutilon crispum*, Int J Pharmaceutical Res Sch.2015 Jan; 4(1):108-11.
  7. Jain PS, Bari SB. Isolation of Stigmasterol and beta Sitosterol from petroleum ether extract of woody stem of *Abeslmsochus manhot*, Asian J of Plant Sci.2009 Nov; 4(2):112-7.
  8. Iman S, Azar I, Hasan MM, Ali MS, Ahmed SW. Two triterpenoids lupanone and lupeol isolation and identification from *Tamarindus indica*, Pak.J. Pharm, Sci.2007; 20:125-7.
  9. Virgilio D, Ebajo JR, Chien-Chang Shen, Consolacion Y, Terpenoids and Steroids from *Hoya multiflora*,J of App Pharmaceutical Sci.2015 Nov;5 (4):33-7.
  10. Patel AJ. Isolation and Characterizat on of Lupeol from *Echinops echinatus*,Euro J Pharmaceutical and Medical Res.2016 Jul;3(9) : 385-7.
  11. Shumaia Parvin Md, Abdul Kader Md, Abdul Muhit Md,Ekramul Haque. Terpenoids and Phytosterols from stem bark of *Crataeva nurvala*, J of App Pharmaceutical Sci.2011 Nov; 1 (9): 47-50.
  12. Jain PS, Bari SB. Isolation of lupeol, stigmasterol and campesterol from petroleum ether extract of woody stem of *Wrightia tinctoria*. Asian Journal of Plant Sciences. 2010;9(3):163.
  13. Rajeswari J, Rani S. Isolation, structural elucidation of flavonoid constituents from *Lawsonia inermis*,Scholar Res Library.2016; 8(7): 203-8.
  14. Anjoo kamboj, Ajay kumar saluja. Isolation of Stigmasteol and  $\beta$ -Sitosterol from petroleum ether extract of aerial parts of *Ageratum conyzods*, Int J of Pharmacy and Pharmaceutical Sci.2011 Oct; 3(1): 94-6.
  15. Saratha V, Iyyam Pillai S, Subramanian S. Isolation and characterization of lupeol, a triterpenoid from *Calotropis gigantea* latex. International Journal of Pharmaceutical Sciences Review and Research. 2011;10(2):54-6.
  16. Snehal B. Bhandare, Kirtis Ladda. Isolation of Kaemperol and Quercetin from *Podophyllum hexandrum* rhizome. Int J Pharma Bio Sci, 2016 Apr;7(2):98-100.

## Reviewers of this article

**Dr. Karka Srinivas Reddy**

Associate Professor, Pharmacognosy and  
Phytochemistry  
Vaagdevi college of Pharmacy  
Ramnagar,hanamkonda  
Warangal Urban



**Prof. Dr. K. Suriaprabha**

Asst. Editor , International Journal  
of Pharma and Bio sciences.



**Asst. Prof. Dr. Sujata Bhattacharya**

Assistant Professor, School of Biological  
and Environmental Sciences, Shoolini  
University, Solan (HP)-173212, India



**Prof. P. Muthuprasanna**

Managing Editor , International  
Journal of Pharma and Bio sciences.

We sincerely thank the above reviewers for peer reviewing the manuscript