



CO-RELATION BETWEEN HEAVY METAL RESISTANCE AND ANTIBIOTIC SUSCEPTIBILITY IN HALOTOLERANT BACTERIA ISOLATED FROM THE ANTARCTIC OCEANIC REGION

ANURADHA R MULIK AND RAMA K BHADEKAR*

Department of Microbial Biotechnology, Rajiv Gandhi Institute of IT and Biotechnology, Bharati Vidyapeeth Deemed University, Katraj, Pune-411046, Maharashtra, India

ABSTRACT

Four bacterial isolates viz *Kocuriasp BRI 36*, *Brevibacillus sp BRI 37*, *Halomonas sp BRI 38* and *Oceanobacillus sp BRI39* from Antarctic oceanic region exhibiting high tolerance to heavy metals like cadmium, chromium, nickel and lead were selected for this study. Their maximum tolerable concentration MTC values were found to be in the range of 300 to 600 ppm. Interestingly, the isolates exhibited growth in presence of multiple metals at various temperatures (15, 30 and 45°C). The characteristic of metal resistance is known to be linked to antibiotic resistance and mostly associated with plasmid DNA. In order to determine role of plasmid DNA, curing experiments were performed using ethidium bromide and Sodium dodecyl sulfate SDS. Cured BRI isolates were again examined for metal tolerance and antibiotic susceptibility pattern. MTC values for the selected four metals were almost similar to those prior to curing. However, antibiotic susceptibility experiments revealed increase in sensitivity to piperacillin and chloramphenicol in cured BRI 36 whereas, remaining isolates demonstrated similar response to the antibiotics tested after curing. The observations indicated possible role of plasmid in piperacillin and chloramphenicol resistance in BRI 36 whereas, in other isolates it may or may not be plasmid linked. However, metal resistance characteristics of BRI isolates appear to be independent of extra chromosomal DNA, since the MTC results remained unaltered after curing.

KEYWORDS: *Multi-metal, Maximum tolerable concentration (MTC), antibiotic resistance, plasmid isolation, curing*



RAMA K BHADEKAR*

Department of Microbial Biotechnology, Rajiv Gandhi Institute of IT and Biotechnology,
Bharati Vidyapeeth Deemed University, Katraj, Pune-411046, Maharashtra, India

Received on: 25-03-2017

Revised and Accepted on: 20-09-2017

DOI: <http://dx.doi.org/10.22376/ijpbs.2017.8.4.b302-306>



[Creative commons version 4.0](#)

INTRODUCTION

Metals like Na, K, Mg, Ca, V, Mn, Fe, Co etc. play a vital role in metabolic processes of living organisms while some like Ag, Cd, Pb, Sn, Au, Hg etc. are toxic¹. Element biotransformation and biogeochemical cycling, decomposition, bioweathering, and soil and sediment formation are some of the activities in which microorganisms are involved. However, increase in metal concentration beyond certain limit activates microbial resistance mechanism which enables them to survive under stress². Microorganisms exhibit various strategies to combat the metal stress. Various adaptation mechanisms include reduction of metals to less toxic form, sequestration, active efflux etc.³ These mechanisms may be attributed to the genetic elements present on chromosomes, transposons or plasmid. In majority of the cases, it had been shown to be associated with plasmid DNA. For example, cadmium resistance mechanism in *Enterobacter* sp and *Staphylococcus aureus* sp,^{4,5} *Bacillus* strain JMAK1⁶, *Halomonas BVR 1*⁷ etc. However, role of chromosomal DNA in metal resistance had been very rarely documented. Moreover, antibiotic resistance is known to be linked to metal resistance. The studies carried out on *Bacillus* sp. showed loss of antibiotic and heavy metal resistance in this organism after plasmid curing⁸. However, among halophilic and/or halotolerant microorganisms very few studies of such type have been published. Keeping this in mind we have selected four halotolerant isolates from Antarctic oceanic region with significant resistance to cadmium, nickel, lead and chromium⁹ to establish the co-relation between heavy metal tolerance, antibiotic resistance and possible role of plasmid DNA. Recently we have reported effect of metals on pigment production by *Kocuri* sp. BRI 36¹⁰.

MATERIALS AND METHODS

Organism

Kocuri sp BRI 36, *Brevibacillus* sp BRI 37, *Halomonas* sp BRI 38 and *Oceanobacillus* sp BRI 39 were used in this work. The isolates were grown in Mineral Salt Medium (MSM) at 25±2°C for 48 h with shaking at 120 rpm¹¹. They were further used for inoculation in all the experiments at 10% concentration.

Chemicals and Reagents

All chemicals used were of analytical grade. The media components were purchased from Hi Media Laboratories Pvt. Ltd. (Mumbai, India). The stock solutions of cadmium, nickel, lead and chromium (1000 ppm each) were purchased from Sigma-Aldrich.

Screening of BRI isolates for multi-metal resistance

All the four isolates were cultivated in MSM amended with combinations of different metals (each at 10 mg/l concentration) at 30°C with shaking at 120 rpm for 48 h. At the end of incubation the samples were used to determine total viable count (TVC).

Effect of temperature on growth in presence of multiple metals

All the four isolates were grown under different

Conditions of temperature (15, 30 and 45°C) in presence of various combinations of metals each at 10 mg/l concentration with shaking at 120 rpm for 48 h.

Plasmid isolation

BRI isolates were grown in Luria-Bertani (LB) broth at 120 rpm for 48 h at room temperature. The cell pellets were harvested by centrifugation at 10000 rpm for 10 min at 4°C. The plasmid DNA was isolated by small scale boiling lysis method¹². The samples were dissolved in Tris-EDTA (T₁₀E₁) buffer and analysed on 1% agarose gel.

Plasmid curing

BRI isolates were grown in sterile LB amended with ethidium bromide (100 mcg/ml) and 2% SDS¹³. After incubation at 30°C for 48 h, the cultures were maintained on Luria agar plates. The cured BRI isolates were examined for i) presence of plasmid DNA ii) their MTC for Cd²⁺, Ni²⁺, Pb²⁺ and Cr³⁺ individually and iii) antibiotic resistance.

Maximum tolerable concentration (MTC)

Cured BRI isolates 36, 37, 38 and 39 were used to determine their maximum tolerable concentration (MTC) to Cd²⁺, Ni²⁺, Pb²⁺ and Cr³⁺ individually in the range of 200-700 ppm as described previously⁹.

Antibiotic resistance

The antibiotic susceptibility of cured BRI isolates was examined as described previously⁹ using 48 h grown cultures and antibiotic discs (Hi Media Pvt. Ltd. India).

RESULTS AND DISCUSSION

The four isolates viz. BRI 36, 37, 38 and 39 were examined to determine the multi-metal resistance and correlation between their metal resistance and antibiotic resistance characteristics. As reported previously, their MTC values for Cd²⁺, Cr³⁺, Ni²⁺ and Pb²⁺ were in the range of 300 to 600 ppm. Moreover, the isolates showed high susceptibility to most of the antibiotics tested⁹. The results of multi-metal resistance experiment showed that all the BRI isolates have ability to resist all four heavy metals in different combinations at 10 mg/l concentration of each metal. Previously studies on multimetal tolerance had been reported in *Thiobacillus ferrooxidans* by Das et al¹⁴ using Cu²⁺, Zn²⁺ and Fe³⁺. The authors had performed the experiments using the metals individually or in possible binary and ternary combinations. *Acinetobacter baumannii* strain HAF – 13 exhibited multi-metal resistance to Hg²⁺, Pb²⁺, Cd²⁺, As⁵⁺ and Cr⁶⁺ at their MTC (75-250 mg/l)¹⁵. The results of TVC experiments indicated maximum viability (23 %) of BRI 37 when exposed to all four metals followed by BRI 38 in presence of Ni²⁺, Pb²⁺, Cd²⁺ and BRI 36 in presence of Cr³⁺, Cd²⁺, Ni²⁺ (Figure 1) at 30°C. We studied the effect of temperature (Table 1) on multi-metal tolerance of all the isolates. All BRI cultures exhibited normal growth at temperatures of 15, 30 and 45°C indicating their ability to survive and proliferate in presence of multiple metals. Kumar¹⁶ had reported growth of *Pseudomonas aeruginosa* at different temperatures (25-45°) in presence of cadmium. He had observed maximum removal at

35°C. In order to confirm the association between these characteristics and plasmid DNA, plasmid curing experiments were carried out (Figure 2 (a, b)). Plasmid mediated resistance mechanism for cadmium, chromium and nickel in *E. coli* was studied by Lazar et al.¹⁷. Authors had discussed phenotypic data showing direct relation between multiple antibiotic resistance and heavy metal resistance. Cured BRI cultures (present work) were then reassessed for MTC values for Cd²⁺, Ni²⁺, Pb²⁺ and Cr³⁺ individually and also for antibiotic susceptibility. We observed MTC results almost similar to those obtained prior to curing⁹. However, antibiotic test results slightly differed from those prior to curing. Among the four BRI isolates, cured BRI 36 showed increased susceptibility to antibiotics in comparison to the previous sensitivity pattern⁹. Initially it had demonstrated resistance to piperacillin and chloramphenicol. However,

same concentration of antibiotics was found to inhibit its growth after curing. On the other hand BRI 37, 38 and 39 did not show any change in their antibiotic susceptibility pattern after curing. The results indicated plasmid mediated antibiotic resistance in halotolerant BRI 36 whereas it may or may not be plasmid derived in other BRI isolates. However, unaltered MTC values suggested absence of association with extra-chromosomal DNA. Thus, our results of heavy metal tolerance in BRI isolates showing susceptibility to majority of antibiotic tested indicate absence of direct correlation between these two characteristics. The present inference is currently under investigation. Similar results representing absence of correlation between antibiotic resistance and metal tolerance were observed by Boga et al.¹⁸ in Gram negative bacteria isolated from Lake Victoria wetlands.

Table 1
Effect of temperature on multi-metal tolerance

Sr.no	Temperature (°C)	15°C				30°C				45°C			
		BRI 36	BRI 37	BRI 38	BRI 39	BRI 36	BRI 37	BRI 38	BRI 39	BRI 36	BRI 37	BRI 38	BRI 39
1	Ni ²⁺ , Pb ²⁺ , Cr ³⁺ , Cd ²⁺	+	+	+	+	+	+	+	+	+	+	+	+
2	Ni ²⁺ , Pb ²⁺ , Cr ³⁺	+	+	+	+	+	+	+	+	+	+	+	+
3	Ni ²⁺ , Pb ²⁺ , Cd ²⁺	+	+	+	+	+	+	+	+	+	+	+	+
4	Pb ²⁺ , Cr ³⁺ , Cd ²⁺	+	+	+	+	+	+	+	+	+	+	+	+
5	Cr ³⁺ , Cd ²⁺ , Ni ²⁺	+	+	+	+	+	+	+	+	+	+	+	+




Figure 1
Percent survival of BRI isolates in presence of different combinations of metals.




Figure2 (a)




Figure2 (b)

Figure2

Agarose gel (1%) electrophoresis showing a) plasmid DNA from BRI isolates before curing and b) after curing. Lane 1- lambda DNA Hind III digest, lane 2 - BRI 36, lane 3- BRI 37, lane 4- BRI 38 and lane 5- BRI 39

CONCLUSION

Four bacterial isolates *Kocuria* sp BRI 36, *Brevibacillus* sp BRI 37, *Halomonas* sp BRI 38 and *Oceanobacillus* sp BRI 39 from the Antarctic oceanic region exhibiting high tolerance to heavy metals like cadmium, chromium, nickel and lead were selected for this study. They exhibited an ability to tolerate multiple heavy metals in different combinations. MTC values for the selected four metals were almost similar before and after curing of BRI isolates. However, antibiotic susceptibility experiments indicated possible role of plasmid in piperacillin and chloramphenicol resistance in BRI 36

whereas, in other isolates it may not be plasmid linked. However, metal resistance characteristics of BRI isolates appear to be independent of extra chromosomal DNA, since the MTC results remained unaltered after curing. Their antibiotic susceptibility test results and unaltered MTC values before and after curing suggest lack of co-relation between heavy metal tolerance and antibiotic resistance in BRI isolates.

CONFLICTS OF INTERESTS

Conflict of interest declared none.

REFERENCES

1. Rajendran P, Muthukrishnan J, Gunasekaran P. Microbes in heavy metal remediation. Indian J. Exp. Biol. 2003 41(9): 935-44.
2. Gadd GM. Metals, minerals and microbes: geomicrobiology and bioremediation. Microbiology. 2010 Mar 1;156(3):609-43..
3. Nies DH. Efflux-mediated heavy metal resistance in prokaryotes. FEMS Microbiol. 2003 27(2-3):313-39.
4. Tynecka Z, Gos Z, Zajac J. Reduced cadmium transport determined by a resistance plasmid in *Staphylococcus aureus*. J Bacteriol. 1998;147:305–12.
5. Bahig El-Deeb. Plasmid Mediated Tolerance and Removal of Heavy Metals by *Enterobacter* sp. Am. J. Biochem. Biotechnol. 2009; 5 (1): 47-53.
6. Guo S and Mahillon. J pGIAK1, a heavy metal resistant plasmid from an obligate alkaliphilic and halotolerant bacterium isolated from the Antarctic Concordia station Confined Environment. Plos one. 2003 8(8): e72461. <http://dx.doi.org/10.1371/journal.pone.0072461>.
7. Manasi, Rajesh.N, VidyaR. Evaluation of the genetic basis of heavy metal resistance in an isolate from electronic industry effluent. J Genet Eng Biotechnol. 2016 <http://dx.doi.org/10.1016/j.jgeb.2016.02.002>
8. Samanta. A, Bera. P, Khatun. M, Sinha. C, Pal. P, Lalee. A, Mandal. A. An investigation on heavy metal tolerance and antibiotic resistance properties of bacterial strain *Bacillus* sp. isolated from municipal waste. J Microbiol Biotech Res. 2012; 1(1):178-89.
9. Mulik AR and Bhadekar RK. Heavy metal removal by bacterial isolates from the Antarctic oceanic region. Int J Pharm Bio Sci. 2017 8: 535 -43.
10. Mulik. A, Kumbhar. P and Bhadekar. R. Production and partial characterization of pigments produced by *Kocuria* sp. BRI 36: influence of heavy metals. Int J Pharm Pharm Sci. 2017 9(10).
11. Pote. S., Chaudhary. Y., Upadhyay. S., Tale. V., Walujkar. S., Bhadekar. R. Identification and biotechnological potential of psychrotrophic marine isolates. Eurasia J Biosci. 2014 (8): 51-60.
12. Sambrook J, Fritsch EF, Maniatis T. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; 1989.
13. Durve A., Naphade S, Bhot M, Varghese J and Chandra N. Plasmid curing and protein profiling of heavy metal tolerating bacterial isolates. Arch Appl Sci Res. 2013; 5 (4):46-54.
14. Das, A., Modak J., Natarajan KA. Studies on multi-metal ion tolerance of *Thiobacillus ferrooxidans*. Miner Eng. 1997; 10 (7): 743-49.
15. Mohamed Helal El-Sayed. Multiple Heavy Metal and Antibiotic Resistance of *Acinetobacter baumannii* Strain HAF – 13 Isolated from Industrial Effluents. Am J Microbiol Res. 2016; 4(1): 26-36.
16. R.R Kumar. Isolation, Molecular identification of metal tolerant bacteria and its heavy metal removal capacity. International Journal of Microbiology, Biochemistry and Molecular Biology. 2014: <http://www.scienceresearchlibrary.com>.
17. Lazar V., Cernat R., Balotsen C., Cotar A., Coipan E., Cojocar C. Correlation between multiple antibiotic resistance and heavy metal tolerance among some *E. coli* strains isolated from polluted waters. Bacteriol. Virusol. Parazitol. Epidemiol. 200; 47: 155–60.
18. Boga. H. I., Okemo. P. O., Mwatha. et al. Heavy metal tolerance and antibiotic resistance profiles of gram negative bacteria isolated from Lake Victoria, Kenya. Journal of Tropical Microbiology and Biotechnology. 2007; 3 (2): 20-26.

Reviewers of this article



Dr Pradnya P. Kanekar , M. Sc., Ph.D. in Microbiology

CSIR Emeritus Scientist, Department of Biotechnology, Modern College of Arts, Science and Commerce, Shivajinagar, Pune 411005, Maharashtra.



Prof. Dr. K. Suriaprabha

Asst. Editor , International Journal of Pharma and Bio sciences.



Asst. Prof. Dr. Sujata Bhattacharya

Assistant Professor, School of Biological and Environmental Sciences, Shoolini University, Solan (HP)-173212, India



Prof. P. Muthuprasanna

Managing Editor , International Journal of Pharma and Bio sciences.

We sincerely thank the above reviewers for peer reviewing the manuscript