

IN VITRO FREE RADICAL SCAVENGING ACTIVITY OF VARIOUS ROOT AND RHIZOME EXTRACTS OF *ACORUS CALAMUS* LINN.**A.ELAYARAJA^{1*}, M.VIJAYALAKSHMI² AND G.DEVALARAO³**

¹. Department of Pharmacology, Sri Chandra Sekharendra Viswamaha Vidyalaya, Enathur, anchipuram. Tamil nadu state. India.

². Department of Microbiology, Acharya Nagarjuna University, Acharyanagar, Guntur, Andhra Pradesh state.

³. KVSRR Siddhartha college of Pharmaceutical Sciences, Vijayawada-520 010. Andhra Pradesh State.

*Corresponding author elayaraja80@rediffmail.com

ABSTRACT

Consecutive ethanolic and hydro alcoholic extracts of roots and rhizomes of *Acorus calamus* Linn (Araceae) were prepared and investigated for antioxidant potential against 2,2 diphenyl-1-picrylhydrazyl (DPPH) free radicals and compared with butylatedhydroxyanisole (BHA) and silymarin. Cumulative concentrations of both the extracts showed free radical scavenging activity of 59.13 ± 18.95 and 56.71 ± 19.54 respectively. It was also concluded that ethanolic extract showed the maximum antioxidant activity. Also increase in concentration of the extract showed enhance in free radical scavenging activity.

KEY WORDS

Acorus calamus Linn, Antioxidant activity, DPPH

INTRODUCTION

Acorus calamus Linn. (fam: Araceae) is a semi-aquatic, perennial, aromatic herb with creeping rhizomes originating in Asia¹ but now widely distributed in Europe, North America, Africa, etc. The rhizome, root and leaf yield a light brown to brownish yellow volatile aromatic oil known as calamus oil. The yield of oil from different part of the plant is upto 1.8% in fresh rhizome, 1.5 to 3.5% in dried rhizome, about 0.2% in leaves and 0.12% in fresh aerial parts². Bach (Hindi name of *Acorus calamus*) is an ingredient of several Ayurvedic recipes, which are used for the treatment of rheumatoid arthritis with pain, swelling and functional disability³⁻⁵ and

low-grade mentally retarded children⁶⁻⁷. Clinical trials of herbal preparations had also been carried out for schizophrenia containing brahmi, bachi and jyotismati⁸. The present study is intended to investigate the phytochemical, antibacterial and antifungal activity of various extracts obtained from the powdered roots and rhizomes of the plant. The present study revealed the first time of investigating the free radical scavenging activity of the roots and rhizomes part of Indian *Acorus calamus*.

MATERIALS AND METHODS

Collection of Plant Materials

The fresh plant materials of *Acorus calamus* which were collected from Krishna delta region of Vijayawada City surroundings of Krishna district in Andhra Pradesh state, was authenticated by Dr.S.M.Khasim, Assistant Professor, Department of Botany, Acharya Nagarjuna University in Guntur. One of the plant specimens had been planted in KVSR Siddhartha College of Pharmaceutical Sciences, Vijayawada and a voucher (No: KVSR/PCRL/No: 0051/BN) had been deposited after planted in the college herbal Garden.

Butylatedhydroxyanisole (BHA) and 2,2 diphenyl-1-picrylhydrazyl (DPPH) were purchased from B.D.H chemicals (P) Limited and S.D.Fine chemicals (P) Limited Mumbai. Silymarin was purchased from Sigma Aldrich, Mumbai. Other chemicals and reagents were procured from chemical laboratory of KVSR Siddhartha college of Pharmaceutical sciences, Vijayawada.

Extraction and Preliminary Phytochemical Screening

About 200 grams of the roots and rhizomes were subjected to hot soxhlete extraction⁹ apparatus with 700ml of pet ether (60-80°C) for 48 hrs. Then followed by 650 ml of ethanol (90%V/V) for another 48 hrs and the dried marc was subjected to cold maceration¹⁰ by using 600ml of hydroalcohol (1:1) for 72 hrs. Finally the obtained extracts were filtered through muslin cloth. Then they were concentrated under reduced pressure and dried in vacuum condition to get a semisolid consistency whose yields are tabulated. The dried extracts were subjected to various chemical tests to detect the presence of different phytoconstituents¹¹⁻¹⁵ present in them.

Experimental work¹⁶

The free radical scavenging activity of the *A.calamus* root and rhizomes extracts were measured and compared with the activity of Butylatedhydroxyanisole (BHA) and silymarin for radical-scavenging potency using the stable

radical 2,2 diphenyl-1-picrylhydrazyl (DPPH). The free radical scavenging activities of extracts, BHA and silymarin were measured by decrease in the absorbance of methanol solution of DPPH at 523 nm.

0.1mM solution of DPPH in spectrum alcohol was prepared and 1.5 ml of this solution was added to 3.5ml of extract solution, BHA and Silymarin in water at different concentrations (50-500mcg/ml). Thirty minutes later the absorbance was measured at 523nm. Lower absorbance of the reaction mixture indicates higher free radical scavenging activity. The capability of the DPPH radical was calculated using the following equation:

$$\text{DPPH Scavenged (\%)} = [(A_{\text{control}} - A_{\text{test}}) / A_{\text{control}}] \times 100$$

Where A_{control} is the absorbance of the control reaction and A_{test} is the absorbance in the presence of the sample of the extracts.

Statistical Analysis

All values are expressed as mean \pm SEM. The data were statistically analysed by one-way ANOVA followed by Turkey kramer multiple comparison test. P values < 0.05 was considered as significant. were considered significant.

RESULTS AND DISCUSSIONS

DPPH (2,2 diphenyl-1-picrylhydrazyl) is a stable free radical system, which is employed as an essential model by *in vitro* antioxidant evaluation. In general *in vitro* methods are preliminary screening methods, which paves way for the *in vivo* evaluation.

In the present study, free radical scavenging activity was carried on only two extracts which are polyphenolic enriched extracts. Ethanol extract (59.13 ± 18.95) and hydro alcoholic extract (56.71 ± 19.54) are showing an excellent scavenging effect and significant to Silymarin (63.55 ± 17.88). But the synthetic drug butylatedhydroxyanisole (BHA) has more activity.

When an antioxidant is mixed with any concentration of the free radical forming sample

such as DPPH, it reduces the free radical formation, which is detected by decrease in the absorbance of DPPH presence at λ_{max} 523nm. Both the extracts are significant to the natural compound Silymarin as showed in Table 1. But the free radical scavenging activity of herbal

standard was found to be less than the synthetic standard drug butylatedhydroxyanisole (BHA) at all dose levels. From this study it was concluded that free radical scavenging effect is tremendous for both the extracts at all the dose levels with that of the standard drugs.

Table1


DPPH radical scavenging activity (antioxidant activities) of root extract of *Acorus calamus* Linn and other standard drugs

S.No	Compound	DPPH radical scavenging activity (% Inhibition \pm SEM)
1.	Control	No activity
2.	BHA	70.37 \pm 14.39
3.	Silymarin	63.55 \pm 17.88
4.	EEAC	59.13 \pm 18.95
5.	HAEAC	56.71 \pm 19.54

BHA: Butylatedhydroxyanisole, EEAC: Ethanolic extract of *A. calamus* Linn.,
HAEAC: Hydro alcoholic extract of *A. calamus* Linn.

Figure 1

Bar Diagram Illustration of DPPH radical scavenging activity



ACKNOWLEDGEMENT

The authors are thankful to Siddhartha Academy for General and Technical Education,

Vijayawada for their valuable help in providing Laboratory facilities to carry out this research work.

REFERENCES

1. Directorate of Ministry of Health and Research Development. In: The Wealth of India. A dictionary of Indian raw materials and Industrial products. Council of Industrial Research, New Delhi.1985: 1; 63-65.
2. Chopra RN, Chopra IC and Nayak SL. In:Glossary of Indian Medicinal Plants. Council of Industrial Research, New Delhi.1986:5.
3. Chandrasekhar HI. Clinical trial on rheumatoid arthritis with vyoshadi guggulu and vachadi kiratha. *Rheumatism*. 1982:17; 127.
4. Rajagopalan V. Effect of ayushman-8 in manasa mandata (mental retardation). Seminar on research in ayurveda and siddha. 1985:20th – 22nd March; 34.
5. Date BB and Kulkarni PH. Assessment of efficacy of “P-tabs” in insomnia and irritability. *Deerghayu International*. 1995:11(04); 29-34.
6. Date BB and Kulkarni PH. Assessment of efficacy of “Prasham” in insomnia and irritability. *Ayurvedic Research Paper-II*. 1995:15-24.
7. Tripathy JS and Singh RH. Clinical evaluation of “Smrtisagara rasa” in cases of residual schizhophrenia. *Journal of research in ayurveda and siddha*. 1994:15(1-2);8-16.
8. Saravana Kumar A, Mazumder A and Saravanan VS. Antihyperlipidemic activity of *Camellia sinensis* leaves in triton-WR1939 induced albino rats. *Pheng Magnum*.2008: 4(3), 60-64.
9. Khandelwal KR. In:Practical Pharmacognosy technique and experiments. 2nd edn. Pune. Nirali Prakashan Publishers. 2000:149-156.
10. Harborne JB.In: phytochemical methods. A guide to modern techniques of plant analysis, Chapman and Hall, London. 1973:182-189.
11. Peach and Tracey MV.In: Modern methods of plants analysis. Spingler and Veriag Publishers Berlin.1955:3;321-322.
12. Franswoth N.R. Phytochemical investigation on *Cassia tora*. *Journal of Pharmaceutical Sciences*. 1966: 55(3); 225-269.
13. Anes Ahmad Siddqui and Mohamad Ali.In: Practical pharmaceutical chemistry, CBS publishers, New Delhi.1997: 127-137.
- 14.Tyler VE, Brady LR and Robert JE. In: Pharmacognosy. Lea and Febiger Publications, Philadelphia, 9th edn, 1988:77-79.
- 15.Kokatae CK.In: Practical Pharmacognosy.Vallabha Prakashan Publications, New Delhi. 3rd edn. 1991:107-111.
- 16.Biosis MS. Antioxidant determinations by the use of a stable free radical. *Nature*. 1958: 181;1199-1200.