



INTERNATIONAL JOURNAL
OF PHARMA AND BIO SCIENCES

Internationally Indexed Journal

Indexed in Chemical Abstract Services(USA),Index Copernicus ,Ulrichs Directory of Periodicals,Google scholar ,Cabi ,DOAJ ,PSOAR, EBSCO ,SCOPUS, EMBASE etc.



Rapid Publishing

The International Journal of Pharma and Bio Sciences (IJPBS) is an international journal published quarterly. The Aim of IJPBS is to publish peer reviewed research and review articles in less time without much delay in the developing field of Pharmaceutical and Biological sciences. One week from the date of manuscript submission author gets the decision of acceptance and if accepted the manuscript will be processed within 3 weeks (approx.) for publication.

Pharmaceutical Sciences

Branches

- Pharmaceutics
- Novel drug delivery sys
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics
- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics

Biological Sciences

Branches

- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Allied Sciences
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology



Impact Factor 0.47*

**Refer Instruction to authors available at www.ijpbs.net*



Indexed In Elsevier Bibliographic Database
(SCOPUS, EMBASE and Sciverse)

Journal Home page
www.ijpbs.net

FOR INSTRUCTION TO AUTHORS VISIT
www.ijpbs.net

FOR ANY QUERIES EMAIL TO

- editorofijpbs@yahoo.com
- editorijpbs@rediffmail.com
- prasmol@rediffmail.com



EVALUATION OF ETHANOLIC EXTRACTS OF *IN VITRO* GROWN *BAUHINIA VARIEGATA* L. FOR ANTIBACTERIAL ACTIVITIES

MADHU KUMARI*

**Department of Botany, B.R.A.Bihar University, Muzaffarpur.*

ABSTRACT

Nodal segments from six years old plants of *Bauhinia variegata* L. were cultured on Murashige and Skoog's (MS) medium supplemented with different concentration and combinations of BAP and IBA. Explants culture in MS basal medium supplemented with 5mg/l BAP showed highest rate of shoot multiplication. When *in vitro* shoots were inoculated on the MS basal medium supplemented with IBA 2mg/l, profused rooting was observed. *Bauhinia variegata*, commonly known as Kachnar is considered as medicinal plant in Nepal and India. The ethanolic extract of this plant was found to have antimicrobial activity against *Bacillus subtilis* (MTCC8), *E coli* (MTCC1), *Staphylococcus aureus* (MTCC98), *Salmonella typhi* (MTCC737). The largest zone of inhibition (16 mm) was found against *B. subtilis*. The extract was found to be more effective against gram- positive than gram-negative bacteria.

KEYWORDS: *Bahuina variegata*, nodal segment culture, *In vitro* multiplication, multiple shoot induction, plantlet regeneration, antimicrobial activity.



DR. MADHU KUMARI

Department of Botany, B.R.A.Bihar University, Muzaffarpur.

INTRODUCTION

Plants have been considered a valuable source of natural products for maintaining human health, with more intensive studies devoted to natural therapies. The use of plant compounds for pharmaceutical purposes has gradually increased in India. People from developed countries now tend to use traditional medicines, which involve compounds derived from medicinal plants. Natural products of higher plants may give a new source of antimicrobial agents. Certain *Bauhinia* species have a long history of traditional medicinal applications¹. Therefore, such plants have to be investigated to understand their properties, safely and efficiently. *Bauhinia variegata* L. lies in the family Ceasalpiniaceae, and is locally called Kachnar. It is a medium sized deciduous tree, found throughout India. *Bauhinia variegata* is a medicinal plant² is widely used in folklore medicine. Its bark, root, leaves, seeds and flowers are used for their medicinal properties. The root and barks are acrid, cooling, constipating, depurative, anthelmintic³, anti-inflammatory and styptic. They are useful in diarrhea, dysentery, skin disease, leprosy, intestinal worms, wound ulcers, tumors⁴, coughs and antidiabetes^{5,6}. Root contains flavanol glycosides⁷ is reported to have anti-inflammatory activity.

Conventional propagation via seed is relatively slow as per increasing requirement, there is an urgent need for applying non-conventional methods of propagation of *Bauhinia variegata*. Micro-propagation ensures not only continuous supply of plant throughout the year but also prevent the destruction of the natural population of medicinal plant^{8,9}. During past few years, there has been an increase interest for in vitro multiplications and germplasm conservation of rare, endangered, aromatic and medicinal plants^{10,11}. Earlier reports on the cultivation and improvement programs of this taxon focus mainly on the development of efficient protocols for its regeneration through tissue culture methods. The available literature on the medicinal properties of this plant

pertains mainly to the field grown plants. Although the plant was successfully raised under *in vitro* tissue culture condition, to best of our knowledge, there is no published work regarding to compare the medicinal value of the *in vitro* raised plant of *Bauhinia variegata*.

The main objective of our study is to establish an efficient, quick and reliable protocol for in vitro regeneration of *Bauhinia variegata* and study of antimicrobial properties of *in vitro* raised plant mass of *Bauhinia variegata*. The later may be scaled up for commercial production of plants for extraction of pharmaceutically important chemicals.

MATERIALS & METHODS

Source of Explants

In order to obtain *in vitro* plants, a protocol was developed for rapid clonal propagation of *B. variegata* L through mature nodal explants. Nodals & internodal explants were collected from the campus of the department of Botany, B R A Bihar University, Muzaffarpur. These nodal segments were cuts into 1.5cm to 2.0cm length with single node & internode intact. These nodal & internodal cutting were washed with 5% (v/v) detergent solution (tepol) for 10minutes followed by rinsed with running tap water for several times. These nodal segments were surface sterilized with sterile bavistin 0.2% (w/v) for 10 minutes and then washed with sterile distilled water. In the laminar chamber the nodal segments were further treated with 70% alcohol for one minute followed by 0.1% (w/v) mercuric chloride treatment for 5 minutes, aseptically explants were washed with sterile distilled water for 3 to 4 times dried using sterile blotting paper which was ultimately used as explants for raising *in vitro* cultures.

Growth Medium & Culture Condition

Murashige and Skoog's ¹²(MS) medium 1962 was used as nutrient source with or without growth hormones such as BAP, IBA either

individually or in combination for culture explants. Sucrose 3% and agar 0.8% (Hi media) were used as carbon source and gelling agent respectively. The pH of the medium was adjusted to 5.8 and autoclaved at 121°C for 20 minutes. The cultures were maintained in the culture room at 25±C 2°C, under white fluorescent light.

Shoot Organogenesis and Elongation

For multiple shoot induction shoot nodal explants were culture on MS medium supplemented with BAP (1- 5 mg/l) alone or combination with Kinetin (1-2 mg/l) (Table 1). Data on percentage of responding explants and number of shoots per explants were recorded after 25 days of initiation culture.

Antibacterial Assay

Preparation of Plant Extract

In vitro grown plants, callus, were weighted and grinded with help of motor & pestle. About 50gm of power grinded material was extracted in soxhlet apparatus with 250ml of ethanolic solvent¹³. The extracted solvent was removed from the extract under reduced pressure with rotary vacuum evaporator. The sticky greenish-brown substance was obtained was stored in refrigerator condition till use¹⁴.

Culture of Microorganism

Both gram positive and gram negative bacteria were used as test organism for this study. The organisms like *Escherichia coli* MTCC 2845, *Salmonella typhi* MTCC 737, *Staphylococcus aureus* MTCC 98, *Bacillus subtilis* MTCC 441, were used for antibacterial assay. The bacteria were grown in nutrient broth media at 37°C, for antibacterial assays. The disc diffusion method¹⁵ was adopted for antimicrobial study

of plant extract. Briefly, 20ml of media was transferred aseptically into each sterile petri dish and allowed to solidify. An over night grown inoculums (100µl) suspension was spread uniformly over the agar medium using sterile glass rod for uniform distribution of bacteria. The readily prepared sterile discs were loaded with different concentrations of about 10-20 µg of plant extract of *B. variegata*. The paper diffuse discs were placed on the medium and plates were incubated at 37°C for 24 hours. The antibacterial activity was recorded by measuring the width of the clear inhibition zone around the disc in millimeter.

Phytochemical Screening

The ethanolic extract of *in vitro* grown plants of *Bauhinia variegata* L were subjected to different tests^{16,17} to identify the nature of chemical constituents present in the plant material.

RESULTS & DISCUSSION

Organogenesis is dependent on the factors like explants type physiological state of donor plant or organ and endogenous level of phytohormones¹⁸. It has been reported by various workers that the balance of auxin to cytokinin is a determining factor¹⁹. The media supplemented with BAP alone triggered shoot induction in nodal explants of *Bauhinia variegata* (fig-A). The appreciable frequency of shoots emerged from each nodal explants when it cultured on MS medium supplemented with high concentration of BAP (5mg/l). Further, increase in concentration of BAP reduced the frequency of shoot induction. BAP alone at the concentration of 5mg/l produced maximum shoot per explants. (fig-B)




Figure 1

***In vitro* clonal propagation of *Bahuina variegata* L from internodal explants. A. shoot buds initiation on MS + BAP (2mg/l); B. multiple leafy shoot emergence in MS + BAP (5mg/l); C. shoots emerged from callus in MS + BA (2.5mg/l) + KN (0.5mg/l). D. Regenerated plantlets after 40 days old culture.**

When the explants were cultured on MS medium with BAP (2.5mg/l) and kinetin (0.5mg/l) the callus as well as shoots induction take place after 30 days (fig-C). When BAP was used in combination with IBA the inductions of shoot as well as root both take place. The induction of roots from callus was observed on MS basal media containing only IBA. The frequency of root induction observed in different concentration of IBA (1mg to 2 mg/l) (Table-1). However, the concentration of IBA higher than this the frequency of root induction decreased. It is well known that root induction depends on the auxin concentration in tissue²⁰.

***In vitro* Rooting Plant Establishment**

The *in vitro* grown plantlets (fig-D) were ready for acclimatization. After hardening it is transferred into the soil. The auxin IBA were used singly to induce rooting from *in vitro* raised shoot lets. A range of concentration was tested (1.0, 2mg/l) for rooting. The superior results on rooting were obtained on MS with IBA (2mg/l). The well rooting plant lets were transferred to plastic cups containing vermiculite for hardening and kept under controlled condition.

Plants started producing fresh shots and roots after one week of transplanting. Later they

were transported to the field condition and the survival rate was 50% (Table-2). The efficient micro propagation technique described here may be highly useful for raising disease free quality planting material of *Bauhinia variegata*

for commercial and off season cultivation which not only help in improving economic condition of the farmer but also fulfill the market demand of herbal industries.

TABLE 1
Effects of Cytokinins and Auxin and their Interaction on Shoot Proliferation from Nodal Explants of *Bauhinia variegata*.

Growth regulator	Concentration (mg/l)	% of response	Induction of shoots per explants	Induction of roots per explants
BAP	0.5	0	0.0	0.0
	1	0	0.0	0.0
	2	1	1	0.0
	3	40	1-2	0.0
	4	60	2-3	0.0
	5	70	4-5	0.0
BAP+ KN	1 + .5	70	2-3	0.0
	2.5+ .5	74	3	0.0
	1+ 1	65	0	0.0
IBA	.5	0.0	0	0.0
	1	40	0	1-2
	2	50	0	1-2

Table-2
Data on Survival of Transferred Regenerated Plants.

	No. of Plants	Plants Survived	% Response
Transferred for Hardening	30	20	66.6
Transferred for Acclimatization	20	10	50

In this investigation, during the process of transfer to field the survival percentage was found to be fifty percent. There are various reasons for low survival of regenerated plants

after transfer to natural soil. Gangopadhyay et al.²¹ (2002) reported that *in vitro* derived roots are often physiological functioning when in contact with the soil.

Table 3

Shows the antibacterial effect of ethanolic extract of *Bauhinia variegata*. The Result obtained in the present study revealed that EBV possesses potential antibacterial activity against all the four tested bacterial organisms (*E. coli*, *S. typhi* and *S. aureus* *B. subtilis*).

S.No.	Microorganisms	Zone of Inhibition(mm) 10µg	Zone of inhibition(mm) 20µg
1	<i>Escherichia coli</i> (-ve)	6mm	10mm
2	<i>Salmonella typhi</i> (-ve)	5mm	8mm
3	<i>Staphylococcu aureus</i> (+ve)	7mm	12mm
4	<i>Bacillus subtilis</i> (+ve)	9mm	16mm

The EBV showed a broad spectrum of activity against all the bacterial strains at the tested concentration of 10 - 20µg/disc. EBV exhibited greater zone of inhibition for *Bacillus subtilis* (16mm). The least activity was observed for *S. typhi* (5mm) at 10 µg/disc.

The plants and their extracts used against microbial infections due to the presence of secondary metabolites such as phenols, flavonoids, tannins, terpenoids etc^{22,23} have

documented the use of natural products as new antibacterial drugs.

Phytochemical Screening

The phytochemical tests revealed the presence of flavonoids, saponin, Alkaloids, fatty acid, Tannins, Glycosides in ethanol extract of *in vitro* grown plants of *Bauhinia variegata* L. The results of phytochemical screening are given in Table 3.

Table 4
Phytochemical Screening

S.NO	Name of the Phytoconstituents	Ethanolic Extract of <i>Bauhinia variegata</i> Plants
1	Alkaloids	+
2	Flavonoids	++
3	fatty acid	+
4	Saponins	+
5	Glycosides	++
6	Tannins	+

(++) presence of compound, (+) moderately presence of compound

The result justify the medicinal use of *Bauhinia variegata* L. as an anti-infectious diseases, as a natural sources. Thus this plant could be utilized as a source of antimicrobial drugs, with little or no side effects as compared to synthetic drugs.

CONCLUSION

The ethanolic extract of *Bauhinia variegata* L. was found to be sensitive to different

REFERENCES

1. Valdir CF, Chemical composition and biological potential of plants from the genus *Bauhinia*. *Phytother Res.* 23:1347-54, (2009).
2. Warriar PK, Knambiar, VP and Amankutty CR, Indian Medicinal Plants, A Compendium of 500 species, vol. 1. Madras, India: Orient Longman Limited, ISBN 0-86311464-4, (1993).
3. Ram PR, Mehrotra BN, In: Compendium of Indian medicinal plants. Vol. 3. New Delhi: Publication and information directorate; 1980.p. 84-91.
4. Raj Kapoor B, Jayakar B, Murugesh N, Antitumour activity of *Bauhinia variegata* on Dalton's ascetic lymphoma. *J Ethnopharmacol* 89:107-9, (2003).
5. Azevedo CR, Maciel FM, Silva LB, Ferreira AT, da Cunha M, Machado OL, Fernandes KV, Oliveira AE and Xavier-Filho J, Isolation and intercellular localization of insulin-like proteins from leaves of *Bauhinia variegata*. *Braz J Med Bio Res.* Nov, 39(11); 1435-44 (2006).
6. Yadava RN, Reddy VM, Anti-inflammatory activity of a novel flavonol glycoside from the *Bauhinia variegata* L. *Nat Prod Res* 2003; 7:165-9.
7. The Wealth of India, A Dictionary of Indian Raw Materials and Industrial products. Vol 2 CSIR New Delhi: p. 56-7, (1959).
8. Datta PC, Biotechnology and Tissue culture of some medicinal plants, In: JN Govil, VK Sing and Hshmi (eds.), Glimpses in plant Research, Medicinal plants: New Vista of Research part 2, Vol XI, pp.337-342, (1993).
9. Hamilton, AC, Medicinal plants, Conservation and livelihoods. *Biodiversity Conser.* 13: 1477-1517, (2004).
10. Tiwari V, Singh BD, Tiwari KN, Shoot regeneration and Somatic embryogenesis from different explants of Brahmi [*Bacopa monniera*(L.) Wettst]. *Plant Cell Rep.* 17: 538-543, (1998).
11. Villalobos VM, Engelmann F, *Ex situ* conservation of plant germplasm using biotechnology. *World J. Microb. Biotech.* 11: 375-382, (1995).
12. Murashige T, Skoog FA, Revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15: 473-497, (1962).
13. Vogel AI, In: Elementary practical organic chemistry (Second edition), Orient Longman Limited, pp. 45-168, (1988).
14. Beyer H and Walter W, Organic chemistry, 4th ed. Published by Jerry March, pp 146-148, (1997).
15. Bayer AW, Kirby. MDK, Sherris. JC, and Turck, Antibiotic susceptibility testing by standard single disc diffusion method, *Am. J. Clinical pathol.*, 493-496, (1966).

ACKNOWLEDGEMENT

The author would like to express her gratitude to the Head, Department of Botany, B.R.A. Bihar University, Muzaffarpur for providing the necessary laboratory facilities.

16. Kokate CK Practical Pharmacognosy, 4 th ed, Vallabh Prakashan, Delhi, pp.107-111, 1994.
17. Khandelwal KR, Practical Pharmacognosy", 11th ed. Nirali Prakashan, Pune, pp. 149-153, 2004.
18. Thanh, KT and Trinh TH, Organogenic differentiation. In: S.S. Bhowani (ed.), Plant Tissue Culture application and limitations, Elsevier, USA. pp. 34-53, (1990).
19. Zheng SS, Yuan HY, Wang LJ, An CC and Chen ZL, The tissue culture of medicinal plant *Trichosanthes kirilowii* and its protein analysis. Sheng wu Gong cheng Xue Bao, 17: 420-422, (2001).
20. Narayanswamy S., Plant Cell and Tissue Culture. Tata McGraw Hill Publishing Company Limited, New Delhi, Chapter 3, pp. 22-51, (1994).
21. Gangopadhyay G, Mitra S Das, Poddar S K, Modak R and Mukherjee, KK, Enhanced rate of multiplication and rooting through the use of coir in aseptic liquid culture media. Plant Cell Tiss. Org. Cult.68: 301-310, (2002).
22. Anandaprakash RL, Natural constituents of *Bauhinia variegata*. Journal of Research into Indian Medicinal, Yoga and Homeopathy 13, 96-97, (1978).
23. Gupta AK, Vidyapati TS and Chauhan JS, Chemical examination of the stem of *Bauhinia variegata*. Planta Medica 38, 174-176, (1980).
24. Silver LL, and Bostian KA, Discovery and development of new antibiotics: the problem of antibiotic resistance, Antimicrob. Agents Chemother. 37, 377-383(1993).