



Internationally indexed journal

Indexed in Chemical Abstract Services (USA), Index copernicus, Ulrichs Directory of Periodicals, Google scholar, CABI ,DOAJ , PSOAR, EBSCO , Open J gate , Proquest , SCOPUS , EMBASE ,etc.



Rapid and Easy Publishing

The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quarterly. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences



Pharmaceutical Sciences

- Pharmaceutics
- Novel drug delivery system
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics



Biological Sciences

- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics
- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

**Indexed in Elsevier Bibliographic Database
(Scopus and EMBASE)**

SCImago Journal Rank 0.288

Impact factor 2.958*

Chemical Abstracts
Service (www.cas.org)



A division of the American Chemical Society

CODEN IJPBJ2



Elsevier Bibliographic databases (Scopus & Embase)

SNIP value – 0.77

SJR - 0.288

IPP - 0.479

SNIP – Source normalised impact per paper

SJR – SCImago Journal rank

IPP – Impact per publication

Source – www.journalmetrics.com

(Powered by scopus (ELSEVIER))



LUND
UNIVERSITY



JACKSONVILLE STATE UNIVERSITY

Jacksonville State University
Houston Cole Library
USA (Alabama)



UNIVERSITY OF
OXFORD

Oxford, United Kingdom

INDEX COPERNICUS
INTERNATIONAL

*And indexed/catalogued in
many more university*



*Instruction to Authors visit www.ijpbs.net

For any Queries, visit "contact" of www.ijpbs.net



SIGNIFICANCE OF TACROLIMUS MONITORING TO OPTIMIZE IMMUNOSUPPRESSIVE THERAPY

PARVEEN POPLI AND PRANAV KUMAR PRABHAKAR*

Department of Paramedical Sciences, Lovely Professional University, Punjab India-144411.

ABSTRACT

Liver transplantation is the surgical treatment of choice for several irreversible liver diseases where a partially or completely diseased liver is replaced with a healthy liver. The continued use of immunosuppressant causes certain serious consequences such as hepatotoxicity, nephrotoxicity and post-transplant hyperglycemia. Inter individual variation has necessitated frequent monitoring of tacrolimus concentration to prevent toxicity, which is the major clinical challenge encountered in the post-transplant period. This study was carried out with the aim to evaluate the correlation between blood concentration of tacrolimus and the risk/degree of toxicity with time after liver transplant. Our data suggests that monitoring effective immunosuppression is essential to minimize toxicity induced by tacrolimus, especially in the 3-6 month period. Therefore, it is necessary to maintain trough blood concentration in liver transplant recipient's 10-12 ng/ml in first trimester, 8-10 ng/ml in next trimester and 5-8 ng/ml thereafter. Hence, monitoring of tacrolimus concentration of the recipient should be done throughout the period of tacrolimus intake i.e. for life time, to enhance clinical evaluation.

KEYWORDS: Tacrolimus, Immunosuppression, Transplant, Liver, Hepatotoxicity

*Corresponding author



PRANAV KUMAR PRABHAKAR

Department of Paramedical Sciences, Lovely Professional University, Punjab India-144411.

INTRODUCTION

Liver transplantation or hepatic transplantation is the replacement of diseased liver with a healthy liver from a donor by the surgical means. Liver transplantation is a viable treatment choice for end-stage liver hepatic disease and acute liver failure. Similar to other allograft, a liver transplant will be rejected and removed by the recipient's immune system unless an immunosuppressive drugs are used. Most of the liver transplant recipient receives corticosteroids plus a calcineurin inhibitor such as tacrolimus or cyclosporine to stop graft rejection¹. Recent clinical results shows that the tacrolimus is a better choice than the cyclosporine during the first year of transplant^{1,2}. Tacrolimus (also known as fujimycin or FK506) is a 23-membered macrolide lactone (Figure 1). It is produced by the fermentation broth of *Streptomyces tsukubaensis*^{3,4}. It is an immunosuppressive agent belongs to a calcineurin inhibitor group which is used in solid organ transplant as a therapeutic drug⁵. Tacrolimus suppresses the natural immunity against a foreign body and thus, prevents rejection in liver, kidney, lung,

pancreas and bone marrow recipients⁶⁻⁸. Immunophilins, a group of highly conserved proteins, mostly participate in protein folding. Inside cytoplasm, tacrolimus binds with respective cytosolic immunophilins such as FKBP-12 and FKBP-52 (an FK506-binding proteins), a component of the glucocorticoid receptor complex. This complex binds and inhibits the activity of the calcineurin, a calcium/calmodulin-dependent protein phosphatase that is expressed in all mammalian tissues⁹. The said complexes interact with calcium-dependent signal pathway in immunological T-cells. Calcineurin then increases the interference of leads to interference with translocation to the nucleus of various nuclear factors involved in the transcription of cytokine genes. As a result of this inhibition, the transcription of the genes involved in T-cell activation is suppressed, affecting the production of interleukin-2 (IL-2) and various other cytokines and interleukins, such as interleukin-3 (IL-3), interferon- γ (INF- γ), and tumor necrosis factor- α (TNF- α)¹⁰.




Figure 1
Tacrolimus structure (Adapted from Tanaka HKA, 1987)

Tacrolimus is metabolized by gastrointestinal cytochrome P450 (CYP)3A isoenzyme, 3A4 and 3A5, through O-demethylation and hydroxylation metabolic reaction in the liver and

intestinal wall. It is then removed by P-glycoprotein¹¹. P-glycoprotein decreases the intracellular concentration of tacrolimus by reversely pumping out the absorbed drug into

intestinal lumen. P-glycoprotein may also regulate access of tacrolimus to CYP3A enzyme and prevent these enzymes from being overcome by high drug concentration in intestine¹²⁻¹⁴. Tacrolimus is repeatedly moved out of intestinal mucosal cell and then passively reabsorbed. 95% tacrolimus metabolites are eliminated by the biliary route¹¹. Increase tacrolimus metabolites concentration in blood can be caused by biliary obstruction¹⁵⁻¹⁶. During drug therapy with several kinds of drugs like antibiotics, immunosuppressants, etc. there is a direct toxic effect on the liver and immune system and under these conditions the hepatocytes are injured, the intracellular enzymes are released into the circulation and, hence, elevated levels of aminotransferases and gamma-glutamyltransferase are seen. This is due to the fact that these drugs are metabolized in the liver. Tacrolimus is metabolized in the liver by cytochrome P450 (CYP3A4) enzyme and cleared by the P-glycoprotein. Hence, elevated circulating concentrations of the hepatic enzymes indicate hepatotoxicity in post-transplant period¹⁷. The resultant effects on the transcription and the mRNA degradation of several cytokine genes of interleukin-2; this plays a critical role in T-cell proliferation associated with alloimmune reactions¹⁸. Kidney contains a high concentration of tacrolimus binding proteins. At the cellular level, it has been suggested that tacrolimus nephrotoxicity targets tubular epithelial cells, vascular endothelial cells, arteriolar myocytes, and interstitial fibroblasts. Tacrolimus also has a direct toxic effect on renal tubules which can be evidenced by the epithelial

vacuolization¹⁹. The vascular toxicity of tacrolimus caused by Vasospasm leading to reduced glomerular filtration²⁰. Interstitial fibrosis are recognized as another mechanism of tacrolimus nephrotoxicity which leads to narrowing of small arteries and arterioles. As the narrowing proceeds, a linear zone of renal parenchymal vessels is deprived of its nutrition and begins to undergo degenerative changes. The renal parenchymal cells are susceptible to injury because of their relatively low oxygen availability²¹⁻²³. The cytokine TGF- β is likely to involve in the pathogenesis of interstitial fibrosis²⁴. The toxic effect of tacrolimus on the hormone producing endocrine pancreas is attributable to the selective localisation of FK-binding protein (FKBP)-12 and calcineurin in the pancreatic islets. The intracellular calcium binding protein calmodulin plays an important role in insulin secretion. Tacrolimus binds with calmodulin which may inhibit insulin secretion and cause diabetes mellitus²⁵⁻²⁶.

MATERIALS AND METHODS

An analysis of data from 36 patients who has gone for liver transplant has been performed. Subjects who received oral tacrolimus as a main immunosuppressant strategy were included in the study. Analysis of whole blood tacrolimus, Liver function tests (Bilirubin, AST, ALT, ALP, GGT) for monitoring hepatotoxicity, Kidney function tests (blood urea nitrogen, creatinine) for monitoring nephrotoxicity and plasma glucose levels for hyperglycemia were evaluated after 3 months, 6 months and 9 months post-transplant.

Toxicity was ascertained on the basis of the following Tacrolimus levels:

Early post-transplant (up to 3 months) > 15 ng/ml
3 to 6 months >10 ng/ml
6 to 9 months >8 ng/ml
After 9 months >7 ng/dl

Hepatotoxicity were diagnosed on the basis of various enzymes level –

AST/ALT > 60 IU/L
ALP > 140 IU/L
GGT > 95 IU/L

At least two enzymes in the toxic range in two successive readings within a week was considered as hepatotoxicity. Nephrotoxicity was diagnosed when serum creatinine level was measured more than 1.3 mg/dl on two successive readings within a week. Hyperglycemia was considered when blood glucose level was found more than 120 mg/dl in two successive readings within a week.

RESULTS

In this study, we observed whole blood tacrolimus concentration of liver transplant recipients for one year period. Also, we estimated biochemically the liver functions, kidney functions and blood glucose levels throughout this one-year period. We then

established correlation between tacrolimus concentration and hepatotoxicity, nephrotoxicity and hyperglycemia. Tacrolimus dose-Liver transplant patients are given tacrolimus post-transplant as per the defined protocol i.e. 0.1-0.3/ mg/kg/day²⁷.

Tacrolimus levels in post-transplant period

Though appropriate use of immunosuppressants is essential in transplant patients, long term use of immunosuppressive drug causes toxicity. Hence, the relative risks of toxicity and efficacy failure are both related to Tacrolimus concentrations. Table 1 shows tacrolimus concentrations and percentage of subject's distribution at 3rd months, 6th months and 9th months.

Table 1
Whole blood tacrolimus concentration range for different time spans of 3rd months, 6th months and 9th months

Tacrolimus concentration	3 rd Month	6 th Month	9 th Month
3 - 5 ng/ml	3.5%	11%	20%
5.1 – 8 ng/ml	16%	28%	45%
8.1- 10 ng/ml	25%	43%	27%
10– 12 ng/ml	36%	13%	5%
above 12 ng/ml	19.5%	5%	3%

Due to tacrolimus, various kinds of toxicity were notices such as hepatotoxic, nephrotoxic and hyperglycemia. In the 3rd month non-toxic subjects were 36.5 % and that of 6th month was 56%. At the end of 9th month non-toxic subjects were almost similar to 6th month i.e., 53 % (Table 2).

Table 2
Prevalence of tacrolimus toxicity for different time spans of 3rd months, 6th months and 9th months

Toxicity	3 rd Month	6 th Month	9 th Month
Hepatotoxic	22%	22%	22%
Nephrotoxic	22%	11%	16.50%
Hyperglycemic	19.50%	11%	8.50%
Nontoxic	36.50%	56%	53%

Three months post-transplant mean tacrolimus concentration was found to be 10.91±4.57 ng/ml; and after 6 months it was 8.00±2.94 ng/ml; where that of after 9 months was 7.13±2.77.

Table 3
Statistical descriptors for the tacrolimus level for 3rd months, 6th months and 9th months

Months	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
3	36	10.914	4.57	0.7617	9.368	12.46	4.7	26.3
6	36	8	2.9419	0.4903	7.005	8.995	2.9	14.7
9	36	7.131	2.7688	0.4615	6.194	8.067	2.7	15.5
Total	108	8.681	3.8487	0.3703	7.947	9.416	2.7	26.3

In each group of patients (patients reviewed during the first 3 months, 3-6 months and 6-9 months), a window of blood tacrolimus levels associated with least toxicity was identified. At the end of 3rd months, 6th months and 9th months this window was identified as 9-11 ng/mL, 8-10 ng/mL and 5-8 ng/mL, respectively. Correlation between tacrolimus levels and toxicity associated parameters were derived in the post-transplant period. During first three months 22% subjects has shown hepatotoxicity, 22% subjects has shown nephrotoxicity and 19.5% subjects has shown hyperglycemia when 53% subject's tacrolimus value were 10-12 ng/ml. After 6 months 22% subjects had hepatotoxicity, 11% subjects had nephrotoxicity and 11% subjects had hyperglycemia when 50% subjects tacrolimus value were 8-10 ng/ml. After 9 months 22% subjects had hepatotoxicity, 16.5% subjects had nephrotoxicity and 8.5% subjects had hyperglycemia when 48.5% subjects tacrolimus value were 5-8 ng/ml. All these levels were lower than the recommended upper limit of trough i.e.20ng/ml. Though the therapeutic range of tacrolimus is defined as 5-20 ng/ml, clinicians are facing dilemma because even within this range several instances of toxicity as well as ineffective immunosuppression were encountered. This could be partly due to inter-individual and intra-individual variations in pharmacokinetics. Moreover, adequate immunosuppression varies with the time after transplant. Hence, long-term compliance is required to ensure graft survival in transplant patients²⁷⁻²⁹. This is achieved by regular therapeutic monitoring of tacrolimus throughout life in the post-transplant period. The

importance of the need for therapeutic monitoring of tacrolimus is further supported by the present study. Drug toxicity in the postoperative period is influenced by the drug-metabolizing capacity of the graft. This capacity depends primarily on the levels and activities of the cytochrome P450 enzymes (P450). P450 enzymes play a central role in the biotransformation of various xenobiotics to more polar compounds, which are readily excreted³⁰. The metabolites of a drug can be inactive or less active than the parent compound, although some biotransformation products show enhanced pharmacological or toxicological activity. Any change in the activity of P450 isoforms influences the rate of activation or inactivation of drugs³¹. Lack of functional activity in P450 enzymes need tailored drug treatment in order to avoid excessive drug concentration and toxic effects. Dose adjustment or selection of an alternative drug, which is not a substrate for the polymorphic enzyme, can prevent the development of side-effects in poor metabolizers. Tacrolimus is known to be toxic to the liver so that when the hepatocytes are injured, the intracellular enzymes are released into the circulation. Hence, this effect of tacrolimus on liver function is best observed by activity of alkaline phosphate (ALP), alanine aminotransferase (ALT), aspartate aminotransferase and γ -glutamyltransferase. Tacrolimus binding proteins are present at a high concentration in the kidney. Calcineurin immunoreactivity and enzyme activity in the kidney can be specifically inhibited by tacrolimus. At the cellular level, it has been suggested that tubular epithelial cells, vascular

endothelial cells, arteriolar myocytes, and interstitial fibroblasts are all targets for tacrolimus nephrotoxicity. The occurrence of epithelial vacuolization is evidence of direct toxic effect of tacrolimus on the renal tubule²⁵. Vasospasm leading to reduced glomerular filtration appears to be a key element in the vascular toxicity tacrolimus²⁶. Interstitial fibrosis are recognized as tacrolimus nephrotoxicity due to narrowing of the small arteries and arterioles. As the narrowing proceeds, a linear zone of renal parenchymal vessels is deprived of its nutrition and, begins to undergo degenerative changes. The renal parenchyma is particularly susceptible to injury because of their relatively low oxygen availability²⁷⁻²⁹. This direct toxicity is characterized by tubular degeneration and atrophy, leading to fibrosis. The cytokine transforming growth factor beta (TGF- β), a well-known activator of collagen transcription, is likely involved in the pathogenesis of interstitial fibrosis. Tacrolimus nephrotoxicity can be said to be present if there is an elevated blood urea nitrogen and/or elevated serum creatinine levels in the post-transplant period. In this study, all subjects with a serum creatinine > 1.3 mg/dl were deemed to have nephrotoxicity. The toxic effect of Tacrolimus on the endocrine pancreas is attributable to the selective localisation of FK-binding protein (FKBP)-12 and calcineurin in the pancreatic islets. The intracellular calcium binding protein calmodulin plays an important role in insulin secretion. Tacrolimus binds with calmodulin which may inhibit insulin secretion and cause diabetes mellitus.

REFERENCES

1. Tanaka HKA, Marusawa H. Physicochemical properties of FK506, a novel immunosuppressant isolated from *Streptomyces tsukubaensis*. *Transplant Proc*, 14:11-16, (1987).
2. Mierke DF, Schmieder P, Karuso P, Kessler H. Conformational Analysis of the cis- and trans-Isomers of FK506 by NMR and Molecular Dynamics. *Helv Chim Acta*, 74:1027-1047, (1991).
3. Plosker GL, Foster RH. Tacrolimus: a further update of its pharmacology and therapeutic use in the management of organ transplantation. *Drugs*, 59:323-389, (2000).
4. Stratta RJ, Taylor RJ, Castaldo P, Sindhi R, Sudan D, Weide LG, *et al.*, FK 506 induction and rescue therapy in pancreas transplant recipients. *Transplant Proc*, 28:991-992, (1996)

In our analysis, 10–12 ng/ml is the safe therapeutic range of trough levels of tacrolimus for liver transplant recipients in first 3 months, 8-10 ng/ml in the next three months i.e upto 6 months, and 5-8 ng/ml thereafter. Hence, maintaining trough concentrations of tacrolimus within the specified ranges may reduce the degree of complications without compromising efficacy.

CONCLUSION

Inadequate dose of immunosuppressants like tacrolimus is associated with an increased risk for rejection, whereas excessive administration of these immunosuppressants is associated with toxicity. Generally, high tacrolimus concentrations are likely to be required in the initial post-transplant period, but target concentrations need to be reduced over time to minimize toxicity. Our data suggest that monitoring effective immunosuppression is essential to minimize toxicity induced by tacrolimus, especially in the 3-6 month period. In liver transplant recipients, it is necessary to maintain trough blood concentration of 10- 12 ng/ml in first 3 months, 8-10 ng/ml in next three months and 5-8 ng/ml thereafter. Hence, it is necessary to monitor tacrolimus concentration of the transplant recipient throughout the period of tacrolimus intake i.e. for life time.

CONFLICT OF INTEREST

Conflict of interest declared none.

5. Kur F, Reichenspurner H, Meiser BM, Welz A, Furst H, Muller C, *et al.*, Tacrolimus (FK506) as primary immunosuppressant after lung transplantation. *Thorac Cardiovasc Surg*, 47:174-178, (1999).
6. Lipson DA, Palevsky HI, Kotloff RM, Edelman J. Conversion to tacrolimus (FK506) from cyclosporine after orthotopic lung transplantation. *Transplant Proc*, 30: 1505-1507, (1998).
7. Almawi WY, Melemedjian OK. Clinical and mechanistic differences between FK506 (tacrolimus) and cyclosporin A. *Nephrol Dial Transplant*, 15:1916-1918, (2000).
8. Paul LC. Overview of side effects of immunosuppressive therapy. *Transplant Proc*, 33: 2089-2091, (2001).
9. Tuteja S, Alloway RR, Johnson JA, Gaber AO. The effect of gut metabolism on tacrolimus bioavailability in renal transplant recipients. *Transplantation*, 7:1303-1307, (2001).
10. Zhang Y, Benet LZ. The gut as a barrier to drug absorption: combined role of cytochrome P450 3A and P-glycoprotein. *Clin Pharmacokinet*, 40:159-168, (2001).
11. Moller A, Iwasaki K, Kawamura A, Teramura Y, Shiraga T, Hata T, *et al.*, The disposition of ¹⁴C-labeled tacrolimus after intravenous and oral administration in healthy human subjects. *Drug Metab Dispos*, 27: 633-636, (1999).
12. Wachter VJ, Silverman JA, Zhang Y, Benet LZ. Role of P-glycoprotein and cytochrome P450 3A in limiting oral absorption of peptides and peptidomimetics. *J Pharm Sci*, 87: 1322-1330, (1998).
13. Benet LZ, Izumi T, Zhang Y, Silverman JA, Wachter VJ. Intestinal MDR transport proteins and P-450 enzymes as barriers to oral drug delivery. *J Control Release*, 62:25-31, (1999).
14. Venkataramanan R, Jain A, Warty VW, Abu-Elmagd K, Furakawa H, Imventarza O, *et al.*, Pharmacokinetics of FK 506 following oral administration: a comparison of FK 506 and cyclosporine. *Transplant Proc*, 23:931-933, (1991).
15. Cadoff EM, Venkataramanan R, Krajack A, Jain AS, Fung JJ, Todo S, *et al.*, Assay of FK 506 in plasma. *Transplant Proc* 1990; 22:50-51, (1990).
16. Christians U, Braun F, Schmidt M, Kosian N, Schiebel HM, Ernst L, *et al.*, Specific and sensitive measurement of FK506 and its metabolites in blood and urine of liver-graft recipients. *Clin Chem*, 38:2025-2032, (1992).
17. Gonschior AK, Christians U, Winkler M, Linck A, Baumann J, Sewing KF. Tacrolimus (FK506) metabolite patterns in blood from liver and kidney transplant patients. *Clin Chem*, 42:1426-1432, (1996).
18. Cardenas ME, Zhu D, Heitman J. Molecular mechanisms of FK506, cyclosporine and rapamycin. *Curr Opin Nephrol Hypertens*, 4:472-477, (1995).
19. McCauley J, Farkas Z, Prasad S, Plummer H, Starzl T, Murray S. Cyclosporine A and FK506 induced inhibition of renal cell (LLC-PKI) proliferation in culture. *Transplant Proc*, 23:2829-2830, (1991).
20. Lieberman KV, Lin WG, Reisman R. FK506 is a direct glomeruloconstrictor as determined by electrical resistant pulse sizing. *Transplant Proc*, 23:3119-3120, (1991).
21. Stillman IE, Andoh TF, Burdmann EA, Bennett WM, Rosen S. FK506 nephrotoxicity: morphologic and physiologic characterization of a rat model. *Lab invest*, 73:794-803, (1995).
22. Andoh TF, Lindsley J, Franceschini N, Bennett WM. Synergistic effects of cyclosporine and rapamycin in a chronic nephrotoxicity model. *Transplantation*, 62:311-316, (1996).
23. Rosen S, Greenfield Z, Brezis M. Chronic cyclosporine-induced nephropathy in the rat: a medullary ray and inner stripe injury. *Transplantation*, 49: 445-452, (1990).
24. Burdmann EA, Andoh TF, Nast CC, Evan A, Connors BA, Coffman TM, *et al.*, Prevention of experimental cyclosporin-induced interstitial fibrosis by losartan and enalapril. *Am J Physiol*, 269: F491-F499, (1995).

25. Davis M. Immunosuppressive drugs. In: Neuberger J, Lucey MR. editors. Liver transplantation: practice and management. London: British Medical Association 1994: 190-209, (1994).
26. Hirano Y, Hisatomi A, Ohara K, Noguchi H. The effects of FK506 and cyclosporine on the exocrine function of the rat pancreas. *Transplantation*, 54: 883-887, (1992).
27. Tsunoda S, Aweeka F. The use of therapeutic drug monitoring to optimize immunosuppressive therapy. *Clin Pharmacokinet*, 30: 107-140, (1996).
28. Kelly P, Burckart G, Venkataramanan R. Tacrolimus: a new immunosuppressive agent. *Am J Health Syst Pharm*, 52:1521-1535, (1995).
29. Jusko W, Thomson A, Fung J, McMaster P, Wong S, Zylber-Katz E, et al., Consensus document: therapeutic monitoring of tacrolimus (FK-506). *Ther Drug Monit*, 17: 606-614, (1995).
30. Lewis DF. 57 varieties: the human cytochromes P450. *Pharmacogenomics*, 5: 305–318, (2004).
31. Solus JF, Arietta BJ, Harris JR, Sexton DP, Steward JQ, McMunn C, et al., Genetic variation in eleven phase I drug metabolism genes in an ethnically diverse population. *Pharmacogenomics*, 5: 895–931, (2004).