



Internationally indexed journal

Indexed in Chemical Abstract Services (USA), Index copernicus, Ulrichs Directory of Periodicals, Google scholar, CABI ,DOAJ , PSOAR, EBSCO , Open J gate , Proquest , SCOPUS , EMBASE ,etc.



Rapid and Easy Publishing

The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quarterly. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences



Pharmaceutical Sciences

- Pharmaceutics
- Novel drug delivery system
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics



Biological Sciences

- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics
- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

**Indexed in Elsevier Bibliographic Database
(Scopus and EMBASE)**

SCImago Journal Rank 0.288

Impact factor 2.958*

Chemical Abstracts
Service (www.cas.org)



A division of the American Chemical Society

CODEN IJPBJ2



Elsevier Bibliographic databases (Scopus & Embase)

SNIP value – 0.77

SJR - 0.288

IPP - 0.479

SNIP – Source normalised impact per paper

SJR – SCImago Journal rank

IPP – Impact per publication

Source – www.journalmetrics.com

(Powered by scopus (ELSEVIER))



LUND
UNIVERSITY



JACKSONVILLE STATE UNIVERSITY

Jacksonville State University
Houston Cole Library
USA (Alabama)



UNIVERSITY OF
OXFORD

Oxford, United Kingdom

INDEX COPERNICUS
INTERNATIONAL

*And indexed/catalogued in
many more university*



*Instruction to Authors visit www.ijpbs.net

For any Queries, visit "contact" of www.ijpbs.net



NUTRITIONAL VALUE OF TRADITIONAL WILD VEGETABLES USED BY THE KINNAURA TRIBALS OF HIMACHAL PRADESH, INDIA.

GEETA*, RICHA AND M.L. SHARMA

Department of Botany, Panjab university, Chandigarh 160014, India

ABSTRACT

After interacting with local people of Kinnaur, nine wild vegetables viz. *Allium wallichii*, *Crambe kotschyana*, *Diplezium esculentum*, *Eremurus himalaicus*, *Fagopyrum tataricum*, *Lepyrodiclis holosteoides*, *Rorippa nasturtium-aquaticum*, *Silene vulgaris*, *Urtica dioica* were selected for proximate composition analysis, including mineral and vitamin content. Young green shoot tips and leaves of *Crambe kotschyana* are very rich in proteins (36.68 mg/100gm); *Rorippa nasturtium-aquaticum* in minerals (16.76 mg/100gm); and *Diplezium esculentum* in vitamin C (56.87mg/100gm) and fiber (6.89mg/100gm). The range of nutrients present in these vegetables is comparable with the commonly used cultivated green leafy vegetables. Infact, the protein (4.20- 36.28mg/100gm) and iron (3- 44.2 mg/100gm) content is much higher than that most of the cultivated vegetables. The present study should be helpful to educate the local youth about the nutritional value of this plant wealth of the region and create an interest in its conservation, collection and sustainable consumption.

KEYWORDS: Proximate composition; Wild vegetables; Kinnaur; Plant wealth, Sustainable.

*Corresponding author



GEETA

Department of Botany, Panjab university, Chandigarh 160014, India

INTRODUCTION

In many culture of India, plants have great ethnobotanical importance, whether it is nutritional, medicinal, and ritual or magical value. There are 1532 edible wild food species in India, mostly from Western Ghats and Himalayan regions (Arora and Pandey, 1996). Himachal Pradesh, a hilly State of Western Himalaya, is a repository of plant diversity and the traditional ethnobotanical knowledge. Because of varied altitudinal gradients and climatic conditions, the State harbours a rich plant diversity, including around 3400 species of flowering plants (Chowdhery, 1999). Utilization of plant resources in the day-to-day life has been an age old practice of the people inhabiting this hilly State. Many wild greens are eaten raw in salads or in boiled mixtures as 'blood cleansing' and 'intestine cleansing' agents. The wild vegetables provide adequate amount of crude fiber, fats, carbohydrates, proteins, water and mineral elements like Ca, Na, K, Fe, Mg, Mn, Cu, Zn etc, in addition to vitamins (Gogoi and Kalita, 2014). Kinnaura tribals are inhabitants of district Kinnaur of Himachal Pradesh and they mostly depend on wild plants for food, medicine, fuel, and house construction. The knowledge about wild plant used as food, vegetables and medicine has been handed down to them for generations, from father to son by means of oral tradition, coupled with observations in the field and at local gatherings of people. Mostly women and children go in groups to forests for the collection of wild foods, usually travelling great distances. However, this interest in the traditional collection of wild food plants is now fast declining due to loss of interest by younger generation of people who now have easy access to conventional vegetables. As a result, there is a complete ignorance among the younger generation about the nutritional and health benefits of the local wild food plants. The wild food plants score over conventional vegetables in that these do not require any formal cultivation, are inexpensive, have high quality nutritional value, and can help in improving malnutrition, so widespread amongst poor tribals. Tribal district Kinnaur being

remotely located, the supply of vegetables comes from far off distances. These vegetables are sold at much higher prices in the local market, while on the contrary the local wild vegetables come free, are easily available in the wild, and are organic and fresh. Integrating wild vegetables into diet has been promoted as the most practical and sustainable way to achieve optimal dietary requirements in many developing countries (Chadha and Oluoch, 2003). There is now growing evidence that traditional green leafy vegetables have higher nutritional value than several known common vegetables (Sundriyal and Sundriyal, 2001; Nordeide *et al.*, 1996; Orech *et al.*, 2007). Dark green vegetables are probably the most concentrated source of nutrition of any food. They are a rich source of minerals (including iron, calcium, potassium, and magnesium), vitamins (including vitamins K, C, E and many of the vitamins from B group), and a variety of phytonutrients (including beta-carotene, lutein, and zeaxanthin) which protect our eyes from age-related problems (such as age-related muscular degeneration) among many other benefits. Dark green leaves also contain small amounts of omega-3 fatty acids which protect us from cardiovascular diseases. The green leafy vegetables can, therefore, provide substantial nutritional and dietary benefits to tribal populations living in remote rural areas and can prevent several chronic diseases caused by malnutrition. The present study on local wild vegetables of District Kinnaur, Himachal Pradesh (Fig.1) was conducted with an aim to check their nutritional value as per the belief of local inhabitants and also to educate the local youth about the nutritional value of this plant wealth of the region and create an interest among the local youth in the conservation, collection and sustainable consumption of local wild vegetables. Assessment of nutritional composition and health benefit of wild food plants is fundamental for promoting these plants amongst the locals and the larger population; important for community health, agriculture and nutrition education initiatives; and important for

improving the management and conservation of local biodiversity.

MATERIALS AND METHODS

2.1 Ethnobotanical study

For the present study, local herbal healers, elderly men and women, and shepherds were consulted, with a view to gain knowledge about the wild vegetables, their nutritional value, health benefits, and medicinal value, if any. Many remote villages were visited by the senior author during this study for an interaction with people about the identity and uses of the various wild vegetable plants collected. A majority of respondents well recognized these plants and the information provided by them about these plants was noted on a specially designed questionnaire and verified by cross-checking with various people from different villages of the area. Different localities of Kinnaur district were visited three times every year, during the years 2012-2013. Information was collected from local sources about the various plants used as vegetables by them and also their medicinal importance, if any. The information collected about each plant included: local name(s) of the plant, part of the plant used, mode of cooking and consumption, and medicinal value, if any. Sampling and collection of plants used by locals were also done. The selected plants were photographed in their natural habitats. All the plants were collected in polythene bags for further study, and also 5-6 specimens of each species were dried and mounted on herbarium sheets for future records. The plants were tentatively identified with the help of keys and descriptions given in the relevant floras and these identifications were later confirmed by comparing with the authentically identified specimens lying in the Herbarium of Panjab University, Chandigarh (PAN); Herbarium of Northern Regional Centre, Botanical Survey of India (BSD); and the Herbarium of Forest Research Institute, Dehra Dun (DD).

2.2 Samples and analysis

Samples of young wild leafy vegetables were collected from the various localities of district

Kinnaur, varying in altitude from 2200-4000m. The collected vegetables were shade dried naturally and preserved for subsequent analysis of their proximate composition (moisture, ash, protein and fat), carbohydrates, dietary fibers, minerals (K, Zn, Fe), vitamin C, K and B₉ (folic acid) contents. Proximate analysis was carried out using standard method (AOAC, 2000). Total, soluble and insoluble dietary fibre constituents were determined by the enzymatic method of Furda (1981). Potassium content was measured by using a flame photometer. Determination of Iron and Zinc was done by Atomic Absorption Spectrophotometer (AAS), using the reference standard of the relevant metals. Vitamin content was determined by HPLC.

RESULTS

3.1 Local wild vegetables

A total of 9 species of wild plants, 8 belonging to 5 families of angiosperms and 1 pteridophyte, were found to be extensively used as vegetables by the locals, mainly in Pooh block area. It was discovered that *Diplezium esculentum*, locally called "lingra", is widely used in Nichar block and is also sold in the local market. Table 1 gives a list of these plants along with information about their botanical name, name of the botanical family, local/common English names, and the mode of consumption. The field photographs of all these plants are collected in figures 2 (a-i). Usually, only the young leaves of these plants are consumed by locals which are collected from the wild during the months of April- May. Besides being consumed fresh, young leaves and tender shoots of 6 species, viz. *Allium wallichii*, *Crambe kotschyana*, *Fagopyrum tataricum*, *Lepyrodiclis holosteoides*, *Silene vulgaris*, *Urtica dioica* are chopped, dried and stored for the winter season when conventional vegetables are not available and the local inhabitants depend entirely on these indigenous vegetables. Of these species, *Lepyrodiclis holosteoides*, *Silene vulgaris*, *Urtica dioica* and straw of *Fagopyrum tataricum* are also used as fodder for domesticated animals. All these vegetables grow wild while

Fagopyrum tataricum is found both as wild and also cultivated. All these vegetables are believed to increase and purify blood and also clean the digestive system.

3.2 Nutritional composition

The nutritional composition of the local wild vegetables examined in this study is presented in Table 2. A perusal of this table would indicate that all the vegetables are source of very high energy except only *Eremurus himalaicus*. *Eremurus himalaicus* has highest (74.47 %) and *Silene vulgaris* has lowest (13.29 %) moisture content. Protein content in various vegetables ranged from 4.20g/100g in *Eremurus himalaicus* to 36.68 g/100g in *Crambe kotschyana*, which is comparable to the common leafy vegetables (Table-3). These vegetables have a very low fat content which ranges from 0.12g/100g to 0.72g/100g. Three species, viz. *Diplezium esculentum* with 6.89g/100g, *Lepyrodiclis holosteoides* with 5.22g/100g, and *Crambe kotschyana* with 5.08g/100g fiber are most rich source of dietary fiber. The ash content ranged from 2.80g/100g (*Eremurus himalaicus*) to 16.76g/100g (*Rorippa nasturtium-aquaticum*). The carbohydrate content in the various species varied considerably, ranging from 4.07g/100g in *Eremurus himalaicus* to 38.86g/100g in *Allium wallichii*. All the species investigated presently contain remarkably high amounts of Potassium (< 1000mg/100g) except only *Eremurus himalaicus* (with 920mg/100g), *Urtica dioica* (with 860mg/100g), and *Lepyrodiclis holosteoides* (with 520mg/100g) which possess lesser amounts of this mineral . The zinc content ranged from 17.8 mg/100g in *Lepyrodiclis holosteoides* to 1.6 mg/100g in *Eremurus himalaicus*. The highest iron content was found to be 41.8 mg/100g in *Allium wallichii* and lowest was 3 mg/100g in *Crambe kotschyana* . The vitamin C content ranged from 62 mg/100 in *Lepyrodiclis holosteoides* to 41.51 in *Urtica dioica*. *Crambe kotschyana* had highest vitamin C content (8.51 mg/100g) while the lowest content of this vitamin was found in *Diplezium esculentum* (1.24 mg/100g). The folic acid content in various species ranged from

0.003 mg/100g in *Lepyrodiclis holosteoides* to 0.103 mg/100g in *Allium wallichii*.

DISCUSSION

The role of wild traditional vegetables as source of food and medicine has been documented by several workers (Ogle *et al.*, 2003; Lee *et al.*, 2003; Adebooye and Opabode, 2004; Ayodele, 2005). In the present study, it was found that the local wild plant species consumed as leafy vegetables contain significantly higher amounts of protein than the commonly cultivated leafy vegetables. Turan *et al.*, (2003) also reported more protein content in wild edible leaves consumed in Eastern Anatoliathan those of cultivated species such as spinach, lettuce, cabbage. The amount of iron is also higher than those of common leafy vegetables compiled by Hoe *et al.*, (1999). Ogle (2001) also reported higher iron content in wild vegetables as compared to the cultivated species. The amount of fiber, ash and carbohydrate was also found to be much higher than the common leafy vegetables as given by Hoe *et al.*, (1999). The results of present study shows that the local wild vegetables have all the essential nutrients essential for human diet for maintaining good health. All the wild leafy vegetables studied here (except only *Lepyrodiclis holosteoides*) also possess medicinal properties as reported by many workers (Kaul, 1997; Pieroni, 2000; Rivera *et al.*, 2005; Khare, 2007; Acharya *et al.*, 2011; Kaushik *et al.*, 2011; Khan *et al.*, 2013; Sayedet *et al.*, 2013) and therefore, these vegetables not only provides the essential nutrients to the consumers but also safeguard them against several ailments. Interestingly, the locals eat most of these green vegetable after frying in oil which makes available vitamin A in absorbable form to our body. This vitamin is a fat-soluble and hence putting oil as dressing on the leafy vegetables if consumed raw as salad, or cooking them in oil, makes this vitamin available in absorbable form. Vitamin K is a star among all the vitamins and a cup of most cooked green vegetable provides at least 9 times the minimum recommended intake of vitamin K. It was observed by the present authors that the knowledge about the uses of

local plants is more common with the locals aged above 35 years as compared to younger residents. Disease of modern world like diabetes and blood pressure were rare among aged local people who are still dependent on ethnic life style. Young adults are also no longer interested in collection of the wild leafy vegetables as conventional vegetables are easily available in the market. This lack of

interest in the local plant wealth by the younger generation can lead to complete loss of traditional knowledge of the local plant diversity. It is in this context that the present study shall help in popularising the importance of local plant diversity amongst the younger generation of local residents, besides creating awareness about the health benefits of these plants.

Table 1
List of local wild plants used as vegetable by local inhabitants.

Sr.No	Botanical name	Local/Common name	Family	Mode of consumption
1	<i>Allium wallichii</i> Kunth	Chiskan/ onion	Himalayan Liliaceae	Chopped leaves are fried in mustard oil and mixed with spices
2	<i>Crambe kotschyana</i> Boiss	Akharo	Brassicaceae	Chopped Leaves are boiled and then fried in mustard oil and mixed with spices
3	<i>Diplezium esculentum</i> (Retz.) Sw.	Lingda/ Paco	Athyriaceae	Fresh immature fronds are wiped with a cloth to remove red petiolar hairs. Then are boiled, cut and fried in mustard oil and mixed with spices
4	<i>Eremurus himalaicus</i> Boker	Prait/ Fox tail Lily	Liliaceae	Chopped leaves are fried in mustard oil and mixed with spices.
5	<i>Fagopyrum tataricum</i> (L.) Gautn.	Bro, Brass, Buck wheat	Fafra/ Polygonaceae	Chopped leaves are fried in mustard oil and mixed with spices
6	<i>Lepydiclis holosteoides</i> Fenz ex Fisch.	Latishaa	Caryophyllaceae	Leaves are fried in mustard oil, mixed with potato and spices.
7	<i>Rorippa nasturtium-aquaticum</i> (L.) Hayek	Tee-soma/ cress	Water Brassicaceae	Fresh leaves and potato are together fried in mustard oil and mixed with spices
8	<i>Silene vulgaris</i> (Moench) Garcke.	Cham- chom/ campion	Bladder Caryophyllaceae	Chopped leaves are fried in mustard oil, mixed with potato and spices.
9	<i>Urtica dioica</i> Linn.	Khorgya/ Nettle	Northern Urticaceae	Young leaves cooked mixed with <i>Silene vulgaris</i> and potato

Table 2
Nutritional composition of local wild vegetables investigated presently (per 100g dry leaves).

Plant name	Energy (Kcal)	Moisture (%)	Ash (%)	Fat (%)	Dietry fiber (%)	Protein (%)	Carbohydrate (%)	K (mg)	Zn (mg)	Fe (mg)	Vit C (mg)	Vit K (mg)	Folic Acid (mg)
<i>A. wallichii</i>	239.3	18.44	11.46	0.22	2.06	20.47	38.86	3160	1.7	41.8	41.84	4.29	0.103
<i>C. kotschyana</i>	225.06	16.64	12.55	0.22	5.08	36.68	19.09	2170	3.2	3	52.50	8.51	0.070
<i>D. esculentum</i>	237.6	16.55	10.27	0.72	6.89	21.02	36.76	1820	15	15	56.87	1.24	0.060
<i>E. himalaicus</i>	36.77	74.47	2.80	0.41	4.23	4.20	4.07	920	1.6	14.7	48.30	1.48	0.037
<i>F. tataricum</i>	226.04	16.80	13.94	0.24	3.76	26.81	29.16	2100	2.9	18.9	41.86	9.52	0.017
<i>L. holosteoides</i>	218.92	15.59	15.81	0.16	5.22	20.73	33.64	520	17.8	27.9	62.00	3.84	0.003
<i>R. nasturtium-aquaticum</i>	229.12	14.65	16.76	0.12	3.05	22.16	34.85	3220	6.5	18.8	48.22	2.87	0.063
<i>S. vulgaris</i>	235.91	13.29	15.50	0.19	2.64	30.24	28.31	6410	3.8	15.7	45.19	2.87	0.063
<i>U. dioica</i>	241.15	17.45	12.79	0.43	3.28	23.92	35.40	860	7.3	44.2	41.59	2.07	0.079

Table 3
Nutritional composition of some common cultivated leafy vegetables

Vegetable	Energy (Kcal)	Moisture (%)	Protein (%)	Fat (%)	CHO (%)	Fibre (%)	Ash (%)	K (mg)	Fe (mg)	Vit. C (mg)	Source of data
Cabbage (<i>Brassica oleraceavar. capitata</i>)	22	93.1	1.6	0.2	3.4	0.9	0.8	103	0.6	53.0	Hoe et al., (1999)
Bathu (<i>Chenopodium album</i>)	59	83	0.5	0.8	8.28	1.92	2.94	-	13	-	Odhav et al., (2007)
Cholai (<i>Amaranthus hybridus</i>)	268.92	83.48	17.92	4.65	52.18	8.61	13.80	54.20	13.58	25.40	Akubugwo et al., (2007)
Chinese mustard (<i>Brassica juncea</i>)	34	91.7	2.1	0.7	4.7	-	0.8	-	6.8	89.0	Hoe et al., (1999)
Lettuce (<i>Lactuca sativa</i>)	17	94.7	1.2	0.1	2.8	0.5	0.7	3545	1.5	27.6	Hoe et al., (1999)
Spinach (<i>Spinacea oleracea</i>)	29	91.3	2.6	0.3	4.0	-	1.8	136	0.8	56.4	Hoe et al., (1999)
Spring onion (<i>Allium fistulatum</i>)	19	95.0	0.6	0.1	3.9	-	0.4	-	0.4	17.0	Hoe et al., (1999)

Figure 1
a) Map of India; b) Himachal Pradesh; c) Study area- tribal district Kinnaur.



Figure 1.

Figure 2
Field photographs of the species studied presently: a) *Diplezium esculentum*; b) *Urtica dioica*; c) *Lepyrodiclis holosteoides*; d) *Rorippa nasturtium- aquaticum*; e) *Silene vulgaris*; f) *Fagopyrum tataricum* (cultivated); g) *Allium wallichii*; h) *Crambe kotschyana*; i) *Eremurus himalaicus*.



Figure 2.

CONCLUSION

Indigenous communities in Kinnaur district nurture rich knowledge about edible and medicinal plants acquired over several generations of experimentation. All such plants possess nutrients comparable with common leafy vegetables and hence have the potential to fulfil the nutritional requirements of local inhabitants. Also, since the traditional knowledge about the importance of local plants

is usually passed on to the next generation just by the word of mouth, this knowledge not only needs to be preserved in written records but also substantiated by modern nutritive and pharmacological experiments for a scientific validation and use as safe healthcare remedies.

ACKNOWLEDGEMENT

The senior author wishes to acknowledge CSIR for providing the fellowship for the study and all

the interviewees for their hospitality and willingness to share their valuable traditional knowledge during the field survey. We are also thankful to the Chairperson, Department of Botany for laboratory facilities.

CONFLICT OF INTEREST

Conflict of interest declared none.

REFERENCES

1. Acharya K, Pokhriol H, Gyawali R., Comparative study of antibacterial and cytotoxic activity of two Nepalese medicinal plants- *Allium wallichii* Kunth and *Allium sativum* L. *International Journal of Pharmaceutical and Biological Archive*, 2(5):1539-1543, (2011).
2. Adebooye OC, Opabode JT., Status of conservation of the indigenous leafy vegetables and fruits of Africa. *African Journal of Biotechnology*, 3: 700–705, (2004)
3. Akubugwo IE, Obasi NA, Chinyere GC, Ugbogu AE., Nutritional and chemical value of *Amaranthushybridus* L. leaves from Afikpo, Nigeria. *African journal of Biotechnonology*. 6(24): 2833-2839, (2007).
William Horwitz, Ed. Official Methods of Analysis of AOAC International, 17th ed. Gaithersburg, MD publisher. (2000)
4. RK Arora and A. Pandey, Ed. Wild edible plants of India, diversity, conservation and use. Botanical survey of India, Calcutta, India, (1996).
5. AE Ayodele. "The medicinally important leafy vegetables of south western Nigeria". *Ethnobotanical leaflets*. Web journal <http://www.siu.edu/~ebl/leaflets/ayodele.htm>, (2005).
6. Chadha ML and Oluoch MO., Home-based vegetable gardens and other strategies to overcome micronutrient malnutrition in developing countries. *Food, Nutrition and Agriculture Series*. 32, (2003).
7. H.J. Chowdhery. Floristic diversity and conservation strategies in India. Himachal Pradesh. In: Mudgal V and Hajra P K (eds). Vol II: in the context of states and union territories, (BSI, Calcutta), 845-894, (1999).
8. I. Furda. Simultaneous analysis of soluble and insoluble dietary fiber. In: W.P.I. James and O. Theander (eds.), *The analysis of dietary fiber in food*, Marcel Dekker, Inc., New York (1981).
9. Gautam AK, Bhatia MK, Bhadauria R., Diversity and usage custom of plants of South Western H.P., India. *Journal of Phytology*. 3(2): 24-36, (2011).
10. Gogoi P and Kalita JC., Proximate analysis and mineral components of some edible medicinally important leafy vegetables of Kamrup district of Assam, India. *International Journal of Pharma and Bio Science*. 5(4): 451-457, (2014).
11. Hoe VB and Siong KH., The nutritional value of Indigenous fruits and vegetables in Sarawak. *Asia Pacific J Clin Nutr*. 8(1): 24-31, (1999).
12. Kala CP and Manjrekar N., Ethno-medico-botany of Indian trans Himalaya: A case study from Spiti. *Journal of Economic and Taxonomic Botany*, 23(1): 177- 183, (1999).
13. Kaul MK., Ed. *Medicinal plants of Kashmir & Ladakh: Temperate & cold arid Himalayas*, (1sted.). Indus Publishing Company, New Delhi, 44-85, (1997).
14. Kaushik A, Kaushik JJ, Das A, Gemal S, Gaim D., Preliminary studies on anti-inflammatory activities of *Diplazium esculentum* in experimental animal models. *International Journal of Pharmaceutical Sciences and Research*. 2(5): 1251-1253, (2011).
15. Khan SM., Page S., Ahmad H., Harper DM. Medicinal flora and ethnoecological knowledge in the Naran valley, Western Himalaya, Pakistan. *Journal of Ethnobiology and Ethnomedicine*. 9(4) (2013).

16. Khare CP., Ed. Indian Medicinal Plants: an illustrated Dictionary. Springer Science and Business Media, LIC., (2007).
17. Lee Y, Cesario T, Wang Y, Shanbrom E, Thrupp L., Antibacterial activity of vegetables and juices. *Nutrition*, 19: 994–996, (2003).
18. Odhav B, Beekrum S, Akul U, Baijnath H., Preliminary assessment of nutritional value of traditional leafy vegetable of Kwa Zulu- Natal, South Africa. *Journal of food composition and analysis*, 20: 430-435, (2007).
19. Ogle BM, Tuyet HT, Duyet HN, Xuan D, Nguyen N., Food, feed or medicine: the multiple functions of edible wild plants in Vietnam. *Economic Botany*. 57: 103–117, (2003).
20. Ogle B. Wild vegetables and micronutrient nutrition. PhD Thesis, Uppsala University, Sweden, (2001).
21. Orech FO, Aagaard-Hansen J, Friis H., Ethnoecology of traditional leafy vegetables of the Luo people of Bondo district, Western Kenya. *Int. J. Food Sci. Nutr.* 58(7): 522-530, (2007).
22. Pal MM, Janifer RX, Phani KG, Gupta S, Bala SS., Phytofoods of Nubra Valley, Ladakh- The cold desert. *Indian Journal of Traditional Knowledge*, 9(2): 303-308, (2010).
23. Pieroni A., Medicinal plants and food medicine in the folk traditions of the upper Lucca Province Italy. *Journal of Ethnopharmacology*, 70: 235-273, (2000).
24. Rivera D, Obon C, Heinrich M, Llorach R., The ethnobotanical study of local Mediterranean food plants as medicinal resource in Southern Spain. *Journal of Physiology and Pharmacology*, 56(1): 97-114, (2005).
25. Rutto LK., Xu Y., Ramirez E., Brandt MM., Mineral Properties and Dietary Value of Raw and Processed Stinging Nettle (*Urtica dioica* L.). *International Journal of Food Science*, 9, (2013).
26. Sarin YK., A survey of vegetable raw material Resources of Lahaul. *Indian Forester*. 93(7): 489-499, (1967).
27. Sayed MJ, Mohmad PW., Ethnobotany of some polypetalous plants from the Kashmir Himalaya. *Journal of Medicinal Plants Research*. 7(36): 2714-2721, (2013).
28. Singh KN and Lal B., Ethnomedicines used against four common ailments by the tribal communities of Lahaul-Spiti. *Journal of Ethnopharmacology*, 115: 147-159, (2008).
29. Thakur Shailza., Medicinal plants used by tribal inhabitants of Sirmourdistric, Himachal Pradesh. *Indian J. Sci. Res.* 2(4): 125-127, (2011).
30. Turan M, Kordali S, Zengin H, Dursun A, Sezen Y., Macro and micro mineral content of some wild edible leaves consumed in Eastern Anatolia. *Acta Agriculturae Scandinavica. Section B. Plant and Soil Science*. 53: 129–137, (2003).
31. Uprety Y, Poudel RC, Shrestha KK, Rajbhandary S, Tiwari NN, Shrestha UB, Asselin H., Diversity of use and local knowledge of wild edible plant resources in Nepal. *Journal of Ethnobiology and Ethnomedicine*. 8: 16, (2012).