



## *Internationally indexed journal*

Indexed in Chemical Abstract Services (USA), Index copernicus, Ulrichs Directory of Periodicals, Google scholar, CABI ,DOAJ , PSOAR, EBSCO , Open J gate , Proquest , SCOPUS , EMBASE ,etc.



### *Rapid and Easy Publishing*

*The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quarterly. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences*



#### **Pharmaceutical Sciences**

- Pharmaceutics
- Novel drug delivery system
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics



#### **Biological Sciences**

- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics
- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

**Indexed in Elsevier Bibliographic Database  
(Scopus and EMBASE)**

**SCImago Journal Rank 0.288**

**Impact factor 2.958\***

Chemical Abstracts  
Service ([www.cas.org](http://www.cas.org))



A division of the American Chemical Society

**CODEN IJPBJ2**



## Elsevier Bibliographic databases (Scopus & Embase)

**SNIP** value – 0.77

**SJR** - 0.288

**IPP** - 0.479

*SNIP – Source normalised impact per paper*

*SJR – SCImago Journal rank*

*IPP – Impact per publication*

*Source – [www.journalmetrics.com](http://www.journalmetrics.com)*

*(Powered by scopus (ELSEVIER))*



**LUND**  
UNIVERSITY



JACKSONVILLE STATE UNIVERSITY

Jacksonville State University  
Houston Cole Library  
USA (Alabama)



UNIVERSITY OF  
OXFORD

Oxford, United Kingdom

INDEX COPERNICUS  
INTERNATIONAL

*And indexed/catalogued in  
many more university*



\*Instruction to Authors visit [www.ijpbs.net](http://www.ijpbs.net)

For any Queries, visit "contact" of [www.ijpbs.net](http://www.ijpbs.net)



## ROLE OF MYCORRHIZAE AS BIOFERTILIZER AND BIOPROTECTANT

**S.K. DWIVEDI, SANGEETA\* AND RAM GOPAL**

*Department of Environmental Science, Babasaheb Bhimrao Ambedkar (A Central) University,  
Raebareli Raod, Lucknow-226025, Uttar Pradesh, India.*

### ABSTRACT

Mycorrhizae are obligate fungi that predominate in the roots and soil of higher plants. They form association with plant roots in a host-non specific manner. Seven types of mycorrhizae have been known i.e. arbuscular, ecto, arbutoid, ectendo, ericoid, monotropoid and orchidaceous mycorrhizae. Out of these, arbuscular and ectomycorrhizae are the most abundant and wide spread. They promote plant growth by enhancing nutrient acquisition and promoting growth hormones. They also increase the resistance in plants against plant pathogens and surface area of root system for better absorption of nutrient from soil. Therefore, they can be used as biofertilizer and as biocontrol agent. This article presents an overview of current knowledge on mycorrhiza and their potential benefits to agriculture as biofertilizer and biocontrol agent.

**KEYWORDS:** Biocontrol agent; Biofertilizer; Plant growth promoter; Soil-borne pathogens; Vesicular-arbuscular mycorrhizae

\*Corresponding author



**SANGEETA**

Department of Environmental Science, Babasaheb Bhimrao Ambedkar (A Central)  
University, Raebareli Raod, Lucknow-226025, Uttar Pradesh, India.

## 1. INTRODUCTION

India is an agricultural based country and in order to feed the increasing population, it has to increase its productivity. After green revolution the crop productivity in India has increased many folds. In last few years the crop yield has increased due to the use of inorganic fertilizers and pesticides (51). Now, agricultural practices totally depend on the chemical fertilizer and pesticides for the productivity of various crops. Use of nitrogen, phosphorus and potassium based fertilizers increase the crop productivity to some extent but, on the other hand, increased doses of chemical fertilizers posed devastating effect on soil fertility, cause environmental pollution and kill the beneficial microorganisms which are present in soil. The indiscriminate use of pesticides has induced resistance in pests against these pesticides. Moreover, the toxic chemicals which are applied in the field are absorbed by plants and enter the food chain through vegetables and cereals causing many health problems. So, in order to cope up with all these problems and make agriculture sustainable development, we should focus on such certain agricultural practices which are ecofriendly. The concept of sustainable development is the management of natural resources for human use and is one of the best ways for the sustainable agriculture and to decrease the negative impact of pesticides and fertilizers on environment (2). Sustainable agriculture means full utilization of environmental resources with no harm to it (61).

One such approach towards sustainable agricultural practice is to move towards the microbial technology such as use of *Mycorrhizae* as biofertilizer and biocontrol agent (27). In agricultural ecosystem, they play a vital role in nutrients fixing, solubilizing and mobilizing. Mycorrhizae have profound effect on the rhizosphere. They alter the microbial community structure of the soil (57). They interact with other soil biota such as phosphate solubilizing bacteria, Plant growth promoting rhizobacteria, plant pathogens and other bioagents, and microfauna which results in a significant positive or negative effects on plant growth (47). Mostly positive effects on plant growth have been reported, although numerous negative effects on plant growth have also been reported (68). Arbuscular mycorrhizal (AM) fungi play a significant role in sustainable farming system because AM fungi are efficient when nutrient availability is low and nutrients are bound to organic matter and soil particles. They directly or indirectly affect plant growth (Fig. 1). Indirectly they promote plant growth by improving the soil quality and by suppressing the pathogens responsible for reduced crop production (78). Mycorrhizae are important for plants and ecosystem. They affect the plant production and soil health (24; 62; 82; 63; 29). AM fungi colonize the roots of many economically important crops and could serve as biofertilizer and bioprotectors in environmentally sustainable agriculture (20).




Figure 1

**Diagrammatic representation showing role of Mycorrhizae as biocontrol and biofertilizers**

## 2. Mycorrhizae

The term mycorrhizae was first time introduced by A.B. Frank in 1885. It literally means “fungal root” used to denote the association between certain soil fungi and plant roots. Mycorrhiza is a type of endophytic, biotrophic, symbiotic, mutualistic relationship special soil fungi and fine plant root (<http://edis.ifas.ufl.edu>). The most ancient and abundant mutualistic on earth is found between majority of land plants and arbuscular mycorrhizal (AM) fungi by forming extensive hyphal network in soil and plants (64). Fossil records indicate that it is evolved 400-450 million years ago (65). It is neither the fungus nor the root, but it is the structure formed by these two partners. Fungus increase the plant nutrition absorption capacity from the soil by increasing the surface area of root system and in exchange fungus receives carbohydrates and growth factors from the plant (<http://edis.ifas.ufl.edu>). In mycorrhizal interaction bidirectional exchange of resources across the mycorrhizal interface takes place. The mycorrhizae provides nutrients (phosphate and nitrogen) to host plants and increases the

abiotic (drought, salinity and heavy metals) and biotic stress (root pathogens) resistance in host plants and in return the host plant provides 4-20% of its photosynthetically fixed carbon to the mycorrhizae (81). There are certain plants which have lost their photosynthetic property and parasitize mycorrhizae that are associated with neighbour autotrophic plants. Such type of plants are mycoheterotrophic plants (20). Vesicular-arbuscular mycorrhizal (VAM) fungi are ubiquitous in nature and act as a natural link between the roots of most plants and soil (52). They form an intimate relationship with plant root system and benefit plant health (17) by contributing to phosphorus uptake (66; 33) and nitrogen uptake (30). Besides these they also play an important role in uptake of K, S, Mg, Fe, Zn, Cu (59). They contribute up to 90% of plant P demand (79). Arbuscular mycorrhizal spores can germinate in the absence of host plant but its hyphal branching and metabolic activity increase only in the presence of strigolactones (root exudates) that induces the pre-symbiotic growth of AM fungal spores (3).




Figure 2

(a) root without Mycorrhizae, (b) root with Endomycorrhizae, (c) root with Ectomycorrhizae

There are three major groups of Mycorrhizae: *Ectomycorrhiza* (Fig 2c), *Ectendomycorrhiza* and *Endomycorrhiza* (Fig 2b), out of these *Ectomycorrhiza* and *Endomycorrhiza* play a vital role in agriculture. *Ectotrophic* types or *Ectomycorrhizal* fungus penetrates between the

cell walls of the cortex and forms a covering sheath of fungal hyphae around the entire root system. Whereas *Endotrophic* types or *Endomycorrhizae* or *Arbuscular Mycorrhizae* do not form mantle or covering sheath of fungal hyphae over the root, they just enter the cortex

cells (<http://edis.ifas.ufl.edu>). The fungi of Glomeromycota (AM fungi) form a unique structure such as arbuscules and vesicles known as arbuscular mycorrhizae (AMs). Out of seven mycorrhizae i.e. arbuscular, ectendo, arbutrid, monotropoid, ericoid and orchidaceous mycorrhizae, the arbuscular and ectomycorrhizae are the most abundant and

widespread (58). The VAM fungi are the most common mycorrhizal type which are involved in agricultural system (16). *Ectomycorrhizae* are found in the roots of pines, birches, willows and oaks and *Endomycorrhizae* are present in most agronomic vegetable crops and fruits (<http://edis.ifas.ufl.edu>).

**Table1**  
**Direct and indirect effects of mycorrhizal fungi on crop productivity**

Direct effects on crops	Indirect effects on crops
Stimulation of plant productivity	Weed suppression
Nutrient acquisition (P, N, Cu, Fe, Zn)	Stimulation of nitrogen fixation by legumes
Enhanced seedling establishment	Stimulation of soil aggregation and soil structure
Drought resistance	Suppression of some soil-borne pathogens
Heavy metal resistance	Stimulation of soil biological activity
	Increased soil carbon storage
	Reduction of nutrient leaching

(Source: van der Heijden et al., 2008)

AM fungi can enhance plant growth (15), affect the cellular and biochemical compositions (44; 37) and suppress plant disease (85). These can also increase the bacterial density in soil, produce siderophore and reduce nematode population in mycorrhizosphere of plants. During the last few decades the government and entrepreneurs have paid special attention towards the setting up of biofertilizers production facilities due to which production and promotion of biofertilizer in agriculture has increased. Now the farmers have realized the benefits of biofertilizers. The role of biofertilizer like AM fungi in the growth and multiplication of crop plant can prove to be the most effective alternative to chemical fertilizers for enhancing growth and biomass production of crop. In addition, it can also improve vigour and nutrient uptake, disease resistance and drought tolerance (76). It has been reported that phosphate solubilizing bacteria (PSB) interact with VAM and release phosphate ions in the soil which have a synergistic effect on P solubilization (48) because the phosphate solubilized by the bacteria can be more efficiently absorbed by plants through the bridge formed by mycorrhizae among the roots and surrounding soil (38). Moreover, it has also been found that PSB and VAM in combination improved mineral (N and P) accumulation in


plants tissues (77). In addition, it has been reported that AM fungi could mobilize phosphorus from rock phosphate (12).

### 2.1. Mycorrhizal fungi as biofertilizer

AM fungi directly influence the growth of its host plant such, for instance, by providing nutrition through plant roots (65). Moreover, some extracellular secretions from fungal hyphae (50) directly affect microflora of the mycorrhizosphere and also play a very significant role in soil aggregation (53). VAM fungi have some special structures known as vesicles and arbuscules. They increase the length of the root 100 fold of the normal because of which the root reach up to wetter soil and help plants to absorb many less available nutrients such as phosphorus, zinc, molybdenum and copper. Some VAM fungi form a covering sheath around the root; they increase the tolerance of seedling to drought, high temperature, pathogens and soil acidity. The greatest growth response to mycorrhizal fungi is found in plants of highly weathered tropical acid soil where P is less and has high toxic value due to presence of Aluminium. The microbiologically solubilized phosphate is taken up by mycorrhizal mycelium which develops a synergistic microbial interaction (14; 25; 21). Besides this, it also stimulates foliar content

(25; 21). Mycorrhizal symbiosis plays a very important role in nutrient cycling in agricultural and natural ecosystem (49; 11). It also increases the plant resistance to different environmental stress (11). The microorganisms present in mycorrhizosphere affect the AM formation and its function (11). The mycorrhizosphere has two zones i.e. mycorrhizosphere and mycosphere. Mycorrhizosphere is the zone which is under the influence of root and fungal partners of the

mycorrhiza whereas mycosphere is the zone which supports bacterial activities (7). Microorganisms in the mycosphere of the AM fungi affect mycorrhizae functions such as nutrients and water uptake carried out by the external hyphae of AM fungi (25) and they are called as 'mycorrhiza helper bacteria' (MHB). These stimulate the growth of mycorrhiza (31) in the soil. The term was coined by Duponnois and Gerbaye (26).



**Figure 3**  
**Nutrient uptake by Mycorrhizae**

Mycorrhizal plants can absorb nutrients by two pathways. Firstly by absorbing nutrients directly from the soil through 'plant pathway', but due to this, a nutrient depletion zone is formed around the root. In nutrient deficient situation 'mycorrhizal pathway' is useful where extraradical mycelium (ERM) grows into the soil, uptake the nutrient and transport it to hartig

net in ectomycorrhizal (ECM) interaction or to intraradical mycelium (IRM) (55). The extraradical mycelium absorbs nutrients from the soil and transfers it to the host root whereas in second pathway the intraradical mycelium releases nutrients into interfacial apoplast and exchange them against carbon from the host (20). The use of rock phosphate as a cheap

alternative source has been used for sustainability purposes (86). Phosphorus is a very important element which is required by the plants for its metabolism and soil microbiological processes. They are present in soil in insoluble form and so, they are unavailable to plants. But AM fungi have the capability to convert insoluble P into soluble (ortho-phosphate) form. They play a very significant role in phosphate solubility through different mechanisms such as production and secretion of organic acids (34). Moreover, under P limiting conditions the external hyphae of AMF extend beyond the P depletion zone and provide nutrients from undepleted zones. It also helps in the formation of lateral roots and root branching which increases the surface area about 100 fold for the absorption of soil nutrients (56). AMF not only dissolves insoluble P but also produce some extracellular phosphatases that help in the mobilization of P from organic sources to plants (74). Besides P, AMF are also responsible for the uptake of nitrogen and sulphur. Sulphur is taken up by AMF in the form of amino acids i.e. cysteine and methionine (6) whereas N is taken in the form of nitrate ( $\text{NO}_3^-$ ) and ammonium ( $\text{NH}_4^+$ ) (19; 71; 73). It also helps in organic nitrogen uptake from soil (65). Increased concentration of both phosphorus and nitrogen in the plant biomass was reported when soybean was inoculated with AMF (36). Sabannavar and Lakshman (54) reported that the use of AMF alongwith *Pseudomonas striata* helps in solubilization of the mineral phosphate and contribute to biogeochemical P cycling, thus promoting sustainable nutrient supply to crop plants for higher yield. Similarly, Yousefi et al. (84) reported that the combined application of phosphate solubilizing bacteria and AMF increased the shoot dry matter yield, seed grain spike number and grain yield. In the same way synergistic relationship between phosphate solubilizing bacteria and AMF have been reported by Baquall and Das (13).

## 2.2 Mycorrhizal fungi as biocontrol agent

There are many disease management methods such as crop rotation, use of resistant varieties and chemical pesticides. However, frequent

and indiscriminate use of these pesticides affects the physical and chemical property of the soil. It also affects the non-target organisms and has developed resistance among the pathogen against these chemicals (8). Biocontrol potential of AM fungi against various phytopathogens is well documented (60; 39; 10; 18; 83; 70; 69). Arbuscular Mycorrhizal Fungi (AMF) are the major component of the rhizosphere of most of the plants and play a very important role as biocontrol agent and help in decreasing plant disease incidence (4). Application of combined VAM and Biological Control Agents (BCAs) reduced the growth of *Fusarium solani* and *Machrophomina phaseolina*, the causal agent of root-rot disease of *Geranium* plants. They had a synergistic effect on disease control (80). Thygesen et al. (75) reported that *Glomus claroideum* and *G. intraradices* induced tolerance in pea against *Aphanomyces euteiches* causing pea root-rot. Mixture of *Glomus intraradices*, *G. mosseae*, *G. clarum*, *Gigaspora margarita* and *G. gigantean* affects the growth of *Sclerotium cepivorum* and significantly reduced the white rot disease in onion when applied in green house and in field experiment (28). Later on Al-Askar and Rashad (5) used the mixture of *Glomus mosseae*, *G. intraradices*, *G. clarum*, *Gigaspora gigantean* and *G. margarita* against *Fusarium* root-rot disease of bean and reported that they significantly reduced the percentage of disease severity and incidence in infected bean plants. Several AM fungi have been reported to have ability to control soil-borne pathogens such as species of *Aphanomyces*, *Cylindrocladium*, *Fusarium*, *Macrophomina*, *Phytophthora*, *Pythium*, *Rhizoctonia*, *Sclerotium* and *Verticillium* (35). *Glomus mosseae* reduced the growth of pathogen *Gaeumannomyces graminis* var. *tritici* (40). Matsubara et al. (43) reported *Glomus fasciculatum* and *Gigaspora margarita* to be antagonistic against *Fusarium oxysporum* f. sp. *asparagi*, the causal agent of root rot disease of *Asparagus*. Similarly, *Glomus clarum* decrease the growth of the pathogen of root rot necrosis in cowpea i.e. *Rhizoctonia solani* (1). Some mycorrhizae as biocontrol agent against several pathogens are presented in Table 2.

**Table 2**  
**Some examples of Mycorrhizae as biocontrol agent**

Brinjal	<i>Fusarium solani</i>	<i>Glomus fasciculatum</i> and Salicylic acid (0.5 and 1.0 mM)	Ojha et al. (46)
Cotton	<i>Verticillium wilt</i>	<i>Glomus etunicatum</i>	Kobra et al. (41)
Sesame	Wilt and root-rot/ <i>Fusarium oxysporum</i> f.sp. <i>sesame</i> (Zap.) cast and <i>Macrophomina phaseolina</i> (Moub) Ashby	<i>Glomus</i> spp. and <i>Lums</i> spp. <i>Trichoderma viride</i> or <i>Bacillus subtilis</i>	Ziedan et al. (87)
Bean	Root-rot / <i>Fusarium solani</i>	Arbuscular Mycorrhizal Fungi (AMF) <i>Glomus mosseae</i> , <i>G. intraradices</i> , <i>G. clarum</i> , <i>Gigaspora gigantea</i> and <i>Gigaspora margarita</i>	Al-Askar and Rashad (5)
Apple	Stem brown canker or <i>Botryosphaeria canker</i>	Arbuscular mycorrhizal fungi	Krishna et al. (42)
Tomato	<i>Fusarium wilt</i> / <i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i>	Fluorescent <i>Trichoderma harzianum</i> and <i>Glomus intraradices</i>	<i>Pseudomonas</i> , Srivastava et al. (67)
Bean	<i>Sclerotinia sclerotiorum</i> (Lib.) de Bary	Arbuscular Mycorrhizal Fungi (AMF) <i>Glomus mosseae</i> (Gm), <i>Glomus fasciculatum</i> (Gf) and <i>Rhizobium leguminosarum</i> biovar <i>phaseoli</i> (Rlp)	Aysan and Demir (9)
Bean	Dry root-rot / <i>Macrophomina phaseolina</i>	AM fungi viz., <i>Glomus fasciculatum</i> , <i>G. mosseae</i> , <i>G. aggregatum</i> , <i>G. claroideum</i> , <i>G. macrocarpum</i> , and <i>G. multicaule</i>	Chandra et al. (22)
French Bean	Root-rot/ <i>Rhizoctonia solani</i>	AM fungi, <i>Pseudomonas fluorescens</i>	Neeraj and Singh (45)
Soybean	Red crown rot/ <i>Cylindrocladium parasiticum</i>	<i>Rhizobia</i> and AM fungi	Gao et al. (32)
Chilli	<i>Meloidogyne incognita</i>	<i>Pasteuria penetrans</i>	Chaudhary and Kaul (23)
Oil palm ( <i>Elaeis guineensis</i> )	Basal stem rot/ <i>Ganoderma boninense</i>	<i>Glomus intraradices</i> UT126, <i>Glomus clarum</i> BR152B and <i>Pseudomonas aeruginosa</i> UPMP3	Sundram et al. (72)

### 3. CONCLUSION

Mycorrhizal fungi play a very significant role in sustainable agriculture which is the urgent need of modern era. Mycorrhizal fungi act as an effective biofertilizer and bioprotectant. These improve the plant vigour and soil quality. They also play a very crucial role in plant nutrient (i.e. N, P, K, S, Mg, Fe, Zn, Cu) uptake, diversity and productivity of plant, biotic (root pathogen) and abiotic (drought, salinity, heavy metal) stress resistance, microbial diversity and population in mycorrhizosphere, soil aggregation and soil structure, reduction of nutrient

leaching, weed suppression and stimulates soil biological activity. Besides all these, mycorrhizal fungi also act as bioagent and protect the host plant root from various soil-borne plant pathogens such as bacteria, fungi and nematodes by using physical, chemical and biological mechanisms.

### ACKNOWLEDGEMENT

The authors are thankful to the Head, Department of Environmental Science, B.B.A.University (A Central University), Lucknow for providing the necessary facilities.

## REFERENCES

1. Abdel-Fattah G.M., Shabana Y.M. Efficacy of the arbuscular mycorrhizal fungus *Glomus clarum* in protection of cowpea plants against root rot pathogen *Rhizoctonia solani*. J. Plant Dis. Prot., 109: 207-215, (2002).
2. Adesemoye A.O., Torbert H.A., Kloepper J.W.. Plant growth promoting rhizobacteria allow reduced application rates of chemical fertilizers. Microb Ecol 58:921-929, (2009).
3. Akiyama K., Hayashi H. Strigolactones: Chemical Signals for Fungal Symbionts and Parasitic Weeds in Plant Roots. Annals of Botany 97: 925-31, (2006).
4. Akthar M.S., Siddiqui Z.A. Arbuscular Mycorrhizal Fungi as Potential Bioprotectants against Plant Pathogens. In: Mycorrhizae; Sustainable Agriculture and Forestry, Siddiqui, ZA, MS Akhtar and K Futai (Eds.). Springer Netherlands, Dordrecht, The Netherla. (2008).
5. Al-Askar A.A., Rashad Y.M. Arbuscular Mycorrhizal Fungi: A Biocontrol agent Common Bean *Fusarium* Root Rot Disease. Plant Pathology Journal 9(1):31-38, (2010).
6. Allen J.W., Shachar-Hill Y. Sulfur transfer through an arbuscular mycorrhiza. Plant Physiol. 149:549–560, (2009).
7. Andrade G., Mihara K.L., Linderman R.G., Bethlenfalvay G.J. Soil aggregation status and rhizobacteria in the mycorrhizosphere. Plant and Soil, 202: 89-96, (1998).
8. Arwry T.A., Quandt S.A. Pesticides at work and at home: Exposure of migrant farmworkers. Lancet, 362: 2021-2021, (2003).
9. Aysan E., Demir S. Using Arbuscular Mycorrhizal Fungi and *Rhizobium leguminosarum* Biovar *phaseoli* Against *Sclerotinia sclerotiorum* (Lib.) de Bary in the Common Bean (*Phaseolus vulgaris* L.). Plant Pathology Journal 8: 74-78, (2009).
10. Azcón-Aguilar C., Jaizme-Vega M.C., Calvet C. The contribution of Arbuscular Mycorrhizal Fungi to the control of soil-borne plant pathogens. In: Gianinazzi S, Schuepp H, Barea JM, Haselwandter K (eds) Mycorrhizal technology in agriculture. Birkhauser, Switzerland, pp 187–197, (2002).
11. Azcón-Aguilar C., Barea J.M. Applying mycorrhizal biotechnology to horticulture: significance and potentials. Scientia Horticulture, 68: 1-24, (1997).
12. Ba A.M., Guissou T. Rock phosphate and vesicular-arbuscular mycorrhiza effects on growth and nutrient uptake of *Faidherbia albida* (Del.) seedlings in an alkaline sandy soil. Afroforestry system, 34: 129-137, (1996).
13. Baquall M.F., Das M.F. Influence of Biofertilizers on Macronutrient uptake by the Mulberry Plant and its Impact on Silkworm Bioassay. Caspian J. Environmental Sci., 4(2): 98-109, (2006).
14. Barea J.M., Azcón-Aguilar C., Azcón. R. Interactions between mycorrhizal fungi and rhizosphere microorganisms within the context of sustainable soil-plant systems. In A.C. Gange and V.K. Brown (ed.), Multitrophic interactions in terrestrial systems. Blackwell Science, Cambridge, England, 1997, pp: 65-77.
15. Bayozen A., Yildiz A. Determination of mycorrhizae interactions and pathogenicity of *Rhizoctonia solani* Kuhn isolated from strawberry and *Xanthium strumarium*, Turkish J. Biol. 33: 53-57, (2009).
16. Bethlenfalvay G.J. Mycorrhizae in the agricultural plant-soil system. Symbiosis 14:413-425, (1992).
17. Bethlenfalvay G.J., Svejcar A.J. Mycorrhizae in plant productivity and soil conservation. In: Proceedings of the IV th International Rangeland Congress, Montpellier, France, p 251-254, (1991).
18. Bodker L., Kjoller R., Kristensen K., Rosendahl S. Interactions between indigenous Arbuscular Mycorrhizal Fungi and *Aphanomyces euteiches* in field-grown pea. Mycorrhiza 12:7–12, (2002).
19. Brandes B., Godbold D.L., Kuhn A.J., Jentschke G. Nitrogen and Phosphorus Acquisition by the Mycelium of the Ectomycorrhizal Fungus *Paxillus involutus* and its Effect on Host Nutrition. New Phytologist 140: 735-43, (1998).
20. Bücking H., Liepold E., Ambilwade P. The Role of the Mycorrhizal Symbiosis in Nutrient Uptake of Plants and the Regulatory Mechanisms Underlying These

- Transport Processes, Plant Science, Dr. Nabin Kumar Dhal and Sudam Charan Sahu (Ed.), InTech Publisher, Croatia, 2012, pp 107-138.
21. Caravaca F., Alguacil M.M., Azcon R., Parlade J., Torres P., Roldan A. Establishment of two ectomycorrhizal shrub species in a semiarid site after *in situ* amendment with sugar beet, rock phosphate and *Aspergillus niger*. Microb. Ecol., 49: 73-82, (2005a).
  22. Chandra S., Khare V., Kehri H. K. Evaluation of Arbuscular Mycorrhizal Fungi against *Macrophomina phaseolina* causing dry root-rot of urd and mung bean. Indian Phytopathology 60(1):42-47, (2007).
  23. Chaudhary K.K., Kaul R.K. Efficacy of *Pasteuria penetrans* and Various Oil Seed Cakes in Management of *Meloidogyne incognita* in Chilli pepper (*Capsicum annum* L.), J. Agr. Sci. Tech. Vol. 15: 617-626, (2013).
  24. Currie A.F., Murray P.J., Gange A.C. Is a specialist root-feeding insect affected by Arbuscular Mycorrhizal Fungi? Applied Soil Ecology 47: 77-83, (2011).
  25. Duponnois R., Colombet A., Hien V., Thioulouse J. The mycorrhizal fungus *Glomus intraradices* and rock phosphate amendment influence plant growth and microbial activity in the rhizosphere of *Acacia holosericea*. Soil Biol. Biochem., 37: 1460-1468, (2005).
  26. Duponnois R., Garbaye J. Effect of dual inoculation of Douglas fir with the ectomycorrhizal fungus *Laccaria laccata* and mycorrhization helper bacteria (MHB) in two bare root forest nurseries. Plant Soil 138:169-176, (1991).
  27. Dwivedi S.K., Dwivedi Padmanabh. Mycorrhizae in ecosystems: an ecofriendly approach for improved plant growth. In: Biotechnology of microbes and sustainable utilization (ed. R.C. Rajak), Scientific Publishers, Jodhpur, 2002, pp. 24-32.
  28. El-Haddad S.A., Abd El-Megid M.S., Shalaby O.Y. Controlling onion white rot by using Egyptian formulated endo-mycorrhiza (Multi-VAM). Ann. Agric. Sci., 49: 733-745, (2004).
  29. Franco-Correa M., Quintana A., Duque C., Suarez C., Rodriguez M.X., Barea J.M. Evaluation of actinomycete strains for key traits related with plant growth promotion and mycorrhiza helping activities. Applied Soil Ecology 45: 209-217, (2010).
  30. Frey B., Schüepp H. Acquisition of nitrogen by external hyphae of Arbuscular Mycorrhizal Fungi associated with *Zea mays* L. New phytol., 124: 221-230, (1993).
  31. Frey-Klett P., Pierrat J.C., Garbaye J. Location and survival of mycorrhiza helper *Pseudomonas fluorescens* during establishment of ectomycorrhizal symbiosis between *Laccaria bicolor* and *Douglas fir*. Appl. Environ. Microbiol., 63: 139-144, (1997).
  32. Gao X., Lu X., Wu M., Zhang H., Pan R., Tian J., Li S., Liao H. Co-Inoculation with Rhizobia and AMF Inhibited Soybean Red Crown Rot: From Field Study to Plant Defense-Related Gene Expression Analysis. Public Library of Science one, 7(3):e33977, (2012).
  33. George E., Haussler K., Vetterlein D., Gorgus E., Marschner H. Water and nutrient translocation by hyphae of *Glomus mosseae*. Can. J. Bot., 70: 2130-2137, (1992).
  34. Gyaneshwar P., Kumar G.N., Parekh L.J., Poole P.S. Role of soil microorganisms in improving P nutrition of plants. Plant and Soil, 245: 83-93, (2002).
  35. Harrier L.A., Watson C.A. The potential role of arbuscular mycorrhizal (AM) fungi in the bioprotection of plants against soil-borne pathogens in organic and/or other sustainable farming systems. Pest Manage. Sci., 60: 149-157, (2004).
  36. Ilbas A.I., Sahin S. *Glomus fasciculatum* inoculation improves soybean production. Acta Agriculture Scandinavica Section B- Soil and Plant Sci., 55(4): 287-292, (2005).
  37. Jaiti F., Kassami M., Meddich A., El Hadrami I. Effect of arbuscular mycorrhization on the accumulation of hydroxycinnamic acid derivatives in date palm seedlings challenged with *Fusarium oxysporum* f. sp. *albedinis*, J. Phytopathol 156: 641-646, (2008).
  38. Jeffries P., Barea J.M. Biochemical cycling and arbuscular mycorrhizas in the sustainability of plant-soil system. In: Gianinazzi S, Schüepp H, editors. Impact of Arbuscular Mycorrhizas on Sustainable Agriculture and Natural Ecosystems. Basel,

- Switzerland: Birkhäuser Verlag, 1994, pp. 101–15.
39. Kasiamdari R.S., Smith S.E., Smith F.A., Scott E.S. Influence of the mycorrhizal fungus, *Glomus coronatum*, and soil phosphorus on infection and disease caused by binucleate *Rhizoctonia* and *Rhizoctonia solani* on mung bean (*Vigna radiata*). Plant Soil 238:235–244, (2002).
  40. Khaosaad T., Garcia-Garrido J.M., Steinkellner, Vierheilig H. Take all disease is systematically reduced in roots of mycorrhizal barley plants. Soil Biol. Biochem., 39:727-734, (2007).
  41. Kobra N., Jalil K., Youbert G. *Arbuscular mycorrhizal* fungi and biological control of *Verticillium* wilted cotton plants. Archives of phytopathology and Plant Protection 44(10): 933-942, (2011).
  42. Krishna H., Das B., Attri B.L., Grover M., Ahmed N. Suppression of *Botryosphaeria* canker of apple by arbuscular mycorrhizal fungi. Crop Protection 29: 1049 – 1054, (2010).
  43. Matsubara Y., Ohba N., Fukui H. Effect of arbuscular mycorrhizal fungus infection on the incidence of Fusarium root rot in asparagus seedlings. J. Jap. Soc. Hortic. Sci., 70: 202-206, (2001).
  44. Neeraj, Singh K. Biochemical changes in *Phaseolus vulgaris* L. dual inoculated with arbuscular mycorrhizal fungi and Rhizobium, Ind. J. Bot. Res. 4:73-88, (2008).
  45. Neeraj, Singh K. Organic amendments to soil inoculated *arbuscular mycorrhizal* fungi and *Pseudomonas fluorescens* treatments reduce the development of root-rot disease and enhance the yield of *Phaseolus vulgaris* L. European Journal of Soil Biology 47:288-295, (2011).
  46. Ojha S., Chakraborty M., Chatterjee N.C. Influence of salicylic acid and *Glomus fasciculatum* on *Fusarial* wilt of tomato and Brinjal. Archives of phytopathology and Plant Protection 45(13):1599-1609, (2012).
  47. Paulitz T.C., Linderman R.G. Mycorrhizal interactions with soil organisms. In: Handbook of Applied Mycology. Vol. 1: Soil and Plants, Arora, D.K., Rai, B., Mukerji KG and Knudsen GR, Eds., Marcel Dekker Inc., New York, 1991, pp 77-129.
  48. Piccini D., Azcón R. Effect of phosphate-solubilizing bacteria and vesicular arbuscular mycorrhizal (VAM) on the utilization of bayoran rock phosphate by alfalfa plants using a Sand-vermiculite medium. Plant Soil 101:45–50, (1987).
  49. Plenchette C., Clermont-Dauphin C., Meynard J.M., Fortin J.A. Managing arbuscular mycorrhizal fungi in cropping systems. Canadian J. Plant Sci., 85: 31-40, (2005).
  50. Ravnskov S., Larsen J., Olsson P.A., Jakobsen I. Effects of various organic compounds growth and phosphorus uptake of an arbuscular mycorrhizal fungus. New Phytol 141: 517-524, (1999).
  51. Reddy M.S., Wang Q. Plant growth promoting rhizobacteria (PGPR) for sustainable agriculture. Proceedings of the 2<sup>nd</sup> Asian PGPR conference August 21-24, Beijing, P.R. China, (2011).
  52. Reid C.P.P. Mycorrhizas. In: The Rhizosphere, Lynch, J.M., Ed., John Wiley & Sons, Chichester, England, 1990, pp 218-315.
  53. Rillig M.C., Mummey D.L. Mycorrhizas and soil structure. New Phytologist 171: 41-53, (2006).
  54. Sabannavar S.J., Lakshman H.C. Effect of Rock Phosphate Solubilization Using Mycorrhizal Fungi and Phosphobacteria on Two High Yielding of *Sesamum indicum* L. World Journal of Agricultural Sciences 5 (4): 470-479, (2009).
  55. Schachtman D.P., Reid R.J., Ayling S.M. Phosphorus Uptake by Plants: From Soil to Cell. Plant Physiology 116(2): 447-53, (1998).
  56. Schellenbaum L., Berta G., Ravolanirina F., Tisserant, Gianinazzi S., Fitter A.H. Influence of endomycorrhizal infection on root morphology in a micropropagated woody plant species (*Vitis vinifera* L.). Ann Bot 68: 135 – 41, (1991).
  57. Secilia J., Bagyaraj D.J. Bacteria and actinomycetes associated with pot cultures of vesicular-arbuscular mycorrhizas. Can. J. Microbiol., 33: 1069-1073, (1987).
  58. Siddiqui Z.A., Pitchel J. Mycorrhizae : an overview ; In : Z.A. Siddiqui , M.S. Akhtar , K. Futai (Eds). Mycorrhizae: Sustainable agriculture and forestry. Springer. 2008, pp. 135.

59. Sieverding E. Vesicular-Arbuscular Mycorrhiza management in tropical Agrosystems. Deutsche Gesellschaft für Technische Zusammenarbeit, Eschborn, (1991).
60. Singh A., Sharma J., Rexer K.H., Varma A. Plant productivity determinants beyond minerals, water and light: *Piriformospora indica* – A revolutionary plant growth promoting fungus. *Curr Sci* 79:1548–1554, (2000).
61. Singh J.S., Pandey V.C., Singh D.P. Efficient soil microorganisms: A new dimension for sustainable agriculture and environmental development. *Agriculture, Ecosystems and Environment* 140: 339-353, (2011).
62. Singh N.K. Organic amendments to soil inoculated arbuscular mycorrhizal fungi and *Pseudomonas fluorescens* treatments reduce the development of root-rot disease and enhance the yield of *Phaseolus vulgaris* L. *European Journal of Soil Biology* 47:288-295, (2011).
63. Smith S.E., Facelli E., Pope S., Smith F.A. Plant performance in stressful environments: interpreting new and established knowledge of the roles of arbuscular mycorrhizas. *Plant and Soil* 326: 3-20, (2010).
64. Smith S.E., Read D.J. *Mycorrhizal symbiosis*, 2nd ed. Academic Press, London, UK, (1997).
65. Smith S.E., Read D.J. *Mycorrhizal Symbiosis*. Third edition. Amsterdam: Academic Press, (2008).
66. Smith S.E., Robson A.D., Abbott L.K. The involvement of mycorrhizas in assessment of genetically dependent efficiency of nutrient uptake and use *Plant Soil*, 146: 169-179, (1992).
67. Srivastava R., Khalid A., Singh U.S. Sharma A.K. Evaluation of arbuscular mycorrhizal fungus, *fluorescent Pseudomonas* and *Trichoderma harzianum* formulation against *Fusarium oxysporum* f. sp. *lycopersici* for the management of tomato wilt. *Biological Control* 53: 24–31 (2010).
68. Staley T.E., Lawrence E.C., Nance E.L. Influence of a plant growth-promoting pseudomonad and vesicular-arbuscular mycorrhizal fungus on alfalfa and birdsfoot trefoil growth and nodulation. *Biol. Fertil. Soils*, 14: 175-180, (1992).
69. St-Arnaud M., Vujanovic V. Effect of the arbuscular mycorrhizal symbiosis on plant diseases and pests. In: Hamel C, Plenchette C (eds). *Mycorrhizae in crop production: applying knowledge*. Haworth, Binghampton, N.Y., USA, 2007, pp 67-122.
70. St-Arnaud M., Elsen A. Interaction of arbuscular mycorrhizal fungi with soil-borne pathogens and non-pathogenic rhizosphere micro-organisms. In: Declerck S, Strullu DG, Fortin JA (eds) *In vitro culture of mycorrhizas*. Springer, Heidelberg, 2005, pp 217–23.
71. Subramanian K.S., Charest C. Acquisition of N by external hyphae of an arbuscular mycorrhizal fungus and its impact on physiological responses in maize under drought-stressed and well-watered conditions. *Mycorrhiza*, 9: 69-75, (1999).
72. Sundram S., Meon S., Seman I. A., Othman R. Application of arbuscular mycorrhizal fungi with *Pseudomonas aeruginosa* UPMP3 reduces the development of *Ganoderma* basal stem rot disease in oil palm seedlings. *Mycorrhiza*, DOI 10.1007/s00572-014-0020-5 (2014).
73. Tanaka Y., Yano K. Nitrogen delivery to maize via mycorrhizal hyphae depends on the form of N supplied. *Plant, Cell and Environment* 28:1247–1254 (2005).
74. Tarafdar J.C., Marschner H. Phosphatase activity in the rhizosphere and hyposphere of VA mycorrhizal wheat supplied with inorganic and organic phosphorus. *Soil Biol Biochem* 26: 387-395, (1994a).
75. Thygesen K., Larsen J., Bodker L. Arbuscular mycorrhizal fungi reduce development of Pea root-rot caused by *Aphanomyces euteiches* using oospores as pathogen inoculum. *Eur. J. Plant Pathol.*, 110: 411-419, (2004).
76. Tiwari P., Prakash A., Adholeya A. Commercialization of arbuscular mycorrhizal biofertilizer. *Handbook of Fungal Biotechnology*, 2003, pp. 195-203.
77. Toro M., Azcón R., Barea J.M. Improvement of arbuscular mycorrhiza development by inoculation of soil with phosphate-solubilizing rhizobacteria to improve rock phosphate bioavailability (32P) and nutrient

- cycling. Appl Environ Microbiol 63:4408–12, (1997).
78. van der Heijden M.G.A., Rinaudo V., Verbruggen E., Scherrer C., Barberi P., Giovannetti M. The significance of mycorrhizal fungi for crop productivity and ecosystem sustainability in organic farming systems. 16th IFOAM Organic World Congress, Modena, Italy, June 16-20, (2008).
79. van der Heijden M.G.A., Streitwolf-Engel R., Riedl R., Siegrist S., Neudecker A., Ineichen K., Boller T., Wiemken A., Sanders I.R. The mycorrhizal contribution to plant productivity, plant nutrition and soil structure in experimental grassland. New Phytol. 172: 739-752, (2006).
80. Wafaa M.H., Abdel-latif, Faten M. Interaction between Vasicular Arbuscular Mycorrhizae and Antagonistic Biocontrol Micro-Organisms on Controlling Root Rot Disease Incidence of Geranium Plants. Journal of biological sciences. 1(12): 1147-1153, (2001).
81. Wright D.P., Read D.J., Scholes J.D. Mycorrhizal Sink Strength Influences Whole Plant Carbon Balance of *Trifolium repens* L. Plant, Cell and Environment 21: 881-91, (1998).
82. Wu Q.S., Zou Y.N. Beneficial roles of arbuscular mycorrhizas in citrus seedling at temperature stress. Scientia Horticulturae 125: 289-293, (2010).
83. Xavier L.J.C., Boyetchko S.M. Arbuscular mycorrhizal fungi in plant disease control. In: Arora DK (ed) Fungal biotechnology in agricultural, food, and environmental applications. Dekker, New York, 2014, pp 183–194.
84. Yousefi A.A., Khavazi K., Moezi A.A., Rejali F., Nadian H.A. Phosphate Solubilizing Bacteria and Arbuscular Mycorrhizal Fungi Impacts on Inorganic Phosphorus Fractions and Wheat Growth. World Applied Sciences Journal 15 (9): 1310-1318, (2011).
85. Zachee A., Bekolo N., Bime Dooh N., Yalen M., Godswill N. Effect of mycorrhizal inoculum and urea fertilizer on diseases development and yield of groundnut crops (*Arachis hypogaea* L.), Afr. J. Biotech. 7:2823-2827, (2008).
86. Zapata F., Axmann H. 32 P isotopic techniques for evaluating the agronomic effectiveness of rock phosphate materials. Fert. Res., 41: 189-195, (1995).
87. Ziedan E.I., Sayed E.I., Mostafa M., Sahab A. Application of Mycorrhizae for controlling Root Diseases of sesame. Journal of Plant Protection Research. 51(4): 355-361, (2011).