



Internationally indexed journal

Indexed in Chemical Abstract Services (USA), Index copernicus, Ulrichs Directory of Periodicals, Google scholar, CABI ,DOAJ , PSOAR, EBSCO , Open J gate , Proquest , SCOPUS , EMBASE ,etc.



Rapid and Easy Publishing

The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quarterly. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences



Pharmaceutical Sciences

- Pharmaceutics
- Novel drug delivery system
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics



Biological Sciences

- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmacoinformatics
- Biopharmaceutics
- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

**Indexed in Elsevier Bibliographic Database
(Scopus and EMBASE)**

SCImago Journal Rank 0.288

Impact factor 5.121*

Chemical Abstracts
Service (www.cas.org)



A division of the American Chemical Society

CODEN IJPBJ2



Elsevier Bibliographic databases (Scopus & Embase)

SNIP value – 0.77

SJR - 0.288

IPP - 0.479

SNIP – Source normalised impact per paper

SJR – SCImago Journal rank

IPP – Impact per publication

Source – www.journalmetrics.com

(Powered by scopus (ELSEVIER))



LUND
UNIVERSITY



JACKSONVILLE STATE UNIVERSITY
Jacksonville State University
Houston Cole Library
USA (Alabama)



Oxford, United Kingdom

INDEX COPERNICUS
INTERNATIONAL

*And indexed/catalogued in
many more university*



*Instruction to Authors visit www.ijpbs.net

For any Queries, visit "contact" of www.ijpbs.net



MEDICINAL PROPERTIES AND GENETIC DIVERSITY OF A MAGICAL SHRUB *TINOSPORA CARDIFOLIA*: AN UPDATE

PRACHI GUPTA AND SAURABH KULSHRESTHA*

*Faculty of Applied Science and Biotechnology Shoolini University of Biotechnology
and Management Sciences, Bajhol, Solan, Himachal Pradesh.*

ABSTRACT

Before the introduction of chemical medicines, man relied on the healing properties of medicinal plants. Perhaps 90% of the world's population still relies completely on raw herbs and unrefined extracts as medicines. Medicinal plants were the "backbone" of traditional medicine, which means more than 3.3 billion people in the less developed countries utilize medicinal plants on a regular basis. *Tinospora cardifolia* is one of the important medicinal plants which is a large and deciduous climbing shrub found at higher altitudes of India cultivated in tropical areas of India and known by names such as Gurchar, Heart leaved moonseed, Heavenly elixir and Jetwatika. About 152 plants have been documented in India for traditional ayurvedic medicines and out of these 152 plants; *T. cardifolia* is one of the most promising plants in the herbal medicine system. The Plant *Tinospora* includes 76 scientific rank species but only 13 of these species are accepted for medicinal uses. Different parts of *T. cardifolia* such as leaves, roots and stem are emetic and used in ayurvedic, "Rasayanas" to improve the immune system and the body resistance against infections such as rheumatism, urinary diseases, dyspepsia, general debility, syphilis, skin diseases, bronchitis, spermatorrhea, impotence, general debility, dyspepsia, fevers, urinary diseases, visceral obstruction and leprosy. This review gives a bird view of the main biological activities, pharmacological actions, medicinal applications of *T. cardifolia* extracts and genetic diversity of *T. cardifolia*.

KEY WORDS: *Tinospora cardifolia*, antimicrobial, genetic diversity, medicinal, Rasayanas.



SAURABH KULSHRESTHA

Faculty of Applied Science and Biotechnology Shoolini University of Biotechnology
and Management Sciences, Bajhol, Solan, Himachal Pradesh.

INTRODUCTION

The tradition to use plant as a medicine is thousands of years old. Recently there has been a shift in universal trend from synthetic to herbal medicine, which we can say 'Return to Nature'. Medicinal plants have been known for millennia and are highly esteemed all over the world as a rich source of therapeutic agents for the prevention of diseases and ailments¹. Many drugs commonly used today are having herbal origin. Some are made from plant extracts while others are synthesized to mimic a natural plant compound. In recent years, pharmaceutical companies were spending a lot of time and money in developing natural products that are extracted from plants. Different parts of the world are working to produce more cost effective remedies that are affordable to the public². In India, medicines based on herbal origin have been the basis of treatment and cure for various diseases and physiological abnormalities under practice such as Ayurveda, Siddha and Unani. Research in herbal medicine has recently been revolutionized with the identification of several botanical plants with established physiological effect and efficacy for clinical condition either alone or combination with pharmaceuticals. Ayurveda was a comprehensive natural health care system that organized in India more than 5000 years ago. Plant antioxidants are composed of a broad variety of different substances like ascorbic acid, tocopherols, polyphenolic compounds and terpenoids. They perform several important functions in plants and humans³. *T. cardifolia* commonly named as "Guduchi" is known for its immense application in the treatment of various diseases in the traditional ayurvedic literature⁴. In India, various extracts of *T. cardifolia* are used as a remedy for many diseases and are included in various polyherbal preparations used for the treatment of diabetes, hepatitis, etc. During the last few decades considerable progress has been achieved regarding its biological activity and medicinal applications. Hence, it can be chosen as a source for the development of industrial products for treatment of various diseases. This review gives a bird view of the

main biological activities, pharmacological actions, medicinal applications of *T. cardifolia* extracts and also biological activities of few *T. cardifolia* compounds isolated⁵.

Medicinal properties of *Tinospora cardifolia*

Ayurvedic literature quoted *T. cardifolia* as a constituent of several compound preparations which has been used in general debility, dyspepsia, fever and urinary diseases⁶. *T. cardifolia* family is rich in various compounds i.e simple sugars, rhamnase, glucosinolates and isothiocyanates⁷. It has multiple actions like stem is a bitter stomachic i.e it strengthens and excites stomach activities and stimulates bile secretion also⁸. Fresh juice of guduchi, when mixed with rock candy, speeds up the recovery in hepatitis patients and leaves of *T. cardifolia* along with honey acts as a tonic, enriches the blood and enlarges the spleen⁹. The juice of the plant is useful in diabetes, vaginal and urethral discharges and root and stem are prescribed in combination with other drugs as an antidote to snake bite and scorpion sting¹⁰. The herb accords longevity, enhances memory, improves health, bestows youth, betters complexion, voice, energy and luster of the skin and hence enjoyed the reputation among ancient Hindu writers of being an aphrodisiac¹¹. Guduchi juice, when taken with cow's milk or lodhra, is effective in combating leucorrhoea and the juice when consumed after mixing with cumin seeds reduces the burning sensation caused due to pitta¹².

BOTANICAL AND PHARMACOGNOSTICAL PROPERTIES

Botanical description

The plant is large, deciduous extensively climbing shrub with several elongated twining branches. Leaves are simple, exstipulate, long petioles upto 15 cm. Both at base and apex they are roundish and pulvinate. Flowers grow separately on small plants and appear only when the plant is leafless, greenish yellow on axillary and terminal racemes. Sepals are 6 in number and smaller ones are on outer side while inner ones are on the inner side. Petals are membranous, smaller than sepals and are obvate. Fruits appear in aggregate of 1-3, ovoid

smooth drupelets on thick stalk with sub terminal style scars, scarlet or orange coloured¹³.

Ethanobotanical and Pharmacognostical description

Various parts of *T. cardifolia* are being prescribed in Ayurveda and other systems of medicine as a monoherbal or polyherbal preparation¹⁴. Since each part of *T. cardifolia* has some medicinal property, it is very much commercially exploitable. The diagnostic pharmacognostical characteristics of medicinal parts are as follows.

Leaf

Single layer of epidermis and a wide zone of cortex composed of 3-4 layers of epidermis are found in cross section of leaf. Trichomes are not found. Petiole is circular in the transverse section. The vascular bundles consist of radial rows of cambium cells on the outer side and a few rows of cambium cells on the inner side followed by phloem¹⁵. The epidermal cells surround the base. Starch is present throughout the tissue¹⁶. Veins are multicostate and prominent on dorsal side^{17,18}. Leaves of *T. cardifolia* acts as Vasorelaxant and relaxes norepinephrine induced contractions¹⁹ and inhibits Ca⁺⁺influx^{20,21}. Leaves acts as anti-inflammatory, anti-microbial, antihypertensive and anti-viral agents²². Leaves of *T. cardifolia* induce apoptosis in leukemia by activating caspase-3 and bax²³.

Stem

Stems are greyish green with smooth surfaces and swelling at nodes, older ones are light brown and surface marked with warty protuberances due to circular lenticels. Outer bark is thin and papery brown to greyish in colour²⁴. Lenticels are circular and prominent. Transverse surface have wheel like structure which is a characteristic feature of family *Menispermaceae*²⁵. Stems of *T. cardifolia* are useful in the Treatment of neurological disorders like Parkinson²⁶, Dementia²⁷, motor and cognitive deficit, neuron loss in spine and hypothalamus²⁸. Stems by Immunomodulation inhibit NF-kB and acts as nitric oxide scavenger to show anticancer activities³⁰.

Root

The young aerial roots are long filiform, threadlike, squarish, which arise from the mature branches or cut bits of stems grow downward and by continuously lengthening sometimes reach the ground and these roots are characterized by tenta to penta primary structure³¹. Oval or elliptical starch grains are present which are mostly simple but sometimes as compound grains of 2-5 components with the presence of central hilum and concentric striations³². Roots are useful in anti-viral infections³³, anticancer³⁴ and anti-diabetes^{35,36}. Roots are helpful in the treatment of inflammation³⁷, Neurological and psychiatric conditions³⁸. Roots acts as Protease inhibitors for HIV and drug resistant HIV^{39,40}.

Flowers

Flowers are unisexual, racemes, greenish yellow in colour, appears when plant is leaf less. Male flowers are clustered together but female flowers exist in solitary inflorescence⁴¹. Flowering occurs during March to June⁴²; Flowers were useful in the treatment of IgA neuropathy and glucocorticoid induced osteoporosis which is in the early stage of inflammatory arthritis⁴³. Flowers also induce cell cycle arrest in G2/M phase and apoptosis through c-Myc suppression⁴⁴. Flowers are useful in the Inhibition of TNF- α , IL-1 β , IL-6 and COX-2⁴⁵.

Fruits

Fruits of *T. cardifolia* grow in aggregates of 1-3 ovoid smooth drupelets on thick stalk with sub terminal style scars. Colour of the fruits is Scarlet or orange red color. Fruits are found during winter season⁴⁶. Fruit was a source of bioactive food colorants⁴⁷. The mature fruits are closely associated with cell metabolism⁴⁸ and growth^{49,50}.

CHEMICAL CONSTITUENTS

Phytochemicals are the chemicals produced by plants. These refer to those chemicals which may have an impact on health, or on flavour, texture, smell or color of the plants, but are not required by humans as essential nutrients. In plants, phytochemicals are the

secondary metabolites such as alkaloids, terpenoids, steroids, tannins, phenolic compounds, flavonoids, resins fatty acids and gums which are capable of producing definite physiological action on body⁵¹.

Alkaloids

Alkaloids are a group of naturally occurring chemical compounds that contain mostly basic nitrogen atoms. *T. cardifolia* is rich in various alkaloids such as Berberine, Palmatine, Tembetarine, Magnoflorine, Choline, Tinosporin^{52,53,54,55,56,57}, Isocolumbin, Palmetine, Tetrahydropalmatine, Magnoflorine⁵⁸, Aporphine, Jatrorrhizine and Tetrahydropalmatine^{52,53,54,55,56}. Alkaloids of *T. cardifolia* possess antitumor activity and are also used as medications or as recreational drug in various pharmacological conditions.

Diterpenoids

Diterpenoids present in *T. cardifolia* are Furanolactone Lactones and Clerodane derivatives⁵⁹, 5R,10R-4R-8R-dihydroxy-2S-3R:15,16-diepoxy-cleroda-13 (16), 14-dieno-17,12S:18,1S-dilactone^{52,53,54,55,56,57}. These terpenoids have medicinal properties such as anti-carcinogenic, antimalarial (e.g. artemisinin), anti-ulcer, hepaticidal, antimicrobial and diuretic (e.g. glycyrrhizin) activity. Terpenoids and their derivatives also belong to other drugs such as validol, menovasin, turpentine, bromkamfora etc.⁶⁰.

Steroids

Steroids of *T. cardifolia* are β -sitosterol, δ -sitosterol, alpha hydroxyacetone, 20 β -hydroxyecdysone, Ecdysterone, Makisterone A and Giloinsterol^{44,45,61}. Steroids stimulate the growth of osteoblasts, increases the differentiation of cells into osteoblastic lineages and also increases the mineralization of bone like matrix⁴.

Glycosides

Glycosides present in *T. cardifolia* are 18-norclerodane glucoside, Furanoid diterpene glucoside^{26,29,62}, Tinocordiside, Tinocordifolioside, Cordioside, Cordifolioside A, Cordifolioside B, Syringin, Syringin-

apiosylglycoside, Palmatosides C, Palmatosides F, Cordifolioside A, Cordifolioside B, Cordifolioside C, Cordifolioside D and Cordifolioside E and Pregnane glycoside^{21,26,29,63}. These all glycosides have many properties such as antioxidant or radical scavenging activity, inhibition of lipid peroxidation and lipoxygenases *in vitro*⁶⁴, antimicrobial and antiviral⁶⁵, antimutagenic⁶⁶, and antidiabetic properties⁶⁷. The antioxidant activity of glycosides provides synergistic benefits for the treatment of diabetes because of their insulin enhancing potential⁶⁸.

Sesquiterpenoid

Tinocordifolin is an important sesquiterpenoid present in *T. cardifolia* which plays a major role in plant development. They act as constitutive protective agents against invading organisms, signal molecules and allelopathic compounds and natural animal toxicants. A regular intake of Sesquiterpenoid decreases the incidence of certain forms of cancer so they were known as chemo-preventive agents⁶⁹.

Miscellaneous compounds

Other compounds in *T. cardifolia* which are immunologically active are Giloin, Giloinin, Tinosporic acid, Tinospororan acetate, Tinosporal acetate, Tinosporidine, Aliphatic Octacosanol, Heptacosanol compound, Nonacosan, Heptacosanol, Cordifolane, Octacosanol, Tinosporic acid, Tinosponone, Tinosporal, 2-Phytoecdysone and Arabinogalactan^{39,40}. Tinosporic acid and Tinospororan acetate showed protection against aflatoxin induced nephrotoxin³⁷. Cordifolane, Arabinogalactan and Giloinin showed radio protective effects¹²².

GENETIC DIVERSITY

The genetic diversity of a species is an important element in establishing conservation programs because the ability of a species to respond adaptively to environmental changes depends on the level of genetic variability it contains^{70,71,72}. Therefore, assessing the level and distribution of genetic diversity are crucial for management and the development of effective conservation strategies. Faced with

the problem of preserving rare and endangered species, great concerns also should be concentrated on the endemic species with restricted geographic distribution⁷³. Medicinal plants are increasingly endangered. The survival index of a species is greatly determined by the percentage of polymorphism and the gene flow between the populations. Molecular markers like RAPD, AFLP, RFLP, SSR SCAR etc are being continuously used for assessing the genetic diversity of plants and these are considered vital for formulating conservation strategies for endangered medicinal plants⁷⁴. In order to develop an efficient identification method, molecular techniques have been used since these are reliable, unaffected by environmental conditions and can aid varietal identification⁷⁵. Among the different types of molecular markers available, random amplified polymorphic DNA (RAPD) is useful for the assessment of genetic diversity among rare species of medicinal plants⁷⁶. RAPD markers have many advantages like the requirement of very small quantity of template DNA and no involvement of radioactive or hybridization techniques. RAPD markers have been extensively used in constructing linkage maps, assessment of genetic variation in population, gene tagging and identification of cultivars and species⁷⁷. Molecular markers are not tissue-specific and can be detected at any stage of plant development. Therefore, DNA-based authentication of medicinal plants can be a useful tool for quality control and safety monitoring of herbal pharmaceuticals and nutraceuticals. Among the polymerase chain reaction (PCR)-based DNA techniques, random amplified polymorphic DNA (RAPD) is easy to perform and requires no information about the DNA sequence to be amplified. It has been used, therefore, to study the genetic diversity in various plant species^{78,79}, authenticate the herbal medicinal materials⁸⁰, and detect adulterants⁸¹. Rasayana churna includes three herbal drugs in powdered Dried stem of *T. cordifolia*, dried fruit of *E. officinalis* and dried fruit of *T. terrestris* were collected from the Pune regions of India. After 120 primers were screened, primer OPA-6 was screened and selected to identify *T. cordifolia*, *E. officinalis*

and *T. terrestris* in Ayurvedic prescription⁸². Identified germplasm is an important component for the efficient and effective management of plant genetic resources. Studies were undertaken for identification and to assess genetic variation within 15 clones of *T. cordifolia* collected from Bhubaneswar through random amplified polymorphic DNA (RAPD) markers. Analysis was made using forty decamer primers. Out of them, 15 primers were selected and used for identification and genetic relationships within 15 clones. The genetic distance was very close within the clones. This polymorphism study with the help of RAPD markers will be helpful to know the genetic background of the medicinal plants with high commercial value and also provides a major input into conservation biology⁸³. The phylogenetic relationships among the three species of *Tinospora* found in India are poorly understood. Morphology alone does not fully help to resolve the phylogeny and therefore a fast approach used molecular analysis was explored. Two molecular approaches viz Random Amplified Polymorphic DNA (RAPD) assay and restriction digestion of ITS1-5.8S-ITS2 rDNA Restriction fragment length polymorphism (RFLP) were used to evaluate the genetic similarities between 40 different accessions of *T. cordifolia*, *T. malabarica* and *T. crispa* collected from Jammu and Kashmir, Assam and South Kerala respectively. Out of 38 random primers used only six generated the polymorphism, while three out of 11 restriction enzymes used gave polymorphic restriction patterns. Three independent clones of each species *T. cordifolia*, *T. malabarica* and *T. crispa* were sequenced. Phylogenetic relationship inferred from ITS sequences was in agreement with RAPD data⁸⁴. Based on morphology, the species status and taxonomic affinities of three species of *Tinospora* (*T. cordifolia*, *T. sinensis*, and *T. crispa*) with ranges in India have been questioned. To evaluate species delimitation and population structure among 40 accessions of the three species, a relatively new marker, cytochrome P450, was used. 40 accessions were collected from multiple locations within India i.e Jammu & Kashmir, Himachal Pradesh, Uttar Pradesh,

Assam, Kerala, Haryana and Delhi. Five out of nine primers generated polymorphisms with 39 out of 47 bands were found to be polymorphic. The resulting phylogenies highly support the Indian *Tinospora* species as part of a clade (expanded *Tinosporeae*), consists of diverse Menispermaceae from around the world. The three Indian species are monophyletic and are most closely related to *Tinospora* species from Australia (*T. esiangkara* and *T. smilacina*)⁸⁵. Genetic variation among the 20 selected samples of *T. cardifolia* from Western Himalayas using RAPD marker was done using 120 decamer oligonucleotide primers. Out of 120 primers used only four RAPD primers i.e OPA-16, OPC-5, OPC-7 and OPC-13 revealed low genetic diversity among *T. cardifolia*⁸⁶. From Gujrat, Genetic diversity was accessed by isoenzymatic method using 25 accessions of the medicinal climber, *T. cardifolia*. Analysis of ten isozymes revealed the presence of 16 gene loci and 33 alleles in 25 accessions. The percentage of polymorphic loci (P) was 45.0% and mean observed number of alleles per locus (A) was 1.57. The average observed heterozygosity (Ho) and expected heterozygosity (He) were 0.443 and 0.270 respectively showed high levels of genetic variations among different accessions. The unweighted paired group method (UPGMA) dendrogram clearly depicted that the spectra of genetic diversity among various accessions of *T. cardifolia*⁸⁷.

TRADITIONAL USES IN INDIGENOUS SYSTEM OF MEDICINE

Traditional medicine has remained as the most affordable and easily accessible source of treatment in the primary healthcare system of resource poor communities in the world. Medicinal plants, since times immemorial, have been used in virtually all cultures as a source of medicine. The widespread use of herbal remedies and healthcare products were described in Vedas and Bible. The local people have a long history of traditional plant usage for medicinal purposes⁸⁸. Furthermore, an increasing reliance on the use of medicinal plants in the industrialized societies has been traced to the extraction and development of

several drugs and chemotherapeutics from these plants as well as from traditionally used rural herbal remedies. The economic significance of medicinal plants stems from the fact that the number of patients suffering from chronic ailments is on the rise and drugs from medicinal plants are proving to be more effective in treating chronic ailments⁸⁹. The three main species of *Tinospora* are: *T. cardifolia*, *Tinospora malabarica* and *Tinospora crispa*. *T. cardifolia* mainly found at tropical region of India and is widely used as a medicine in folk and ayurvedic systems. *T. cardifolia* is of great interest to researchers across the globe because of its reported medicinal properties like anti-diabetic, anti-periodic, anti-spasmodic, anti-inflammatory, anti-arthritic, anti-oxidant, anti-allergic, anti-stress, anti-leprotic, anti-malarial, hepatoprotective, immunomodulatory and anti-neoplastic activities. *Tinospora malabarica* found in Konkan, Karnataka, Tamilnadu, Bengal, and Orissa is mixed as adulterant or substitute with *T. cardifolia* and is also used as an immunomodulator⁹⁰.

ANTIMICROBIAL ACTIVITY OF TINOSPORA CARDIFOLIA

The antibacterial activity of the aqueous, ethanol and chloroform extracts were tested with the disc diffusion method against a number of Gram-negative and Gram-positive bacteria, the results suggested that the ethanolic extract has significant antibacterial activity, which justified the uses of *T. cardifolia* in traditional medicine in various parts of the countries to treat various infectious diseases⁹¹. Secondary infections are the most common cause of mortality in the immunocompromised host. In Ayurveda, *T. cardifolia*, *Asparagus racemosus* and *Withania somnifera* are prescribed as agents to strengthen host defence mechanisms so that immunocompromised host can fight with the secondary infections. Administration of *T. cardifolia*, *Asparagus racemosus* and *Withania somnifera* helped in faster bacteremia clearance in the rats and mice suffering from *Escherichia coli* peritonitis. Mortality due to *Staphylococcus aureus* sepsis in neutropenic

and hemisplenectomized mice was reduced in animals treated with *T. cardifolia*, *Asparagus racemosus* and *Withania somnifera*. *Candida albicans* sepsis induced mortality was also reduced by these plants. In view of their oral efficacy and lack of adverse effects, *T. cardifolia*, *Asparagus racemosus* and *Withania somnifera* show promise as useful immunomodulators⁹².

USES OF T. CARDIFOLIA IN FOLK AND TRIBAL MEDICINE OF INDIA

T. cardifolia is widely used in veterinary folk medicine system of India for its general tonic, antiperiodic, anti-spasmodic, anti-inflammatory, antiarthritic, anti-allergic and anti-diabetic properties⁹³. The tribal Baiga living in the interior areas of Naugarh and Chakia Block of Varanasi district, Uttar Pradesh make the paste of the stems of Guduchi and the roots of Bhatkatiaya. The tribals of Bombay and its neighbouring areas and the fisherman along the sea coast use *T. cardifolia* in the treatment of fever, jaundice, chronic diarrhea and dysentery. Similarly, the tribals of North Gujrat used the powder of root and stem bark of the plant along with milk in day to day life for the treatment of dysentery and diarrhea. This paste was used in the treatment of fever for three days. Decoction of stem is administered orally by the people of Jammu and Kashmir for the treatment of fever⁹⁴. In dahanu forest division of Maharashtra, tribal races, viz. Agaris, Bhils, Dhodias, Khakarais, Dublas and Rimoshis uses the stem decoction with cold and hot water in morning in an empty stomach as a tonic in general debility. The muslim tribals of Rajouri, Jammu comprising Gujjar and Backwals used the plant in bone fracture⁹⁴. The Inhabitants of Bhuvneshwar uses the warm juice of root of *T. cardifolia* orally for the treatment of high fever. In Punjab juice of leaves along with honey is used to lower the body temperature and the ear pain. In Haryana powder of *Terminalia chebula*, *T. cardifolia* and *Trachyspermum ammi* administered orally, along with salt in the morning for the treatment of cough. The local women in the tribes of Rajasthan use the paste of 5 seeds of *Piper nigrum* along with few leaves of *T. cardifolia* in the treatment of

leucorrhoea. Inhabitants of Hisar take the juice of stem orally with honey for the treatment of asthma⁹⁵.

CHARACTERIZATION OF COMPOUNDS OF TINOSPORA CARDIFOLIA USING VARIOUS TECHNIQUES

Natural products, such as plants extract, either as pure compounds or as standardized extracts, provide unlimited opportunities for a new drug discoveries because of the unmatched availability of chemical diversity⁹⁶.

Due to the development of adverse effects and microbial resistance to the chemically synthesized drugs, men turned to ethnopharmacognosy. Researchers found literally thousands of phytochemicals from plants as safe and broadly effective alternatives with less adverse effect. Many beneficial biological activities such as anticancer, antimicrobial, antioxidant, antidiarrhoeal, analgesic and wound healing activity were reported in the medicinal plants⁹⁷. Botanicals and herbal preparations for medicinal usage contain various types of bioactive compounds and these compounds must undergo extraction, isolation and characterization so that they can be used in drug development.

Thin layer chromatography (TLC) of *Tinospora cardifolia*

In the methanolic stem extract of *T. cardifolia* six molecules were reported having different Rf values but the major compound found was Tinosporoside with the Rf value of 0.35¹³. Chloroformic extract of *T. cardifolia* showed Rf values of 0.24 and 0.35 in thin layer chromatography (TLC) identity test which was characterized as tinosporoside. The solvent system consists of chloroform and methanol. The spraying reagent was anisaldehyde-sulphuric acid⁹⁸. Quality of plants grown in tissue culture was ensured using the juice of the plants collected from various supports from the same garden and were analyzed by thin layer chromatography (TLC). A total number of 11 spots were observed. Out of 11 spots three Rf value i.e 0.192 (Tinosporoside), 0.551 (berberine) and 0.717 were found to be universally present in the plants and these are

used as chemical markers of the plant. Tinosporaside contents were found to be significantly high in *T. cardifolia*⁹⁹.

High-performance liquid chromatographic (HPLC) of *Tinospora cardifolia*

HPLC analysis of *T. cordifolia* showed six different phenolic acids (Benzoic acid, Cinnamic acid, Caffeic acid, Ferulic acid, Gallic acid and Tannic acid) present in varying amount in the ethanolic stem extract¹⁰⁰. A high-performance liquid chromatographic (HPLC) method for the estimation of berberine in the stem of *T. cordifolia* and *Tinospora sinensis* was performed. HPLC analysis gave peaks at 24.54 min and 28.48 min, in the methanolic extract of *T. sinensis*. *T. cardifolia* and *T. sinensis* showed differences in chemical constituents. The physicochemical studies of *T. cardifolia* stem crude drug were evaluated. The phytochemical screening was carried out and the berberine alkaloid was quantified in different fractions of the extract by HPLC. The result showed that the methanolic extracts have a higher concentration of berberine when compared to other solvent fractions¹⁰¹. The high performance liquid chromatography (HPLC) method was validated for accuracy, precision, reproducibility and robustness for the estimation of cordifolioside A in 60% methanolic extract of *T. cardifolia*. Statistical analysis proved that the method is precise, reproducible, selective, and accurate for the estimation of cordifolioside A. HPLC method was validated for accuracy, precision, reproducibility, and robustness as per International Conference on Harmonisation (ICH) guidelines^{102,103}.

IMMUNO-MODULATORY ACTIVITY OF TINOSPORA CARDIFOLIA

T. cardifolia is well researched for its immunomodulatory activities since many years and few compounds have been isolated and subjected for their possible mode of immuno-modulatory activity. According to a recent report, arabinogalactan polysaccharide (G1-4A) from the stem, appear to induce tolerance against endotoxic shock by modulation of cytokines and nitric oxide (NO). In addition, it also modulated the release of nitric oxide by marine

macrophages¹⁰⁴. The immunostimulating signaling mechanism of the novel (1,4)-alpha-D-glucan reported to be through the activation of macrophages that occurred through TLR6 signaling, NF-kappa B translocation and cytokine production¹⁰⁵. Polysaccharide fraction from *T. cardifolia* is found to be very effective in reducing the metastasis potential of B16F-10 melanoma cells¹⁰⁶. Arabinogalactan polysaccharide has good protective effect against iron mediated lipid peroxidation of rat brain homogenate as revealed by the thiobarbituric acid reactive substances and lipid hydroperoxide assays¹⁰⁷. The effect of aqueous extract of *T. cordifolia*, an immunomodulator with antimalarial activity along with chloroquine was studied in the treatment of three cases of hyper-reactive malarious splenomegaly in District Hospital, Daltonganj town, Jharkhand, India. Addition of extract of *T. cordifolia* for the first six weeks to chloroquine showed regression of spleen by 37–50% after six weeks and 45–69% after six months from the start of treatment¹⁰⁸.

ANTIHYPERGLYCAEMIC ACTION

T. cardifolia has been extensively studied for its hypoglycemic activity in support of its usefulness in the treatment of diabetes mellitus. An Ayurvedic polyherbal formulation 'Ilogen-Excel', which contains *T. cardifolia* as one of the constituent, administered at the dose of 50 mg/kg and 100 mg/kg for 60 days has shown a significant decrease in the blood glucose levels and increase in the plasma insulin, hepatic glycogen and total hemoglobin¹⁰⁰. One more herbomineral formulation "Hyponidd" is reported for its possible hypoglycemic as well as antioxidant activity and the results are comparable with earlier reports on this plant¹⁰⁹. Diabetics are prone to the development of cataract; alcohol extract of *T. cardifolia* has preventive effect on the development of cataract and produces a significant reduction of plasma glucose levels in alloxan induced diabetic rats¹¹⁰. The effects of *T. cardifolia* extracts on blood glucose concentrations as well as on kidney functions in streptozotocin induced diabetic rats have been investigated. The extract when administered orally for 40

days was found to decreased plasma glucose concentration and prevented polyuria, rise in urinary albumin levels and renal hypertrophy as well¹¹¹. In India, the decoction of kernels of *Eugenia jambolana* and extracts of *T. cardifolia* are used as a household remedy for diabetes. The anti-hyperglycemic effect of aqueous and alcoholic extracts as well as lyophilized powder of these two plants was evaluated in diabetic animals¹¹². The stem of *T. cardifolia* is widely used in the therapy of diabetes by regulating the blood glucose in traditional folk medicine of India and it has been reported to mediate its anti-diabetic potential through mitigating oxidative stress (OS), promoting insulin secretion and also by inhibiting gluconeogenesis and glycogenolysis, thereby regulating blood glucose¹¹³. The isoquinoline alkaloid rich fraction from stem, including, palmatine, jatrorrhizine, and magnoflorine have been reported for insulin-mimicking and insulin-releasing effect both *in vitro* and *in vivo*¹¹⁴. The protective effects of *T. cardifolia* were reported in the presence of higher levels of anti-oxidant molecules and enzymes³⁷. *T. cardifolia* has been shown to significantly counterbalance the diabetes-associated OS in the maternal liver by lowering the levels of malondialdehyde and ROS and the increased levels of GSH and total thiols¹¹⁵.

THERAPEUTIC APPLICATIONS

The ethanolic extract of the root of *T. cardifolia* was observed to induce a marked protective action against restrain stress induced ulcerization. The antiamebic activity of crude drug formulation containing *T. cardifolia* against *Entamoeba histolytica* showed varying degrees of inhibition of the enzymes i.e DNase, RNase, Aldolase, alkaline phosphatase, acid phosphatase and protease activities of crude extracts of axenically cultured amoeba. The water and ethanolic extracts of stem of the plant inhibited immunosuppression produced by cyclophosphamide. The ethanolic extract of stem of the plants inhibits cyclophosphamide induced anemia. Water extract was more potent than any other extract of the plant¹¹⁶.

ANTIOXIDANT/HEPATOPROTECTIVE ACTIVITY

The antioxidant activity of an arabinogalactan polysaccharide i.e Tamarind seed polysaccharide (TSP) isolated from *T. cordifolia*, an Indian medicinal plant, was studied. The polysaccharide showed good protection against iron-mediated lipid peroxidation of rat brain homogenate as revealed by the thiobarbituric acid reactive substances (TBARS) and lipid hydroperoxide (LOOH) assays. TSP also provided significant protection to protein against γ -ray induced damage. The protective action can possibly be explained by its very high reactivity towards 2,2-diphenyl-1-picrylhydrazyl (DPPH), superoxide radicals and the most damaging of radicals i.e the hydroxyl radical¹¹⁷. The 50% alcoholic extracts of *Swertia chirayita*, *Cedrus deodar*, *Boerhavia diffusa*, *Berberis aristata*, *Withania somnifera*, *Pongamia glabra*, *Petrocarpus santalinus*, *T. cardifolia* and *Acrois calamus* were screened for antioxidant and possible anti-inflammatory potential. Among the tested plants, only *Pongamia glabra*, *Petrocarpus santalinus*, *T. cardifolia*, and *Acrois calamus* were found to possess hydroxyl radical scavenging activity¹¹⁸.

CYTOTOXIC EFFECT

The anti tumor activity of *T. cardifolia* may be due to decreased lipid peroxidation, glutathione-S-transferase (GST) activity or due to the release of lactate dehydrogenase. Dichloromethane extracts showed concentration dependent decline in the clonogenicity, glutathione-S-transferase activity as well as increased in lipid peroxidation with a peak at 4 h and a lactate dehydrogenase release with a peak at 2 h¹¹⁹. Exposure of HeLa cells to 0, 5, 10, 25, 50 and 100 $\mu\text{g/ml}$ of guduchi extracts (methanol, aqueous and methylene chloride) resulted in a dose-dependent but significant increase in cell killing, when compared to non-drug-treated controls. The effect of guduchi extracts was comparable or better than doxorubicin treatment. The results demonstrate that guduchi killed the cells very effectively *in vitro* and deserves attention as an antineoplastic agent³⁸. Administration of

T. cordifolia stem methanolic extract to mice (200 mg/kg, i.p, daily for 5 days) increased the total white blood cell count significantly ($P < 0.001$). It also increased bone marrow cellularity ($18.16 \times 10^6/\text{femur}$) and α -esterase positive cells (1423/4000 cells) in bone marrow indicating increased maturation of stem cells. *Tinospora* extract reduced solid tumor growth and synergistically acted with cyclophosphamide in reducing (83%) the animal tumors. *T. cordifolia* extracts have been shown to up regulate the anti tumor activity of macrophages (TAM). Evidence has shown that an alcoholic extract of *T. cordifolia* (ALTC) enhances the differentiation of TAM to dendritic cells (DC) in response to granulocyte/macrophage-colony-stimulating factor, interleukin-4, and tumour necrosis factor 7¹²⁰. Cyclophosphamide (CP) an anti-cancer drug has been reported to reduce the GSH content in both bladder and liver and lowered levels of cytokines Interferon- γ and IL-2 an increased levels of pro-inflammatory cytokine TNF- α . This effect could be reversed on *T. cordifolia* treatment indicating the role of *T. cordifolia* in overcoming CP induced toxicities in cancer treatment¹²¹.

OTHER ACTIONS

The usefulness of *T. cordifolia* as a cognitive enhancer was substantiated by its potent *in vitro* acetylcholinesterase inhibitory activity. Methanolic and successive water extracts have been investigated, whereby, methanolic extract was found to be more active than water extract¹⁰². The efficacy of the extract in patients of allergic rhinitis was assessed in a randomized double blind placebo controlled trial which demonstrated a significant decrease in all the symptoms of allergic rhinitis⁹⁸. Methanolic (70%) stem extract has significant male antifertility activity. In male rats, it has no effect on body weight loss but decreases the weight of testes, epididymis, seminal vesicle and ventral prostate in a significant manner. It reduces the sperm motility as well as sperm density significantly, which result in reduction of male fertility by 100%⁵³. *T. cordifolia* can be exploited for the protection against gamma irradiation in mice. This property can be

exploited to human applications for radioprotective manifestation. A preparation of *T. cordifolia* administered to mice at a dose of 200 mg/kg and an aqueous extract at 5 mg/kg for 1 hour before whole body gamma irradiation, exhibits a significant protective effect. The direct and indirect antioxidant actions of *T. cordifolia* acts in corroboration to manifest the overall radioprotective effects^{100,122}.

ANTI-ARTHRITIC AND ANTI-OSTEOPOROTIC EFFECTS

Single or synergistic formulations of *T. cordifolia* with *Zingiber officinale* has been used in rheumatoid arthritis treatment in traditional medicine¹²³. *T. cordifolia* have been reported to affect the proliferation, differentiation and mineralization of bone like matrix on osteoblast model systems *in vitro* and hence finds potential application as an anti-osteoporotic agent. Alcoholic extract of *T. cordifolia* have been shown to stimulate the growth of osteoblasts, increasing the differentiation of cells into the osteoblastic lineage and also increasing the mineralization of bone like matrix. Beta-Ecdysone (Ecd) from *T. cordifolia* extracts have been reported to induce a significant increase in the thickness of joint cartilage, induce the osteogenic differentiation in mouse mesenchymal stem cells and to relieve osteoporosis in osteoporotic animal models¹²⁴.

ANTI-HIV EFFECTS

T. cordifolia has been shown to demonstrate a decrease in the recurrent resistance of HIV virus thus improving the therapeutic outcome¹²⁵. Anti-HIV effects of *T. cordifolia* was revealed by the reduction in eosinophil count, stimulation of B lymphocytes, macrophages and polymorphonuclear leucocytes and hemoglobin percentage thus, revealing its promising role of application in management of the disease¹²⁶.

NUTRIENT-NUTRIENT INTERACTIONS

Bacopa monnieri (Brahmi) and *Evolvulus alsinoides* (Shankhapushpi) are two ayurvedic herbs used for the purpose of cognitive

enhancement. The combination of these two herbs along with *T. cardifolia* appears to be most effective remedy against amnesia which suggests synergism¹²⁷. Ginger is a Spice that is

also traditionally used as an ayurvedic medicine and it is used alongside *T. cordifolia* for the treatment of rheumatism^{123,128}.

REFERENCES

- Sharma AC, Shanker LK, Tyagi, Singh M and CV Rao, Herbal medicine for market potential in India: An overview. African Journal of Plant Science, 1: 26-36 (2008).
- Sujata G, Dastidar A, Manna K, Asok K, Kaushiki M, NK, Thakur RS, Puri HS and Akhtar H, Major Medicinal plants of India. Science Research Corporation Dacca, 1: 177 (2004).
- Panchabhai TS, Ambharkhane SV, Joshi AS, Samant BD and Rege NN, Protective effects of *Tinospora cordifolia*, *Phyllanthus emblica* and their combination against antitubercular drugs induced hepatic damage: an experimental study. Phytotherapy Research, 22(5): 646 – 650 (2008).
- Saha S and Ghosh S, *Tinospora cordifolia*: One plant, many roles. Ancient Science of Life, 31(4): 151–159 (2012).
- Jagruti P, Krishna K, Jigar B and Jagruti P, Biological and Medicinal properties, a review. The International Journal of Alternative Medicine, 6(2): 160-168 (2009).
- Edoga HO, Okwu DE and Mbaebie BO, Phytochemicals constituents of some Nigerian medicinal plant. African Journal of Biotechnology, 4(7): 685-688 (2005).
- Bennett RN, Mellon FA, Foidl N, Pratt JH, Dupont MS, Perkins L and Kroon P, Profiling glucosinolates and phenolics in vegetative and reproductive tissues of the multi-purpose trees *Moringa oleifera* L. (horseradish tree) and *Moringa stenopetala* L. Journal of Agricultural and Food Chemistry, 51: 3546–3553 (2003).
- Adhvaryu MR, Reddy MN and Vakharia BC, Prevention of hepatotoxicity due to anti tuberculosis treatment: A novel integrative approach. World Journal of Gastroenterology, 14(30): 4753-4762 (2008).
- Mary NK, Babu BH and Padikkala J, Antiatherogenic effect of Caps HT2, a herbal Ayurvedic medicine formulation. International Journal of Phytotherapy and Phytopharmacology, 45: 457-565 (2003).
- Kaneria M, Kanani B and Chanda S, Assessment of effect of hydroalcoholic and decoction methods on extraction of antioxidants from selected Indian medicinal plants. Asian Pacific Journal of Tropical Biomedicine, 2(3): 195 – 20 (2012).
- Odukoya, Olukemi A, Ilori, Olukemi, Sofidiya, Oluwatoyin M, Aniunoh O, Austin, Lawal B, Mansurat and Tade I Olufunmilola, Antioxidant activity of Nigerian dietary spices. Electronic Journal of Environmental, Agricultural and Food Chemistry, 6: 1086-1093 (2005).
- Sudhakaran DS, Srirekha P, Devasree LD, Prem Singh S and Michael RD, Immunostimulatory effect of *Tinospora cardifolia* Miers leaf extract in *Oreochromis mossambicus*. Indian Journal of Experimental Biology, 44: 726–32 (2006).
- Sinha K, Mishra NP, Singh J and Khanuja SPS, *Tinospora cardifolia* (Guduchi), a reservoir plant for therapeutic applications: A review. Indian Journal of Traditional Knowledge, 3(3): 257-270 (2004).
- Ganesh CJ and Shaival KR, Evaluation of the Antineoplastic activity of Guduchi (*Tinospora cordifolia*) in Ehrlich Ascites Carcinoma bearing Mice. Biological and Pharmaceutical Bulletin, 29(3): 460-466 (2006).
- Purandare H and Supe A, Immunomodulatory role of *Tinospora cardifolia* as an adjuvant in surgical treatment of diabetic foot ulcers: a prospective randomized controlled study. Indian Journal of Medical Sciences, 61 (6): 347-55 (2007).

16. Sausa SM, Silva PS, Campos JMS and Viccini LF, Cytotoxic and genotoxic effects of two medicinal species of Verbenaceae. *Caryologia*, 62(4): 326-333 (2009).
17. Deole S Y, Sulakshan SC, Ashok BK, Ravishankar B, Thakar AB and Chandola HM, Evaluation of anti-depressant and anxiolytic activity of *Rasayana Ghana* Tablet (A compound Ayurvedic formulation) in albino mice. *Ayurveda*, 32(3): 375–379 (2013).
18. Choudhry N, Singh S, Badruzzaman MS and Khatoon S, Impact of Seasons and Dioccy on Therapeutic Phytoconstituents of *Tinospora cordifolia*, a Rasayana Drug. *BioMed Research International*, 2014: 1-11 (2014).
19. Sriramaneni RN, Omar AZ, Ibrahim SM, Amirin S and Zaini MA, Vasorelaxant effect of diterpenoid lactones from andrographis paniculata chloroform extract on rat aortic rings. *Pharmacognosy Research*, 2: 242–6 (2010).
20. Yang JH, Kondratyuk TP, Marler LE, Qiu X, Choi Y and Cao H, Isolation and evaluation of kaempferol glycosides from the fern *neochlopteris palmatopedata*. *Phytochemistry*, 71: 641–7 (2010).
21. Dhanasekaran M, Baskar AA, Ignacimuthu S, Agastian P and Duraipandiyan V, Chemopreventive potential of Epoxy clerodane diterpene from *Tinospora cardifolia* against diethylnitrosamine-induced hepatocellular carcinoma. *Invest New Drugs*, 27(4): 347-355 (2009).
22. Zhao TF, Wang X, Rimando AM and Che C, Folkloric medicinal plants: *Tinospora sagittata* var. *cravaniensis* and *Mahonia bealei*. *Planta Medicine*, 57: 505 (2008).
23. Kohno H, Maeda M, Tanino M, Tsukio Y, Ueda N and Wada KA, Bitter diterpenoid furanolactone columbin from *Calumbae Radix* inhibits azoxymethane-induced rat colon carcinogenesis. *Cancer Letter*, 183: 131–9 (2007).
24. Nadig PD, Revankar RR, Dethle SM, Narayanswamy SB and Aliyar MA, Effect of *Tinospora cardifolia* on experimental diabetic neuropathy. *Indian Journal of Pharmacology*, 44(5):580-3 (2012).
25. Singh A, Jain D, Upadhyay MK, Khandelwal N and Verma HN, Green Synthesis of Silver Nanoparticles using *Argemone mexicana* Leaf Extract and Evaluation of their Antimicrobial Activities. *Digest Journal of Nanomaterials and Biostructures's*, 5(2): 483-489 (2013).
26. Chen S, Wu K and Knox R, Structure-function studies of DT-diaphorase (NQO1) and NRH: Quinone oxidoreductase (NQO2). *Free Radical Biology and Medicine*, 20(29); 276–84 (2001).
27. Yang S, Evens AM, Prachands, Singh AT, Bhalla S, Devid K et al., Diterpenoid lactone and rographolide, the active component of andrographis paniculata. *Clinical Cancer Research*, 16: 4755-4768 (2010).
28. Singh RP, Banerjee S, Kumar PV, Raveesha KA and Rao AR, *Tinospora cardifolia* induces enzymes of carcinogen/drug metabolism and antioxidant system, and inhibits lipid peroxidation in mice. *Phytomedicine*, 13: 74–84 (2012).
29. Baldwin AS, Control of oncogenesis and cancer therapy resistance by the transcription factor NF-kappaB. *Journal of Clinical Investigation*, 107: 241–6 (2001).
30. Kim SK, Kim HJ, Choi SE, Park KH, Choi HK and Lee MW, Anti-oxidative and inhibitory activities on nitric oxide (NO) and prostaglandin E2 (COX-2) production of flavonoids from seeds of *Prunus tomentosa* Thunberg. *Archives of Pharmacology Research*, 31: 424–428 (2008).
31. Khatoon S, Singh H and Goel HK, Use of HPTLC to establish the chemotype of parasitic plant, *Dendrophthoe Falcata* etting, growing on different substrates. *Journal of planar chromatography*, 24(1): 60-65 (2011).
32. Sinha K, Mishra NP, Singh J and Khanuja SPS, *Tinospora cardifolia*(Guduchi), a reservoir plant for therapeutic applications: A review. *Indian Journal of Traditional Knowledge*, 3(3): 257-270 (2003).
33. Upadhyay AK, Kumar K, Kumar A and Mishra HS, *Tinospora cardifolia* (Willd.) Hook and Thoms. (Guduchi)-validation of

- the Ayurvedic pharmacology through experimental and clinical studies. International Journal of Ayurveda Research, 1: 112–21 (2010).
34. Swain SS, Rout KK and Chand PK, Production of triterpenoid anti-cancer compound taraxerol in Agrobacterium-transformed root cultures of butterfly pea (*Clitoria ternatea* L.). Applied Biochemistry and Biotechnology, 168(3):487-503 (2012).
 35. Patel SS, Shah RS and Goyal RK, Antihyperglycemic, antihyperlipidemic and antioxidant effects of Dihar, a polyherbal ayurvedic formulation in streptozotocin induced diabetic rats. Indian journal of experimental biology, 47: 564–70 (2009).
 36. Patel J, Kumar GS, Qureshi MS and Jena PK, Antihelmintic activity of ethanolic extract of whole plant of *Eupatorium odoratum*. International Journal of Phytomedicine, 2: 127-132 (2011).
 37. Gupta R and Sharma V, Ameliorative effects of *Tinospora cardifolia* root extract on histopathological and biochemical changes induced by aflatoxin-b (1) in mice kidney. International Journal of Toxicology, 18: 94–8 (2011).
 38. Jagetia GC and Rao SK, Evaluation of Cytotoxic Effects of Dichloromethane Extract of Guduchi (*Tinospora cardifolia* Miers ex Hook F & THOMS) on Cultured HeLa Cell. Evidence Based Complementary and Alternative Medicine, 3(2): 267-72 (2006).
 39. Ghosh AK, Chapsal BD, Weber IT and Mitsuya H, Design of HIV protease inhibitors targeting protein backbone: An effective strategy for combating drug resistance. Accounts of Chemical Research, 41: 78–86 (2008).
 40. Mukherjee R, De UK and Ram GC, Evaluation of mammary gland immunity and therapeutic potential of *Tinospora cardifolia* against bovine subclinical mastitis. Tropical Animal Health and Production, 42: 645-651 (2010).
 41. Jain SK, Dictionary of folk medicine and ethanobotany, 2nd Edn, Vol 21, Deep publications, New Delhi, India: 179-180, (2010).
 42. Mittal J, Sharma MM and Batra A, *Tinospora cordifolia*: a multipurpose medicinal plant- A review. Journal of Medicinal Plants Studies, 2: 32-47 (2014).
 43. Perkovic V, Lv J, Xu D, Ma X, Johnson D.W and Woodward M, Corticosteroid therapy in IgA nephropathy. Journal of the American Society of Nephrology, 23: 1108–16 (2012).
 44. McKeown E, Bykerk VP, De Leon F, Bonner A, Thorne C and Hitchon CA, Quality assurance study of the use of preventative therapies in glucocorticoid-induced osteoporosis in early inflammatory arthritis: Results from the catch cohort. Rheumatology (Oxford), 51: 1662–9 (2012).
 45. Sundarraj S, Thangam R, Sreevani V, Kaveri K, Gunasekaran P and Achiraman S, Y-Sitosterol from *acacia nilotica* L induces G2/M cell cycle arrest and apopyosis through c-Myc suppression in MCF-7 and A549 cells. Journal of Ethnopharmacology, 141: 803-809 (2012).
 46. Sharma V and Pandey D, Beneficial effects of *Tinospora cardifolia* on blood profiles in male mice exposed to lead. International Journal of Toxicology, 17: 8–11 (2010).
 47. Khan MI, Harsha PS, Giridhar P and Ravishankar GA, Pigment identification, antioxidant activity, and nutrient composition of *Tinospora cardifolia* (willd.) Miers ex Hook and Thoms fruit. International Journal of Food Science and Nutrition, 62: 239–49 (2011).
 48. Hazlewood CF, A view of the significance and understanding of the physical properties of cell-associated water. In Cell Associated Water, 165–259 (2001).
 49. Chougale AD, Ghadyale VA, Panaskar SN and Arvindekar AU. Alpha glucosidase inhibition by stem extract of *Tinospora cordifolia*. Journal of Enzyme Inhibition and Medicinal Chemistry, 24: 998–1001 (2010).
 50. Bramley PM, Regulation of carotenoid formation during tomato fruit ripening and development. Journal of Experimental Botany, 53: 2107–2113 (2002).
 51. Mojab F, Kamalinejad M, Ghaderi N and Vanidipour HR, Phytochemicals screening

- of some Iranian plants. Iranian Journal of Pharmaceutical Sciences, 3: 77-88 (2003).
52. Kumar S, Verma NS, Pande D and Srivastava PS, *In vitro* regeneration and screening of berberine in *Tinospora cordifolia*. Journal of Medicinal and Aromatic Plant Science, 22: 61 (2000).
 53. Rout KK, Pradhan S and Mishra SK, Estimation of berberine in ayurvedic formulations containing *Berberis aristata*. Journal of the Association of Official Agricultural Chemists, 91: 1149-1153 (2008).
 54. Jagetia GC and Rao SK, Evaluation of the antineoplastic activity of guduchi (*Tinospora cordifolia*) in Ehrlich ascites carcinoma bearing mice. Biological A and Pharmaceutical bulletin, 29(3): 460-6 (2006).
 55. Patel SR, Goyal RK and Shah DS, Studies on the pharmacological effects of *Tinospora cordifolia*. Indian Journal of Medical Research, 13(2): 46-57 (2010).
 56. Gupta RS and Sharma A, Antifertility affect of *Tinospora cardifolia* (Willd.) stem extract in male rats. Indian Journal of Experimental Biology. 41(8): 885-89 (2003).
 57. Patel MB and Mishra S, Hypoglycemic activity of alkaloidal fraction of *Tinospora cordifolia*. Phytomedicine, 18: 1045–52 (2011).
 58. Sarma DNK, Padma P and Khosa RL, Constituents of *Tinospora cardifolia* root. Fitoterapia, 69: 541-2 (1998).
 59. Zhao TF, Wang X, Rimando AM and Che C, phytochemistry of *Tinospora cardifolia*. Medicine, 57: 505 (2010).
 60. Lin CN, Huang AM, Lin KW, Hour TC, Ko HH, Yang SC and Pu YS, Xanthine oxidase inhibitory terpenoids of *Amentotaxus formosana* protect cisplatin-induced cell death by reducing reactive oxygen species (ROS) in normal human urothelial and bladder cancer cells, Phytochemistry, 71: 17-18 (2010).
 61. Khaleque A, Miah MAW, Huq MS and Abdul BK, *Tinospora cardifolia* IV Isolation of heptacosanol, b sitosterol and three other compounds tinosporidine, cordifol and cordifolone. Pakistan Journal of Scientific and Industrial Research, 14: 481-3 (2000).
 62. Lv J, Xu D, Perkovic V, Ma X, Johnson DW and Woodward M, Corticosteroid therapy in IgA nephropathy. Journal of the American Society of Nephrology, 23: 1108-16 (2012).
 63. Shaw CA, Ly PT and Singh S, Novel environmental toxins: Steryl glycosides as a potential etiological factor for age related neurodegenerative diseases, Journal of Neuroscience, 85: 231-237 (2007).
 64. Amarowicz R, Karamac M and Shahidi F, Antioxidant activity of phenolic fractions of lentil (*Lens culinaris*). Journal of Food Lipids, 10(1): 1–10 (2003).
 65. Serafini M, Bellocco R, Wolk A and Ekstrom A. M, Total antioxidant potential of fruit and vegetables and risk of gastric cancer. Gastroenterology, 123: 985–991 (2002).
 66. Carlsen MH, Halvorsen BL, Holte K, Bohn SK, Dragland S, Sampson L, et al., The total antioxidant content of more than 3100 foods, beverages, spices, herbs and supplements used worldwide. Journal of Nutrition, 9(3): 1-11 (2010).
 67. Matsui T, Ueda T, Oki T, Sugita K, Terahara N and Matsumoto K, Alpha-Glucosidase inhibitory action of natural acylated anthocyanins 2 alpha-Glucosidase inhibition by isolated acylated anthocyanins. Journal of Agricultural and Food Chemistry, 49(4): 1952-6 (2001).
 68. Madhujith and Shahidi, Optimization of the extraction of antioxidative constituents of six barley cultivars and their antioxidant properties. Journal of Agricultural and Food Chemistry, 54: 8048–8057 (2006).
 69. Ndakidemi PA and Dakora FD, Legume seed flavonoids and nitrogenous metabolites as signals and protectants in early seedling development. Functional Plant Biology, 30: 729-745 (2003).
 70. Qian W, Ge S and Hong DY, Genetic variation within and among populations of a wild rice *Oryza granulata* from China detected by RAPD and ISSR markers. Theoretical and Applied Genetics, 102 (2001).

71. Bafna PA and Balaraman R, Anti-ulcer and anti-oxidant activity of pepticare, a herbomineral formulation. *Phytomedicine*, 12: 264–70 (2005).
72. Kim SC, Lee C and Santos GA, Genetic analysis and conservation of the endangered Canary Island woody sow-thistle, *Sonchus gandogerii* (Asteraceae). *International Journal of Plant Research*, 118: 147–153 (2005).
73. Wang XM, Hou XQ, Zhang YQ, Yang R, Feng SF, Yan L and Ren Y, Genetic Diversity of the Endemic and Medicinally Important Plant *Rheum officinale* as Revealed by Inter-Simple Sequence Repeat (ISSR) Markers. *International Journal of Molecular Sciences*, 13: 3900-3915 (2012).
74. Sebastian VA, D'Cruz L, Subramanian RB and Braganza VJ, Assessment of genetic diversity within and among populations of *Tylophora rotundifolia* using RAPD markers. *Gene conserve*, 9: 94-117 (2010).
75. Warude D, Chavan P, Joshi K and Patwardhan B, DNA isolation from fresh and dry plant samples with high acidic tissue extracts. *Plant Molecular Biology Reporter*, 21: 467–467 (2003).
76. Ikbal, Boora KS and Dhillon RS, Evaluation of genetic diversity in *Jatropha curcus* using RAPD markers. *Indian Journal of Biotechnology*, 9: 2–57 (2010).
77. Samy RP, Pushparaj PN and Gopalakrishnakone P, A compilation of bioactive compounds from Ayurveda. *Biotransformation*, 3: 100–110 (2008).
78. Liu CW, Bi ZM, Zhu YF and Li P, Simultaneous determination of four kinds of bioactive components in *Radix Scrophulariae* by HPLC. *Journal of China Pharmaceutical*, 42(21): 1614–1616 (2007).
79. Zhang YB, Ngan FN, Wang ZT, Ng TB, Butt PP and Shaw PC, Random primed polymerase chain reaction differentiates *Codonopsis pilosula* from different localities. *Planta Medica*, 65(2): 157–160 (2008).
80. Shinde VM and Dhalwal K, DNA Fingerprinting of *Tinospora cardifolia* using RAPD analysis. *Journal of Global Pharma Technology*, 56: 751-767 (2008).
81. Khan S, Mirza KJ, Tayyab and Abdin MZ, RAPD profile for authentication of medicinal plant *Glycirriza glabra* L. *Internet Journal of Food Safety*, 11: 24-28 (2009).
82. Shinde VM, Dhalwal K, Mahadik KR, Joshi KS and Patwardhan BK, RAPD Analysis for Determination of Components in Herbal Medicine. *Evidence Based Complementary and Alternative Medicine*, 4(1): 165-190 (2001).
83. Rout GR, Identification of *Tinospora cardifolia* (Willd.) Miers ex Hook F & Thomas using RAPD markers. *Z Naturforsch C*, 61:118–22 (2006).
84. Palombo EA and Semple SJ, Antibacterial activity of traditional Australian medicinal plants. *Journal of Ethnopharmacology*, 77: 151-157 (2001).
85. Ahmad SM, Hoot SB, Qazi PH and Verma V, Phylogenetic patterns and genetic diversity of Indian *Tinospora* species based on chloroplast sequence data and cytochrome P450 polymorphisms. *Plant Systematics and Evolution*, 56: 751-767 (2009).
86. Khandelwal N, Singh A, Jain D, Upadhyay MK and Verma HN, Green Synthesis of Silver Nanoparticles using *Argemone mexicana* Leaf Extract and Evaluation of their Antimicrobial Activities. *International Journal of Nanomaterials and Biostructures*, 5(2): 483-489 (2006).
87. Ishnava K and Mohan JSS, Assessment of Genetic Diversity in Medicinal Climber of *Tinospora cordifolia*(Willd.) Miers (Menispermaceae) from Gujarat, India. *Asian Journal of Biotechnology*, 1(3): 93-103 (2009).
88. Alfred M, Traditional use of medicinal plants in south-central Zimbabwe: review and perspectives. *Journal of Ethnobiology and Ethnomedicine*, 9: 31 (2013).
89. Deshpande RS, Neelakanta NT and Hegde N, Cultivation of Medicinal Crops & Aromatic Crops as a Mean of Diversification in Agriculture. *Research Report: IX/ADRT/115*. (2006).
90. Patel N and Krishnamurthy SR, Indian *Tinospora* species: natural immunomodulators and therapeutic agents.

- International Journal of Pharmaceutical Biological and Chemical Sciences, 2 (2): 01-09 (2013).
91. Jeychandran RT, Xavier TF and Anand SP, Antimicrobial activity of stem extracts of *Tinospora cardifolia* (Willd.) Mierr ex Hook. F. and Thoms. Ancient science of life, 23(1): 40-43 (2003).
 92. Dahanukar SA and Rege NN, Quantitation of microbicidal activity of mononuclear phagocytes: an in vitro technique. Postgraduate Medical Journal, 39(1): 22-25 (2006).
 93. Singh SS, Pandey SC, Srivastava S, Gupta VS, Patro B and Ghosh AC, Chemistry and medicinal properties of *Tinospora cardifolia* (Guduchi). Indian Journal of Pharmacology, 35: 83-91 (2003).
 94. Anonymous, Wealth of India: A dictionary of Indian Raw Materials and Industrial Products, 1st Edn, Vol 10, New Delhi: CSIR: 2: 251, (2003).
 95. Thakur RS, Puri HS and Hussain A, Major Medicinal plants of India. Science Research Dacca, 1:177 (2000).
 96. Cosa P, Vlietinck AJ, Berghe DV and Maes L, Anti-infective potential of natural products, How to develop a stronger in vitro 'proof-of-concept'. The Journal of Ethnopharmacology, 106: 290-302 (2006).
 97. Sasidharan S, Chen Y, Saravanan D and Sundram KM, Extraction, Isolation and Characterization of Bioactive Compounds from Plants Extracts. African Journal of Traditional, Complementary and Alternative medicines, 8(1): 1-10 (2011).
 98. Badar VA, Thawani VR, Wakode PT, Shrivastava MP, Gharpure KJ, Hingorani LL and Khiyani, Efficacy of *Tinospora cardifolia* in allergic rhinitis. Journal of Ethnopharmacology, 96(3): 445-449 (2005).
 99. Choudhary BG, Patel NM, Shah PB and Modi KP, Development and validation of a HPTLC method for the simultaneous estimation of atorvastatin calcium and ezetimibe. Indian Journal of Pharmaceutical Sciences, 68:793 (2014).
 100. Pahadiya S and Sharma J, Alteration of lethal effects of gamma rays in Swiss albino mice by *Tinospora cordifolia*. Phytotherapy Research, 17(5): 552-54 (2003).
 101. Sivakumar V and Rajan MS, Standardization & Characterization of *Tinospora cardifolia* (Willd.) Miers ex Hook.F. & Thoms Plant stem extract in different solvent fractions. Asian Journal of Biochemical and Pharmaceutical Research, 4 (1): 56-90 (2011).
 102. Vinutha B, Prashanth D, Salma K, Sreeja SL, Pratiti D, Padmaja R, Radhika S, A A, Venkateshwarlu K, Screening of selected Indian medicinal plants for acetylcholinesterase inhibitory activity. Journal of Ethnopharmacology, 109(2):359-63 (2007).
 103. Alam P, Ali M, Singh R, Madhurima, Ahmad S and Shakeel, A Validated HPLC Method for Estimation of Cordifolioside A in *Tinospora cordifolia*, Miers and Marketed Formulations. Journal of Chromatographic Science, 34: 47 (2009).
 104. Desai VR, Ramkrishnan R, Chintalwar GJ and Sainis KB, G1-4A-an immunomodulatory polysaccharide from *Tinospora cordifolia*, modulates macrophage responses and protects mice against lipopolysaccharide induced endotoxic shock. International Immunopharmacology, 7: 1375-1386 (2007).
 105. Nair PK, Rodriguez S, Ramachandran R, Alamo A, Melnick SJ and Escalon E, Immune stimulating properties of a novel polysaccharide from the medicinal plant *Tinospora cordifolia*. International Immunopharmacology, 4: 1645-59 (2004).
 106. Leyon PV and Kuttan G, Inhibitory effect of a polysaccharide from *Tinospora cardifolia* on experimental metastasis. Journal of Ethnopharmacology, 90 (2): 233-37 (2004).
 107. Subramanian M, Chintalwar GJ and Chattopadhyay S, Antioxidant properties of a *Tinospora cardifolia* polysaccharide against iron-mediated lipid damage and

- gamma-ray induced protein damage. Redox Report, 7(3): 137-43 (2002).
108. Singh N, Singh SM, Prakash and Singh G, Restoration of thymic homeostasis in a tumor-bearing host by in vivo administration of medicinal herb *Tinospora cordifolia*. Immunopharmacology Immunotoxicology, 27(4): 585-99 (2005).
 109. Rao PR, Kumar VK, Viswanath RK and Subbaraju GV, Cardioprotective activity of alcoholic extract of *Tinospora cardifolia* in ischemia-reperfusion induced myocardial infarction in rats. Journal Biological and Pharmaceutical Bulletin, 28(12): 2319-22 (2005).
 110. Rathi SS, Grover JK and Vats V, Amelioration of experimental diabetic neuropathy and gastropathy in rats following oral administration of plant (*Eugenia jambolana*, *Mucuna pruriens* and *Tinospora cordifolia*) extracts. Indian Journal of Experimental Biology, 40(3): 273-76 (2002).
 111. Grover JK, Vats V, Rathi SS and Dawar R, Traditional Indian anti-diabetic plants attenuate progression of renal damage in streptozotocin induced diabetic mice. Journal of Ethnopharmacology, 76(3):233-38 (2001).
 112. Grover JK, Vats Vand and Rathi SS, Anti-hyperglycemic effect of *Eugenia jambolana* and *Tinospora cardifolia* in experimental diabetes and their effects on key metabolic enzymes involved in carbohydrate metabolism. Journal of Ethnopharmacology, 73(3): 461-70 (2000).
 113. Sangeetha MK, Raghavendran, Balaji HR, Gayathri V and Vasanthi HR, *Tinospora cardifolia* attenuates oxidative stress and distorted carbohydrate metabolism in experimentally induced type 2 diabetes in rats. Journal of Natural Medicines, 65: 544–550 (2011).
 114. Patel J, Kumar GS, Qureshi MS and Jena PK, Antihelmintic activity of ethanolic extract of whole plant of *Eupatorium odoratum*. International Journal of Phytomedicine, 2: 127-132 (2010).
 115. Shivananjappa MM and Muralidhara, Abrogation of maternal and fetal oxidative stress in the streptozotocin-induced diabetic rat by dietary supplements of *Tinospora cordifolia*. Nutrition, 28: 581–7 (2012).
 116. Manjrekar PN, Jolly CI and Narayanan S, Comparative studies of Immunomodulatory activity of *Tinospora cardifolia* and *T. sinensis*. Fitoterapia, 71: 254 (2000).
 117. Subramanian M, Chintalwar GJ and Chattopadhyay S, Antioxidant properties of a *Tinospora cardifolia* polysaccharide against iron-mediated lipid damage and gamma-ray induced protein damage. Redox Report, 7:137–43 2002.
 118. Gacche RN and Dhole NA, Antioxidant and possible anti-inflammatory potential of selected medicinal plants prescribed in the Indian traditional system of medicine. Journal of Pharmaceutical Biology, 44: 383-395 (2006).
 119. Baliga MS and Jagetia GC, The evaluation of nitric oxide scavenging activity of certain Indian medicinal plants *in vitro*: A preliminary study. Journal of Medicinal Food, 7: 343–8 (2006).
 120. Singh N, Singh SM and Shrivastava P, Effect of *Tinospora cardifolia* on the antitumor activity of tumor-associated macrophages-derived dendritic cells. Immunopharmacology and Immunotoxicology, 27(1):1-14 (2005).
 121. Hamsa TP and Kuttan G, *Tinospora cardifolia* ameliorates urotoxic effect of cyclophosphamide by modulating GSH and cytokine levels. Experimental and Toxicologic Pathology, 64: 307–14 (2012).
 122. Goel HC, Kumar IP and Rana S, Free radical scavenging and metal chelation by *Tinospora cordifolia*, a possible role in radioprotection, Indian Journal of Experimental Biology. 40(6): 727-34 (2002).
 123. Chopra A, Saluja M, Tillu G, Venugopalan A, Narsimulu G and Handa R, Comparable efficacy of standardized Ayurveda formulation and hydroxychloroquine sulfate (HCQS) in the treatment of rheumatoid arthritis (RA): a randomized investigator-

- blind controlled study. *Clinical Rheumatology*, 31(2): 259-69 (2012).
124. Kapur P, Pereira BM, Wuttke W and Jarry H. Androgenic action of *Tinospora cardifolia* ethanolic extract in prostate cancer cell line LNCaP. *Phytomedicine*, 16 (6-7): 679-82 (2010).
125. Kalikar MV, Thawani VR, Varadpande UK, Sontakke SD, Singh RP and Khiyani RK Immunomodulatory effect of *Tinospora cardifolia* extract in human immunodeficiency virus positive patients. *Indian Journal of Pharmacology*, 40(3): 107-10 (2008).
126. Akhtar S, Use of *Tinospora cardifolia* in HIV infection. *Indian Journal of Pharmacology*, 42 (1): 57 (2010).
127. Gupta A, Raj H, Karchuli MS and Upmanyu N, Comparative evaluation of ethanolic extracts of *Bacopa monnieri*, *Evolvulus alsinoides*, *Tinospora cardifolia* and their combinations on cognitive functions in rats. *Current Aging Science*, 6(3): 239-43 (2013).
128. Lekurwale PS, Pandey K and Yadaiah P, Management of Amavata with 'Amrita Ghrita': A clinical study. *Ayurveda*, 31(4): 430-5 (2010).